

REFERENCES

- Abrass, C.K., K.M. Nies, J.S.Louie, W.A.Border, and R.J. Glassock. 1980. Correlation and predictive accuracy of circulating immune complexes with disease activity in patients with systemic lupus erythematosus. Arthritis. Rheum. 23 : 273-282.
- Alcocer - Varela, J. and D. Alarcon - Segovia. 1982. decreased production of and response to interleukin-2 by cultured lymphocytes from patients with systemic lupus erythematosus. J.Clin.Invest. 69 : 1388-1392.
- Alcocer - Varela, J., A. Laffon, and D. Alarcon - Segovia. 1984. Defective monocyte production of, and T lymphocyte response to, interleukin-1 in the peripheral blood of patients with systemic lupus erythematosus. Clin.Exp.Immunol. 55 : 125-132.
- Alvarez, J.M., A. Silva, and M. O. Delandazuri. 1979. Human T cell growth factor : I. optimal conditions for its production. J. Immunol. 123 : 977-983.
- Arbeit, R.D., P.L.Leary, and M.J.Levin. 1982. Gamma interferon production by combinations of human peripheral blood lymphocytes, monocytes, and cultured macrophages. Infect. Immun. 35 : 383-390.
- Ashman, R.F. 1984. Lymphocyte activation : in Fundamental Immunology. W.E. Paul (ed). New York Raven Press. P. 267.

- Ayanlar-Batuman, O., J. Shevitz, U.C. Traut, S. Murphy, and D. Sejewski. 1987. Lymphocyte interleukin 2 production and responsiveness are altered in patients with primary myelodysplastic syndrome. Blood. 70 : 494-500.
- Bach, F.H., C. Grillot - Courvalin, and O.J. Kuperman. 1977. Antigenic requirements for triggering of cytotoxic T lymphocytes. Immunol. Rev. 35 : 76-96.
- Bakke, A.C., P.A. Kirkland, R.C. Kitridou, F.P. Quismorio, Jr, T. Rea, G.R. Ehresmann, and D.A. Horwitz. 1983. T lymphocyte subsets in systemic lupus erythematosus : correlations with corticosteroid therapy and disease activity. Arthritis. Rheum. 26 : 745-750.
- Balow, J.E. and A.S. Rosenthal. 1973. Glucocorticoid suppression of macrophage migration inhibitory factor. J. Exp. Med. 137 : 1031 - 1041.
- Barada, F.A., Jr, B.S. Andrews, J.S. Davis, and R.P. Taylor. 1981. Antibodies to Sm in patients with systemic lupus erythematosus : correlation of Sm antibody titers with disease activity and other laboratory parameters. Arthritis. Rheum. 24 : 1236-1244.
- Bast, R.C., Jr, E.L. Reinherz, C. Maver, P. Lavin, and S.F. Schlossman. 1983. Contrasting effects of cyclophosphamide and prednisolone on the phenotype of human peripheral blood leukocytes. Clin. Immunol. Immunopathol. 28 : 101-104.

- Bidulo, L., R.J.DeHoratius, C.Monihan, and A.T.Levinson. 1985. Restoration of the immunoglobulin secreting cells response to mononuclear cells to pokeweed mitogen by interleukin 1 and interleukin 2. Arthritis. Rheum. 28 : 571-575.
- Bishop, C.R., J.W.Athens, D.R.Boggs, H.R.Warner, G.E.Cartwright, and M. M. Wintrobe. 1968. Leukokinetic studies : XIII. a non-steady - state kinetic evaluation of the mechanism of cortisone - induced granulocytosis. J.Clin.Invest. 47 : 249-260.
- Blaese, R.M., J.Grayson, and A.D.Steinberg. 1980. Elevated immunoglobulin secreting cells in the blood of patients with active systemic lupus erythematosus : correlation of laboratory and clinical assessment of disease activity. Am.J.Med. 69 : 345-350.
- Bobrove, A.M. and P.Miller. 1977. Depressed in vitro B-lymphocyte differentiation in systemic lupus erythematosus. Arthritis. Rheum. 20 : 1326-1333.
- Boyum, A. 1968. Isolation of mononuclear cells and granulocytes from human blood. Scan.J.Clin.Lab.Invest. 21 (supple.97) : 77.
- Burns, G.F., C.L.Librach, I.M.Frazer, I.J.Kronborg, and I.R.Mackay. 1982. Spontaneous reverse hemolytic plaque formation. III. monocyte - mediated suppression of elevated plaque formation in autoimmune disease. Clin.Immunol.Immunopathol. 24 : 386-395.

- Campen, D.H., D.A.Horwitz, F.P.Quismorio, Jr, G.R.Ehresmann, and W.J.Martin. 1988. Serum levels of interleukin-2 receptor and activity of rheumatic diseases characterized by immune system activation. Arthritis. Rheum. 31 : 1358-1363.
- Cantor, H. and E.Simpson. 1975. Regulation of the immune response by subclasses T lymphocytes : I. interaction between pre-killer T cells and regulatory T cells obtained from peripheral lymphoid tissues of mice. Eur.J.Immunol. 5 : 330-336.
- Cantrell, D.A., and K.A.Smith. 1984. The interleukin-2 T-cell system : a new cell growth model. Science. 224 : 1312-1316.
- Cerottini, J.C. and K.T.Brunner. 1974. Cell-mediated cytotoxicity, allograft rejection, and tumor immunity. Adv.Immunol. 18 : 67-132.
- Charpentier, B., C.Carnaud, and J.F.Bach. 1979. Selective depression of the xenogeneic cell - mediated lympholysis in systemic lupus erythematosus. J.Clin.Invest. 64:351-360.
- Chopra, R.K., N.J.Holbrook, D.C.powers, M.T.McCoy, W.H.Adler, and J.E.Nagel. 1989. Interleukin 2, interleukin 2 receptor, and interferon- γ synthesis and mRNA expression in phorbol myristate acetate and calcium ionophore A 23187 - stimulated T cells from elderly humans. Clin.Immuno.Immunopatho. 53 : 297-308.
- Cochrane, C.E. and D. Koffler. 1973. Immune complex disease in experimental animals and man. Adv. Immunol. 16 : 185-221.

- Cooper, D.A., V.Petts, E.Luckhurst, and J.B.Ziegler. 1977. The effect of acute and prolonged administration of prednisolone and ACTH on lymphocyte subpopulations. Clin. Exp. Immunol. 28 : 467-473.
- Crabtree, G.R., S.Gillis, K.A.Smith, and A. Munck. 1979. Glucocorticoid and immune response. Arthritis. Rheum. 22 : 1246-1256.
- Cruickshank, B. 1987. The basic pattern of tissue damage and pathology of systemic lupus erythematosus : in Dubois' Lupus Erythematosus 3rd. Wallace. D.J. and E.L.Dubois (eds). Lea and Febiger publishing. Philadelphia. p. 53.
- Dale, D.C., A.S.Fauci, D. Guerry, IV, and S.M.Woff. 1975. Comparison of agents producing a neutrophilic leukocytosis in man : hydrocortisone, prednisone, endotoxin, and etiocholanolone. J. Clin. Invest. 56 : 808-813.
- Dauphinée, M.J., S.B.Kipper, D.Wofsy, and N. Talal. 1981. Interleukin 2 deficiency is a common feature of autoimmune mice. J.Immunol. 127 : 2483-2487.
- Davidson, W.F. and C.R.Parish. 1975. A procedure for removing red cells and dead cells from lymphoid cell suspensions. J.Immunol.Methods. 7 : 291-300.
- Decker, J.L., A.D. Steinberg, J.L.Reinertsen, P.H.Plotz, J.E.Balow, and J.H.Klippel. 1979. Systemic lupus erythematosus : evolving concepts.Ann.Intern.Med. 91:589-604.

- DeHoratius, R.J. 1982. Lymphocyte subsets in patient with systemic lupus erythematosus. Arthritis.Rheum. 25 : 828-832.
- Delfraissy, J.F., P.Segond, P.Galanand, C.Wallon, P.Massias, and J.Dormant. 1980. Depressed primary in vitro antibody response in untreated systemic lupus erythematosus : T helper cell defect and lack of defective suppressor cell function. J.Clin.Invest. 66 : 141-148.
- Depper, J.M., W.J.Leonard, M.Krönke, P.D.Noguchi, R.E.Cunningham, T.A.Waldmann, and W.C.Greene. 1984. Regulation of interleukin 2 receptor expression : effects of phorbol diester, phospholipase c, and reexposure to lectin or antigen. J.Immunol. 133 : 3054-3061.
- Dixon, F.J. 1985. Murine lupus : a model for human autoimmunity. Arthritis. Rheum. 28 : 1,081-1,088.
- Djeu, J.Y., J.A.Heinbaugh, H.T.Holden, and R.B.Herberman. 1979. Augmentation of mouse natural killer cells activity by interferon and interferon induces. J.Immunol. 122 : 175-181.
- Djeu, J.Y., N.Stocks, K.Zoon, G.J.Stanton, T.Timonen, and R.B.Herberman. 1982. Positive self-regulation of cytotoxicity in human natural killer cells by production of interferon upon exposure to influenza and herpes virus. J.Exp.Med. 156 : 1222-1234.

- Dos Reis, G.A. and E.M.Shevach. 1982. Effect of cyclosporin A on T cell function in vitro : the mechanism of suppression of T cell proliferation depends on the nature of the T cell stimulus as well as the differentiation state of the responding T cell. J.Immunol. 129 : 2360-2367.
- Draeger, A.M., A.J.G. Swaak, H.G.Van Den Brink, and L.A.Aarden. 1986. T cell function in systemic lupus erythematosus : normal production of and responsiveness to interleukin 2. Clin. Exp. Immunol. 64 : 80-87.
- Drutz, D.J. and J.R.Graybill. 1987. Infectious disease : in Basic and Clinical Immunology 6th. Stites.D.P., J.D.Stobo, and J.V.Wells (eds). Prentice-Hall publishing. East Norwald, Connecticut. P.534.
- Dutton, R.W., R.Falkoff, J.A.Hirst, M.Hoffman, J.W.Kappler, J.R. Keetman, J.R.Lesley, and P.Vann.1971. Is there evidence for a non-antigen specific diffusible chemical mediator from the thymus-derived cell in the initiation of the immune response? Prog.Immunol. 1 : 355-359.
- Dutton, R.W. 1975. Suppressor T cells. Transplant. Rev. 26 : 39-55.
- Elkon, K.B., M.J.Walport, R.I.Rynes. C.M.Black, J.R.Batchelor, and G.R.V. Hughes. 1983. Circulating C_{1q} binding immune complexes in relatives of patients systemic lupus erythematosus. Arthritis. Rheum. 26 : 921-927.

- Epstein, L.B., M.J.Cline, and T.C.Merigan. 1971. The interaction of human macrophages and lymphocytes in the phytohemagglutinin - stimulated production of interferon. J.Clin.Invest. 50 : 744-753.
- Erad, F., P.Corthesy, M.Nabholz, J.W.Lowenthal, P.Zaech, P.Plaetinck, and H.R.MacDonald. 1985. Interleukin 2 is both necessary and sufficient for the growth and differentiation of lectin stimulated cytotoxic T lymphocyte precursors. J.Immunol. 134 : 1644-1652.
- Falk, W., D.N.Mannel, and W. Droge. 1983. Activation of cytotoxic T lymphocytes requires at least two spleen cell-derived helper factors besides interleukin 2. J.Immunol. 130 : 2214-2218.
- Falk, W., D. N. Mannel, B. Katzer, B. Kaltman, P. H. Kramer, T. Diamanstein, and W. Droge. 1985. Induction of IL-2 receptor expression and cytotoxicity of thymocytes by stimulation with TCF 1. J.Immunol. 135 : 1160-1164.
- Farrar, J.J., P.L.Simon, W.J.Koopman, and J. Fuller - Bonar. 1978. Biochemical relationship of thymocyte mitogenic factor and factors enhancing humoral and cell-mediated immune responses. J.Immunol. 121 : 1353-1360.
- Fathman, C.G. and F.W.Fitch. 1984. Long-term culture of immunocompetent cells : in Fundamental Immunology. W.E.Paul(ed). New York, Raven Press. P.781.

- Fauci, A.S. and D.C.Dale. 1975. The effect of hydrocortisone on the kinetics of normal human lymphocytes. Blood. 46 : 235-243.
- Fauci, A.S., A.D.Steinberg, B.F.Haynes, and G.Whalen. 1978. Immunoregulatory aberrations in systemic lupus erythematosus. J.Immunol. 121 : 1473-1479.
- Fauci, A.S. and H.M.Moutsopoulos. 1981. Polyclonally triggered B cells in the peripheral blood and bone marrow of normal individuals and in patients with systemic lupus erythematosus and primary Sjögren's syndrome. Arthritis.Rheum. 24 : 577-584.
- Finke, J.H., J.Scott, S.Gillis, and M.L.Hilfiker. 1983. Generation of alloreactive cytotoxic T lymphocytes : evidence for a differentiation factor distinct from IL-2. J.Immunol. 130 : 763-767.
- Fournier, C. and J.Charreire. 1978. Activation of a human T cell subpopulation bearing receptors for autologous erythrocytes by concanavalin A. J.Immunol. 121 : 771-776.
- Friou, G.J. 1957. Clinical application of lupus serum nucleoprotein reaction using fluorescent antibody technique. J.Clin. Invest. 36 : 890.
- Fujii, M., K.Sugamura, K.Sano, M.Nakai, K.Sugita, and Y.Hinuma. 1986. High - affinity receptor - mediated internalization and degradation of interleukin 2 in human T cells. J.Exp.Med. 163 : 550-562.

- Garman, R.D. and D.P.Fan. 1983. Characterization of helper factor distinct from interleukin 2 necessary for the generation of allospecific cytolytic T lymphocytes. J.Immunol. 130 : 756-762.
- Gearing, A.J.H., A.P. Johnstone, and R.Thorpe. 1985. Production and assay of the interleukins. J.Immunol. Methods. 83 : 1-27.
- Gilliam, J.N., D.E.Cheatum, E.R.Hurd, P.Stastny, and M. Ziff. 1974. Immunoglobulin in clinically uninvolved skin in systemic lupus erythematosus : association with renal disease. J.Clin.Invest. 53 : 1434-1440.
- Gillis, S., M.M.Ferm, W.Ou, and K.A.Smith. 1978a. T cell growth factor : parameters of production and a quantitative microassay for activity. J.Immunol. 120 : 2027-2032.
- Gillis, S., P.E.Baker, F.W.Ruscetti, and K.A.Smith. 1978b. Long-term culture of human antigen-specific cytotoxic T-cell lines. J.Exp.Med. 148 : 1093-1098.
- Gillis, S., G.R.Crabtree, and K.A.Smith. 1979a. Glucocorticoid-induced inhibition of T cell growth factor production : I. the effect on mitogen-induced lymphocyte proliferation. J.Immunol. 123 : 1624-1631.
- Gillis, S., G.R.Crabtree, and K.A.Smith. 1979b. Glucocorticoid-induced inhibition of T cell growth factor production: II. the effect on the in vitro generation of cytolytic T cells. J.Immunol. 123 : 1632-1638.

- Gillis, S., K.A.Smith, and J.Watson. 1980. Biochemical and biological characterization of lymphocyte regulatory molecules : II. purification of a class of rat and human lymphokines. J.Immunol.124 : 1954-1962.
- Gillis, S., D.Y.Mochizuki, P.J.Conlon, S.H.Hefeneider, C.A.Ramthun, A.E.Gillis, M.B.Frank, C.S.Henney, and J.D.Watson. 1982. Molecular characterization of interleukin 2. Immunol.Rev. 63 : 166-208.
- Ginsberg, W.W., F.D. Finkelman, and P.E.Lipsky. 1979. Circulating and pokeweed mitogen - induced immunoglobulin - secreting cells in systemic lupus erythematosus. Clin. Exp. Immunol. 35 : 76-88.
- Glaser, M. and L.W.Law. 1978. T-T cell collaboration in rejection of a syngeneic SV 40-induced sarcoma in mice. Nature. 273 : 385-387.
- Gluckman, J.C. and P. Montambault. 1975. Spontaneous autorosette forming cells in man : a marker for a subset population of T lymphocytes? Clin.Exp.Immunol. 22 : 302-310.
- Gottlieb, A.B., R.G.Lahita, N.Chiorazzi, and H.G.Kunkel. 1979. Immune function in systemic lupus erythematosus : impairment of in vitro T-cell proliferation and in vivo antibody response to exogenous antigen. J. Clin. Invest. 63 : 885-892.

- Gray, P.W. and D.V.Goeddel. 1982. Structure of the human immune interferon gene. Nature. 298 : 859-863.
- Greene, W.C., R.J.Robb, P.B.Svetlik, C.M.Rusk, J.M.Depper, and W.J.Leonard. 1985. Stable expression of cDNA encoding the human interleukin 2 receptor in eukaryotic cells. J.Exp. Med. 162 : 363-368.
- Grimm, E.A., A.Mazumder, H.Z.Zhang, and S.A.Rosenberg. 1982. Lymphokine-activated killer cell phenomenon : lysis of natural killer-resistant fresh solid tumor cells by interleukin 2-activated autologous human peripheral blood lymphocytes. J.Exp.Med. 155 : 1823-1841.
- Grote, W., J.Klaar, P.F.Muhlradt, and D.A.Monner. 1987. Large scale production and purification of human IL-2 from buffy coat lymphocytes stimulated with 12-0-tetradecanoylphorbol 13-acetate and calcium ionophore A 23187. J.Immunol.Methods. 103 : 15-25.
- Hamaguchi, T., Y.Kotani, S.Imanaka, S.Morito, Y.Kawamura, and M. Shimizu. 1970. Lupus erythematosus and herpes zoster. MIE. Med.J. 19 : 189-192.
- Hamilton, M.E., D.E.Normansell, M.A.Garrett, and J.S.Davis IV. 1982. Mitogen stimulation of human lymphocytes : II. effect of exogenous DNA on lymphocyte function in systemic lupus erythematosus. Clin. Immunol. Immunopathol. 22 : 238-246.

- Handa, K., R.Suzuki, H.Matsui, Y.Shimizu, and K.Kumagai. 1983. Natural killer (NK) cells as a responder to interleukin 2 (IL2) : II. IL-2 induced interferon- γ production. J.Immunol. 130 : 988-992.
- Hardt, C., T. Diamanstein, and H.Wagner. 1985. Signal requirements for the in vitro differentiation of cytotoxic T lymphocytes (CTL) : distinct soluble mediators promote preactivation of CTL-precursors, clonal growth and differentiation into cytotoxic effector cells. Eur. J. Immunol. 15 : 472-478.
- Henney, C.S., K.Kuribayashi, D.E.Kern, and S.Gillis. 1981. Interleukin 2 augments natural killer cell activity. Nature. 291 : 335-338.
- Hirohata, S., L.S.Davis, and P.E.Lipsky. 1989. Role of IL-2 in the generation of CD4⁺ suppressors of human B cell responsiveness. J.Immunol. 142 : 3104-3112.
- Hoffman, T. 1980. Natural killer function in systemic lupus erythematosus. Arthritis. Rheum. 23 : 30-35.
- Horwitz, D.A. 1975. Mechanism producing decreased mitogenic reactivity in patients with systemic lupus erythematosus and other rheumatic diseases : in Mitogens in Immunobiology. Oppenheim. J.J. and D.L. Rosenstreich (eds). Academic Press, New York. P.625.

- Howard, M., J.Farrar, M.Hilfiker, B. Johnson, K.Takatsu, T.Hamaoka, and W.E.Paul.1982. Identification of T cell-derived B cell growth factor distinct from interleukin 2. J.Exp.Med. 155 : 914-923.
- Howard, M. and W.E. Paul. 1983. Regulation of B-cell growth and differentiation by soluble factors. Annu.Rev.Immunol. 1 : 307-334.
- Huang, Y.P., P.A.Miescher, and R.H.Zubler. 1986. The interleukin 2 secretion defect in vitro in systemic lupus erythematosus is reversible in rested cultured T cells. J.Immunol. 137 : 3515-3520.
- Hughes, G.R.V. 1982. Systemic lupus erythematosus in Lachmann : in Clinical Aspects in Immunology vol 2 (4th). P.J. and D.K. Peters (eds). Blackwell Scientific publication. Oxford. P. 1196.
- Inman, R.D. and N.K.Day. 1981. Immunologic and clinical aspects of immune complex disease. Am. J. Med. 70 : 1097-1106.
- Ishida, H., S.Kumagai, H.Umehara, H.Sano, Y.Tagaya, J.Yodoi, and H. Imura.1987. Impaired expression of high affinity interleukin 2 receptor on activated lymphocytes from patients with systemic lupus erythematosus. J.Immunol. 139 : 1070-1074.

- Kaplan, J. 1975. Human T lymphocytes form rosettes with autologous and allogeneic human red blood cells. Clin. Immunol. Immunopathol. 3 : 471 - 475.
- Kappler, J. and P. Marrack. 1986. Lymphokines : in Hand Book of Experimental Immunology in four volumes, vol 2 : cellular immunology. D.M.Weir(ed). Backwell scientific publications. P.59.1.
- Kasahara, T., J.J.Hooks, S.F.Dougherty, and J.J.Oppenheim. 1983. Interleukin 2-mediated immune interferon (IFN- γ) production by human T cells and T cell subsets. J.Immunol. 130 : 1784-1789.
- Katz, P., A.M.Zaytoun, and J.H.Lee, Jr. 1984. The effects of in vivo hydrocortisone on lymphocyte - mediated cytotoxicity. Arthritis. Rheum. 27 : 72-78.
- Kehrl, J.H., A.Muraguchi, J.L.Butler, R.J.M.Falkoff, and A.S.Fauci. 1984. Human B cell activation, proliferation and differentiation. Immunol.Rev. 78 : 75-96.
- Klein, J.R., D.H.Raulet, M.S.Pasternak, and M.J.Bevan. 1982. Cytotoxic T lymphocytes produced immune interferon in response to antigen or mitogen. J.Exp.Med. 155 : 1198-1203.
- Kunkel, M.G. 1983. The immunopathology of SLE : in the biology of Immunologic Disease. Diseon, F.J. and D.W.Fisker (eds). Sinauer associates publishing. Sauderland, Massachusetts. P. 247.

- Kuo, L.M., C.M.Rusk, and R.J.Robb. 1986. Structure - function relationships for the IL 2-receptor system : II. localization of an IL 2 binding site on high and low affinity receptors. J.Immunol. 137 : 1544-1551.
- Kurnick, J.T., K.O.Grönvik, A.K.Kimura, J.B.Lindblom, V.T.Skoog, O.Sjöberg, and H.Wigzell. 1979. Long term growth in vitro of human T cell blasts with maintenance of specific and function. J.Immunol. 122 : 1255-1260.
- Lahita, R.G., N.Chiorazzi, A.Gibofsky, R.J.Winchester, and H.G.Kunkel. 1983. Familial systemic lupus erythematosus in males. Arthritis. Rheum. 26 : 39-44.
- Lakhanpal, S. and B.S.Handwerger. 1986. Stimulation of resting normal human peripheral blood mononuclear cells by fetal calf sera. J.Immunol.Methods. 94 : 197-200.
- Lakhanpal, S. and B.S.Handwerger. 1987. Interleukin 2 receptors and responsiveness to recombinant human interleukin 2 in patients with systemic lupus erythematosus. Mayo.Clin.Proc. 62 : 3-7.
- Langford, M.P., G.J.Stanton, and H.M. Johnson. 1978. Biological effects of staphylococcal enterotoxin A on human peripheral lymphocytes. Infect. Immun. 22 : 62-68.

Leonard, W.J., J.M.Depper, T.Uchiyama, K.A.Smith, T.A.Waldmann, and W.C.Greene. 1982. A monoclonal antibody that appears to recognize the receptor for human T-cell growth factor : partial characterization of the receptor. Nature. 300 : 267-269.

Linker-Israeli, M., A.C.Bakke, R.C.Kitridou, S.Gendler, S.Gillis, and D.A.Horwitz. 1983. Defective production of interleukin-1 and interleukin-2 in patients with systemic lupus erythematosus (SLE). J.Immunol. 130 : 2651-2655.

Linker-Israeli, M., A.C.Bakke, F.P.Quismorio, Jr, and D.A. Horwitz. 1985. Correction of interleukin-2 production in patients with systemic lupus erythematosus by removal of spontaneously activated suppressor cells. J.Clin.Invest. 75 : 762-768.

Luger.T.A., J.S.Smolen, T.M.Chused, A.D.Steinberg, and J.J.Oppenheim. 1982. Human lymphocytes with either the OKT4 or OKT8 phenotype produce interleukin 2 in culture. J.Clin.Invest. 70 : 470-473.

Marcucci, F., M.Waller, H.Kirchner, and P.Kramer. 1981. Production of immune interferon by murine T-cell clones from long term cultures. Nature. 291 : 79-81.

McKenna, R.M., J.A.Wilkins, and R.J.Warrington. 1988. Lymphokine production in rheumatoid arthritis and systemic lupus erythematosus. J.Rheumatol. 15 : 1639-1642.

- Mendelsohn, J., M.M.Multer, and J.L.Bernheim. 1977. Inhibition of human lymphocytes stimulation by steroid hormones : cytokinetic mechanisms. Clin.Exp.Immunol. 27 : 127-134.
- Meuer, S.C., R.E.Hussey, A.C.Penta, K.A.Fitzgerald, B.M.Stadler, S.F.Schlossman, and E.L.Reinherz. 1982. Cellular origin of interleukin 2 (IL2) in man : evidence for stimulus-restricted IL-2 production by T4⁺ and T8⁺ T lymphocytes. J.Immunol. 129 : 1076-1079.
- Meuer, S.C., R.E.Hussey, D.A.Cantrell, J.C.Hodgdon, S.F.Schlossman, K.A.Smith, and E.L.Reinherz. 1984. Triggering of the T3-Ti antigen - receptor complex results in clonal T-cell proliferation through an interleukin 2 - dependent autocrine pathway. Proc.Natl.Acad.Sci.USA. 81 : 1509-1513.
- Miyasaka, N., T.Nakamura, I.J.Russel, and N.Talal. 1984. Interleukin 2 deficiencies in rheumatoid arthritis and systemic lupus erythematosus. Clin.Immunol.Immunopathol. 31 : 109-117.
- Moretta, A. 1985. Frequency a surface phenotype of human T lymphocytes producing interleukin-2:analysis by limiting dilution and cell cloning. Eur.J.Immunol. 15 : 148-155.
- Morgan, D.A., F.W.Ruscetti, and R.C.Gallo. 1976. Selective in vitro growth of T lymphocytes from normal human bone marrows. Science. 193 : 1007-1008.

- Morimoto, C., J.A.Distaso, Y.Borel, S.F.Schlossman, and E.L.Reinhery. 1982. Communicative interactions between subpopulation of human T lymphocytes required for generation of suppressor effector function in a primary antibody response. J.Immunol. 128 : 1645-1650.
- Morimoto, C., N.L.Letvin, J.A.Distaso, W.R.Aldrich, and S.F.Schlossman. 1985a. The isolation and characterization of the human suppressor - inducer T cell subset. J.Immunol. 134 : 1508-1515.
- Morimoto, C., N.L.Letvin, A.W.Boyd, M.Hagan, H.M.Brown, M.M.Kornacki, and S.F.Schlossman. 1985b. The isolation and characterization of the human helper inducer T cell subset. J.Immunol. 134 : 3762-3769.
- Morimoto, C., N.L.Letvin, J.A.Distaso, H.M.Brown, and S.F.Schlossman. 1986. The cellular basis for the induction of antigen specific T8 suppressor cells. Eur.J.Immunol. 16 : 198-204.
- Morimoto, C., A.D.Steinberg, N.L.Letvin, M.Hagan, T.Takeuchi, J.Daley, H.Levine, and S.F.Schlossman. 1987. A defect of immunoregulatory T cell subsets in systemic lupus erythematosus patients demonstrated with anti-2H4 antibody. J.Clin. Invest. 79 : 762-768.

- Mosmann, T.R., T.Yokota, R.Kastelein, S.M.Zurawski, N.Arai, and Y.Takebe. 1987. Species-specificity of T cell stimulating activities of IL-2 and BSF-1 (IL-4) : comparison of normal and recombinant, mouse and human IL-2 and BSF-1 (IL-4). J.Immunol. 138 : 1813-1816.
- Murakawa, Y., S.Takada, Y.Ueda, N.Suzuki, T.Hoshino, and T.Sakane. 1985. Characterization of T lymphocyte subpopulations responsible for deficient interleukin 2 activity in patients with systemic lupus erythematosus. J.Immunol. 134 : 187-195.
- Murakawa, Y. and T.Sakane. 1988. Deficient phytohemagglutinin-induced interleukin-2 activity in patients with inactive systemic lupus erythematosus is correctable by the addition of phorbil myristate acetate. Arthritis. Rheum. 31 : 826-833.
- Nakagawa, T., T.Hirano, N.Nakagawa, K.Yoshiyake, and T.Kishimoto. 1985. Effect of recombinant IL-2 and IFN- γ on proliferation and differentiation of human B cells. J.Immunol. 134 : 959-966.
- Nathan, I., J.E.Croopman, S.G.Quan, N.Bersch and D.W.Golde. 1981. Immune (gamma) interferon produced by a human T-lymphoblast cell line. Nature. 292 : 842-844.

- Nelson, D.L., L.A.Rubin, C.C.Kurman, M.E.Fritz, and B.Boutin. 1986. An analysis of the cellular requirements for the production of soluble interleukin-2 receptors in vitro. J.Clin.Immunol. 6 : 114-120.
- Nies, K.M., R.H.Stevens, and J.S.Louie. 1980. Normal T cell regulation of Ig G synthesis in systemic lupus erythematosus. J.Clin.Lab.Immunol. 4 : 69-75.
- Northoff, H., C.Carter, and J.J.Oppenheim. 1980. Inhibition of concanavalin A-induced human lymphocyte mitogenic factor (interleukin-2) production by suppressor T lymphocytes. J. Immunol. 125 : 1823-1828.
- O'Malley, J.A., A.Nussbaum-Blumenson, D.Sheedy, B.J.Grossmayer, and H.Ozer. 1982. Identification of the T cell subset that produce human γ -interferon. J.Immunol. 128 : 2522-2526.
- Ortaldo, J.R., A.T.Mason, J.P.Gerard, L.E.Henderson, W.Farrar, R.F.Hopkins III, R.B.Herberman, and H.Rabin. 1984. Effects of natural and recombinant IL-2 on regulation of IFN- γ production and natural killer activity : lack of involvement of the Tac antigen for these immunoregulatory effects. J.Immunol. 133 : 779-783.
- Palacios, B. and G. Möller. 1981. T cell growth factor abrogates concanavalin A induced suppressor cell function. J.Exp.Med. 153 : 1365-1365.

- Parrillo, J.E. and A.S.Fauci. 1979. Mechanism of glucocorticoid action on immune processes. Ann.Rev.Pharmacol.Toxicol. 19 : 179-201.
- Ralph, P., G.Jeong, K.Welte, R.Mertelsmann, H.Rabin, L.E.Henderson, L.M.Souza, T.C.Boone, and R.J.Robb. 1984. Stimulation of immunoglobulin secretion in human B lymphocytes as a direct effect of high concentration of IL-2. J.Immunol. 133 : 2442-2445.
- Raulet, D.H. and M.J.Bevan. 1982. A differentiation factor required for the expression of cytotoxic T-cell function. Nature. 296 : 754-757.
- Raziuddin, S., M.S.Nur, and A.A.Alwabel. 1989. Selective loss of the CD4⁺ inducer of suppressor T cell subsets (2H4⁺) in active systemic lupus erythematosus. J.Rheumatol. 16 : 1315-1319.
- Reinherz, E.L., P.C.Kung, G.Goldstein, and S.F.Schlossman. 1979. Separation of functional subsets of human T cells by a monoclonal antibody. Proc. Natl.Acad.Sci.USA. 76 : 4061-4065.
- Reske-Kunz, A.B., H.Osawa, O.Josimavic-Alasevic, E.Rude, and T.Diamantstein. 1987. Soluble interleukin 2 receptors are released by long term-cultured insulin-specific T cells transiently after contact with antigen. J.Immunol. 138 : 192-196.

- Robb, R.J., A. Munck, and K.A.Smith. 1981. T cell growth factor receptors : quantitation, specificity, and biological relevance. J.Exp.Med. 154 : 1455-1474.
- Robb, R.J. 1984. Interleukin 2 : the molecule and its function. Immunol. Today. 5 : 203-209.
- Robb, R.J., W.C.Greene, and C.M.Rusk. 1984. Low and high affinity receptors for interleukin 2 : implications for the level of Tac antigen. J.Exp.Med. 160 : 1126-1146.
- Robb, R.J. 1985. Human interleukin 2. Methods in Enzymo. 116 : 493-525.
- Robb, R.J. and W.C.Greene. 1987. Internalization of interleukin 2 is mediated by the β chain of the high - affinity interleukin 2 receptor. J.Exp.Med. 165 : 1201-1206.
- Robb, R.J. and R.M.Kuty. 1987. Structure-function relationships for the IL-2 receptor system : IV. analysis of the requence and ligand-binding properties of soluble Tac protein. J.Immunol. 139 : 855-862.
- Robb, R.J., C.M.Rusk, J.Yodoi, and W.C.Greene. 1987. Interleukin 2 binding molecule distinct from the Tac protein : analysis of its role in formation of high - affinity receptors. Proc. Natl.Acad.Sci.USA. 84 : 2002-2006.
- Roder, J.C. and T.Haliotis. 1980. Do NK cells play a role in anti-tumor surveillance. Immunol.Today. 1 : 96-100.

- Rosenthal, C. J. and E. C. Franklin. 1975. Depression of cellular - mediated immunity in systemic lupus erythematosus : relation to disease activity. Arthritis. Rheum. 18 : 207-217.
- Rubin, L.A., C.C.Kurman, M.E.Fritz, W.E.Biddison, B.Boutin, R. Yarchoan, and D.L.Nelson. 1985. Soluble interleukin 2 receptors are released from activated human lymphoid cells in vitro. J.Immunol. 135 : 3172-3177.
- Rubin, L.A., G.Jay, and D.L.Nelson. 1986. The released interleukin 2 receptor binds interleukin 2 efficiently. J.Immunol. 137 : 3841-3844.
- Sagawa, A. and N.I.Abdou. 1978. Suppressor-cell dysfunction in systemic lupus erythematosus : cells involved and in vitro correction. J.Clin.Invest. 62 : 789-796.
- Saiki, O., Y.Saeki, and S.Kishimoto. 1985. Spontaneous immunoglobulin A secretion and lack of mitogen-responsive B cells in systemic lupus erythematosus. J.Clin.Invest. 76 : 1865-1870.
- Sakane, T., M.Honda, Y.Taniguchi, and H.Kotani. 1981. Separation of concanavalin A -induced human suppressor and helper T cells by the autologous erythrocyte rosette technique. J.Clin. Invest. 68 : 447-453.
- Salmon, M., G.D.Kitas, J.S.Hill Gaston, and P.A.Bacon. 1988. Interleukin-2 production and response by helper T-cell subsets in man. Immunol. 65 : 81-85.

- Sandilands, G.P., K.G.Gray, A.E.Cooney, J.D.browning, and J.R. Anderson. 1974. Autorosette formation by human thymocytes and lymphocytes. Lancet. 1 : 27-28.
- Sato, K., N.Miyasaka, K.Yamaoka, M.Okuda, J.Yata, and K.Nishioka. 1987. Quantitative defect of CD4⁺ 2H4⁺ cells in systemic lupus erythematosus and sjögren's syndrome. Arthritis.Rheum. 30 : 1407-1411.
- Schimpl, A. and E.Wecker. 1972. Replacement of T-cell function by a T-cell product. Nature. 237 : 15-17.
- Schrader, J.W. and G.M.Edelman. 1976. Participation of the H-2 antigens of tumor cells in their lysis by syngeneic T cells. J.Exp.Med. 143 : 601-614.
- Schwartz, R.S. and S.K.Datta. 1989. Autoimmunity and autoimmune disease; in Fundamental Immunology 2nd. Paul.W.E. (ed). Raven Press. New York. P.819.
- Severinson, E., T.Naito, H.Tokumoto, D.Fukushima, A.Hirano, K.Hama, and T.Honjo. 1987. Interleukin-4 (Ig G1 induction factor) : a multifunctional lymphokine acting also on T cells. Eur.J. Immunol. 17 : 67-72.
- Shou, L., S.A.Schwartz, and R.A.Good. 1976. Suppressor cell activity after concanavalin A treatment of lymphocytes from normal donors. J.Exp.Med. 143 : 1100-1110.

- Sibbitt, W.L., Jr, C.Kenny, C.W.Spellman, K.D.Ley, and A.D.Bankhurst. 1984. Lymphokines in autoimmunity : relationship between interleukin-2 and interferon- γ production in systemic lupus erythematosus. Clin.Immunol.Immunopathol. 32 : 166-173.
- Siegel, J.P., M.Sharon, P.L.Smith, and W.J.Leonard. 1987. The IL-2 receptor β chain (p70) : role in mediating signals for LAK, NK, and proliferative activities. Science. 238 : 75-78.
- Sinigaglia, F., D.Scheidegger, K.Talmadge, and G.Garotta. 1985. A sensitive and quantitative microassay for the detection of mycoplasma contamination : inhibition of IL-2 dependent cell line proliferation. J.Immunol.Meth. 76 : 85-92.
- Slade, J.D. and B.Hepburn. 1983. Prednisone-induced alterations of circulating human lymphocyte subsets. J.Lab.Clin.Med. 101 : 479-487.
- Smith, K.A., S.Gillis, P.E.Baker, D.McKenzie, and F.W.Ruscetti. 1979. T-cell growth factor-mediated T-cell proliferation. Ann. NY.Acad.Sci. 322 : 423-432.
- Smith, K.A., P.G.Baker, S.Gillis, and F.W.Ruscetti. 1980. Functional and molecular characteristic of T-cell growth factor. Mol. Immunol. 17 : 579-589.
- Smith, K.A. and F.W.Ruscetti.1981. T-cell growth factor and the culture of cloned functional T cells. Adv.Immunol. 31 : 137-175.

- Smith, K.A. 1984. Interleukin 2. Ann. Rev. Immunol. 2:319-333.
- Smith, K.A. and D.A.Cantrell. 1985. Interleukin 2 regulates its own receptors. Proc.Natl.Acad.Sci.USA. 82 : 864-868.
- Smith, K.A. 1987. The two-chain structure of high-affinity IL-2 receptor. Immunol. Today. 8 : 11-13.
- Smith, K.A. 1988. The interleukin 2 receptor. Adv.Immunol. 42 : 165-179.
- Staples, P.J., D.N.Gerding, J.L.Decker, and R.S.Gordon. 1974. Incidence of infection in systemic lupus erythematosus. Arthritis. Rheum. 17:1-4.
- Stohl, W. and J.Z.Singer. 1987. Correlation between systemic lupus erythematosus and T4 epitope phenotype. Arthritis. Rheum. 30 : 1412-1415.
- Symington, F.W. and H.S.Teh. 1980. A two-signal mechanism for the induction of cytotoxic T lymphocytes. Scand. J. Immunol. 12 : 1-12.
- Takada, S., Y.Ueda, Y.Murakawa, N.Suzuki, and T. Sakane. 1983. Functional heterogeneities among concanavalin A-activated OKT4⁺ and OKT8⁺ cells by using autologous erythrocyte rosette technique. J.Clin.Invest. 72 : 2060-2071.
- Takeuchi, T., C.E.Rudd, S.F.Schlossman, and C.Morimoto. 1987. Induction of suppression following autologous mixed lymphocyte reaction : role of a novel 2H4 antigen. Eur.J. Immunol. 17 : 97-103.

- Tan, E.M. and H.G.Kunkel. 1966. Characteristics of a soluble nuclear antigen precipitating with sera of patients with systemic lupus erythematosus. J.Immunol. 96 : 464-471.
- Tan, E.M., A.L.Cohen, J.F.Fries, A.T.Masi, D.J.McShane, N.F. Rothfield, J.G.Schaller, N.Talal, and R.J.Winchester. 1982. The 1982 revised criteria for the classification of systemic lupus erythematosus. Arthritis. Rheum. 25 : 1271-1277.
- Tan, P., G.Pang, and J.D.Wilson. 1981. Immunoglobulin production in vitro by peripheral blood lymphocytes in systemic lupus erythematosus : helper T cell defect and B cell hyperactivity. Clin Exp. Immunol. 44 : 548-554.
- Tanaka, S., T.Matsuyama, A.D.Steinberg, S.F.Schlossman, and C.Morimoto. 1989. Antilymphocyte antibodies against CD4⁺ 2H4⁺ cell populations in patients with systemic lupus erythematosus. Arthritis. Rheum. 32 : 398-405.
- Taniguchi, T., H.Matsui, T.Fujita, C.Takaoka, N.Kashima, R.Yoshimoto, and J.Hamuro. 1983. Structure and expression of a cloned cDNA for human interleukin-2. Nature. 302 : 305-310.
- TenBerg, R.J., H.P.Sauerwein, S.L.Yong, and P.T.Schellekens. 1984. Administration of prednisolone in vivo affects the ratio of OKT4/OKT8 and the LDH-isoenzyme pattern of human T lymphocytes, Clin.Immunol.Immunopathol. 30 : 91-103.

- Teshigawara, K., H.M.Wang, K.Kato, and K.A.Smith. 1987. Interleukin 2 high-affinity receptor expression requires two distinct binding proteins. J.Exp.Med. 165 : 223-238.
- Theofilopoulos, A.N. and F.J.Dixon. 1979. The biology and detection of immune complexes. Adv. Immunol. 28 : 89-220.
- Theofilopoulos, A.N. and F.J.Dixon. 1985. Murine models of systemic lupus erythematosus. Adv.Immunol. 37 : 269-390.
- Timonen, T., J.R.Ortaldo, and R.B.Herberman. 1981. Characteristics of human large granular lymphocytes and relationship to natural killer and K cells. J.Exp.Med. 153 : 569-582.
- Timonen, T., J.R.Ortaldo, B.M.Stadler, G.D.Bonnard, S.Sharrow, and R.B.Herberman. 1982. Cultures of purified human natural killer cells : growth in the presence of IL-2. Cell.Immunol. 72 : 178-185.
- Tsokos, G.C. and J.E.Balow. 1983. Phenotypes of T lymphocytes in systemic lupus erythematosus : decreased cytotoxic/suppress or subpopulation is associated with deficient allogeneic cytotoxic responses rather than with concanavalin A-induced suppressor cells. Clin.Immunol.Immunopathol. 26 : 267-276.
- Tsokos, G.C., P.L.Smith, C.B.Christian, R.N.Lipnick, J.E.Balow, and J.Y.Djeu. 1985. Interleukin-2 restores the depressed allogeneic cell-mediated lympholysis and natural killer cell activity in patients with systemic lupus erythematosus. Clin.Immunol.Immunopathol. 34 : 379-386.

- Tsuda, M., R.W.Kozuaka, C.K.Goldman, and T.A.Waldman. 1986. Demonstration of a non-Tac peptide that binds interleukin 2; a potential participant in a multichain interleukin 2 receptor complex. Proc.Natl.Acad.Sci.USA. 83 : 9694-9698.
- Tsuda, M., R.W.Kozak, C.K.Glodman, and T.A.Waldmann. 1987a. Contribution of a p75 interleukin 2 binding peptide to a high - affinity interleukin 2 receptor complex. Proc.Natl. Acad.Sci.USA. 84 : 4215-4218.
- Tsuda, M., C.K.Goldman, K.F.Bongiovanni, W.C.Chan, E.F.Winton, M.Yagita, E.A.Grimm, and T.A.Waldman. 1987b. The P75 peptide is the receptor for interleukin 2 expressed on large granular lymphocytes and is responsible for the interleukin 2 activation of these cells. Proc.Natl.Acad.Sci.USA. 84 : 5394-5398.
- Utsinger, P.D. and W.J.Yount. 1977. Phytohemagglutinin response in systemic lupus erythematosus : reconstitution experiments using helper purified lymphocyte subpopulations and monocytes. J.Clin.Invest. 60 : 622-638.
- Vilcek, J., D.Henriksen-Destefano, D.Siegel, A.Klion, R.J.Robb, and J.Le. 1985. Regulation of IFN-gamma induction in human peripheral blood cells by exogenous and endogenously produced interleukin 2. J.Immunol. 135 : 1851-1856.

- Vohr, H.W. and T.Hunig. 1985. Induction of proliferative and cytotoxic responses in resting Lyt-2⁺ T cells with lectin and recombinant interleukin 2. Eur.J.Immunol. 15 : 332-337.
- Volk, H.D., J.Kopp, I.J.Körner, S.Jahn, R.Grunow, H.Barthelmes, and H.Fiebig. 1986. Correlation between the phenotype and the functional capacity of activated T cells in patients with active systemic lupus erythematosus. Scand.J.Immunol. 24 : 109-114.
- Wagner, H. and M.Röllinghoff. 1978. T-T-cell interactions during the in vitro cytotoxic allograft responses : I. soluble products from activated Ly1⁺ T cells trigger autonomously antigen-primed Ly23⁺ T cells to cell proliferation and cytolytic activity. J.Exp.Med. 148 : 1523-1538.
- Wagner, H., C.Hardt, B.T.Rouse, M.Rollinghoff, P.Scheurich, and K.Pfiffermaier. 1982. Dissection of the proliferative and differentiative signals controlling murine cytotoxic T lymphocytes responses. J.Exp.Med. 155:1876-1881.
- Wang, H.M. and K.A.Smith. 1988. The interleukin 2 receptor : functional consequences of its bimolecular structure. J.Exp.Med. 166 : 1055-1069.
- Watson, J. and D.Mochizuki. 1980. Interleukin 2 : a class of T cell growth factors. Immunol. Rev. 51 : 257-278.

- Webel, M.L. and R.E.Aritts, Jr. 1977. The effects of corticosteroid concentrations on lymphocyte blastogenesis. Cell.Immunol. 32 : 287-292.
- Weigent, D.A., G.J.Stanton, and H.M.Johnson. 1983. Interleukin 2 enhances natural killer cell activity through induction of gamma interferon. Infect.Immunol. 41 : 992-997.
- Weston, W.L., M.J.Mandel, and J.A.Yeckley. 1973. Mechanism of cortisol inhibition of adoptive transfer of tuberculin sensitivity. J.Lab.Clin.Med. 82 : 366-371.
- Wheelock, E.F. 1965. Interferon-like virus-inhibitor induced in human leukocytes by phytohemagglutinin. Science. 149 : 310-311.
- Wolf, R.E. and W.G.Brelsford. 1988. Soluble interleukin-2 receptors in systemic lupus erythematosus. Arthritis.Rheum. 31 : 729-735.
- Yip, Y.K., H.L.Pang, C.Urban, and J.Vilcek. 1981. Partial purification and characterization of human γ (immune) interferon. Proc.Natl.Acad.Sci.USA. 78 : 1601-1605.