

REFERENCES

- Ames , B.N., Durston, W.E., Yamasaki, E., and Lee, F.D.(1973)
 "Carcinogens are mutagens : a simple test system
 combining liver homogenates for activation and
 bacteria for detection." Proc Natl Acad Sci U.S.A.,
 70 : 2281-2285.
- Ames, B.N.(1989) "Mutagenesis and Carcinogenesis : Endogenous
 and exogenous factors." Environ Mol. Mutagen., 14 Suppl.
 16:66-77.
- Balbinder, E., Stick, S.M., and Sharma, O.K. (1983) " Complex
 effects of retinol on the metabolic activation of 2-
 aminofluorene." Environ. Mutagen., 5 : 665-678.
- Busk, L., and Ahlborg, U.G.(1980) "Retinol (vitamin A) as an
 inhibitor of the mutagenesis of aflatoxin B₁." Toxicol Lett., 6 : 243 - 249.
- Carmeron, E., Pauling, L. and Leibovitz, B. (1979) "Ascorbic
 acid and cancer : A review." Cancer Res., 39 : 663-
 681.
- Chaplin, M.F., and Kenedy, J.F., (1986) Carbohydrate analysis
 : a practical approach., Oxford, England.

Chopra, D.P., and Wilkoff L.J. (1975) "Inhibition and reversal of carcinogen-induced lesions in the mouse of prostate in vitro by all trans-retinoic acid." Pro Am Assoc Cancer Res., 16 : 35.

Chopra, D.P., and Wilkoff, L.J. (1977) "Reversal by vitamin A analogues (retinoids) of hyperplasia induced by N²-methyl- N²- nitroguandine in mouse prostate organ cultures." J Natl Cancer Inst., 58 : 922-930.

Czeczot, H., Tudek, B., Kusztelak, J., Szymczyk, T., Dobrowolska, B., Glinkowska, G., Malinowski, J., and Strzelecka, H., (1990) "Isolation and studies of the mutagenic activity in the Ames test of flavonoids naturally occurring in medical herbs." Mutation Res., 240 : 209-216.

Elias, P.M., and Williams, M.L. (1981) "Retinoids, cancer and skin." Arch. Dermatol., 117 : 160-181.

Fiddler, W., Pensabine, J.W., Piotrowski, E.G., Doerr, R.C., and Wasserman, A.E. (1978). "Use of sodium ascorbate or erythrobrate to inhibit formation of N-nitrosodimethylamine in frankfurters." J. Food Sci., 38 : 1084 (Abstract).

Garner, R.C., Miller, E.C., and Miller, J.A. (1972). "Liver microsomal metabolism of aflatoxin B₁ to a reactive derivative toxic to Salmonella typhimurium TA 1530." Cancer Res., 32:2058-2066.

- Goodman, D.S. (1979) "Vitamin A and retinoids : Recent advances, introduction , background and general overview." Fed. Proc., 38 : 2501 - 2503.
- Hill, D.L., and Shih, T.W. (1974) "Vitamin A compounds and analogs as inhibitors of mixed-function oxidases that metabolize carcinogenic polycyclic hydrocarbon and other compounds." Cancer Res., 34 : 564-570.
- Hodgson, E., and Guthrie, F.E. (1980) "Introduction to biochemical toxicology, New York:Elsevier, pp. 62-104.
- Kasper, C.B., and Henton, D. (1980) : Glucuronidation. In Jakoby, W.B. "Enzymatic basis of detoxication, vol.II" New York : Academic Press. pp 3-36.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L., and Randall, R.J. (1951) "Protein measurement with the folin phenol reagent" J.Biol.Chem., 193:265-275.
- MacGregor, J.T. (1984) "Genetic and carcinogenic effects of plant flavonoids : An overview." Adv. Exp. Med. Biol., 177 : 497-526
- Malaveille, C., Brun, G., Hantefeulle, A. and Bartsch, H. (1980) "Effect of glutathione and uridine 5'-diphosphoglucuronic acid on benzo (a) pyrene mutagenesis in the Salmonella/ microsome assay." Dev Toxicol Environ Sci, 8 : 175 - 180.

Maron, M.D., and Ames, B.N. (1983) Revised methods for the Salmonella mutagenicity test. Mutation Res., 113 : 173-215.

Marquardt, H., Ruffing, F. and Weisburger, J.H. (1977) "Mutagenic activity of nitrite treated foods: human stomach cancer may be relates to dietary factors." Science, 196: 1000-1001.

Matsushima, T., Sawamura, K., Hara, K., and Sugimura, T. (1976) A safe substitute for polychlorinated biphenyls as an inducer of metabolic activation system, in F.J. de Serres, JR. Fouts, JR. Fouts, JR. Bend and RM Philpot (Eds). In vitro metabolic activation in mutagenesis testing. Elsevier/North-Holland. Amsterdam, p 85-88.

Matsushima, T., Sugimura, T., Nagao, T., Shirai, A., and Sawamura, M. (1980) Factors modulating mutagenicity in microbial test, in K.H. Norpoth and RC. Garner (Eds) Short-term test system for detecting carcinogens. Springer, Berlin, p 273-285.

Meester, C.D. (1989) "Bacterial mutagenicity of heterocyclic amines found in heat-processed food." Mutation Res., 221: 235-262.

Merriman, R.L., and Bertram, J.S. (1979) "Reversible inhibition by retinoids of 3-methylcholanthrene-induced neoplastic transformation in C3H/10T 1/2 clone 8 cells." Cancer Res., 39 : 1661-1666.

Pariza, M.W., Loretz, L.J., Storkson, J.M., and Holland, N.C. (1983) "Mutagens and modulators of mutagenesis in fried ground beef." Cancer Res., 43:244-246.

Purintrapiban, J. (1989) "Study of mutagenic activity of some chilli-paste samples and of their ingredients." Thesis, Chiang Mai University, Thailand.

Qin, S., and Huang, C.C. (1985) "Effect of retinoids on carcinogen-induced mutagenesis in *Salmonella* tester strains." Mutation Res., 142:115-120.

Raina, V., and Gurtoo, H.L. (1985) "Effects of vitamins A, C, and E on aflatoxin B₁-induced mutagenesis in *Salmonella typhimurium* TA -98 and TA-100." Teratogenesis Carcinog Mutagen., 5:29-40.

Recio, L., and Hsie, A.W. (1984) "Glucuronide conjugation reduces the cytotoxicity but not the mutagenicity of benzo(a) pyrene in the CHO/HGPRT assay." Teratogenesis, Carcinogenesis, and Mutagenesis., 4:391-402.

Shephard, S.E., Schlatter, CH., and Lutz, W.K. (1987) "Assessment of the risk of formation of carcinogenic N-nitroso compounds from dietary precursors in the stomach." Food chem. toxicol., 25:91-108.

Slaga, T.J., and Braken, W.M. (1977) "The effect of antioxidants on skin tumor initiation and aryl hydrocarbon hydroxylase." Cancer Res., 37:1631-1635.

Stiltz, D.R., Starvic, B., Stapley, R., Kalssen, R., Bendall, R., and Krewski, D. (1984). "Mutagenicity screening of foods. II Result with fruits and vegetables." Environ. Mutagen., 6:343-354.

Sugimura, T. (1988) "Successful use of short-term tests for academic purposes: Their use in identification of new environmental carcinogens with possible risk for humans." Mutation Res., 205:33-39.

Tannenbaum, S.R., Weisman, M., and Fett, D. (1976) "The effect of nitrate intake on nitrite formation in human saliva." Fd. Cosmet. Toxicol., 14:549.

Todaro, G.J., Delacro, J.W., and Sporn, M.B. (1978) "Retinoids block phenotypic cell transformation produced by sarcoma growth factor." Nature., 276:272-274.

Vinitketkumnuen, U., Suttajit, M., and Matsushima, T. (1989)
 "Mutagenic activity of chilli-pastes and their
 ingredients." Environ. Mol. Mutagen., 14 suppl. 15:210.

Vinitketkumnuen, U. (1991) "Analytical studies of mutagens/
 carcinogens in common Thai foods." Ph.D. Dissertation,
 University of Tokyo, Japan.

Wakabayashi, K., Ochiai, M., Saito, H., Tsuda, M., Suwa, Y.,
 Nagao, M., and Sugimura, T. (1983) "Precence of 1-methyl
 -1,2,3,4-tetrahydro- β -carboline-3-carboxylic acid, a
 precursor of a mutagenic nitroso compound, in soy
 sauce." Proc Natl Acad Sci U.S.A., 80:2912-2916.

Waziers, I.D., and Decloitre, F. (1984) "Effect of glutathione
 and uridine-5'-diphosphoglucuronic acid on the
 mutagenicity of tryptophan pyrolysis products (Trp-P-1
 and Trp-P-2) by rat-liver and -intestine S9 fraction."
Mutation Res., 139:15-19.

Williams, G.M., Mori, H., and McQueen, C.A. (1989) "Structure-ac-
 tivity relationships in the rat hepatocyte DNA-repair
 test for 300 chemicals." Mutation Res., 221:263-286.