

## VII. REFERENCES

Abbas, A.K., Lichtman, A.H. and Pober, J.S., (1991). The major histocompatibility complex, p.108-110. *In* Abbas, A. K., Lichtman, A. H. and Pober, J. S. (Ed.), Cellular and molecular immunology. W.B.Saunders.

Abel, L., Demenais, F., Baule, M-S., Blanc, M., Muller, A., Raffoux, C., Millan, J., Bois, E., Babron, M-C. and Feingold, N., (1988). Genetic susceptibility to leprosy on a Caribbean island : linkage analysis with five markers. *Inter. J. Leprosy.*, 57: 465-471.

Altmann, D.M., Sansom, D. and Marsh, S.G.E., (1991). What is the basis for HLA-DQ associations with autoimmune disease?., 12: 267-270.

Amos, D.B., Ward, F.e., (1975). Immunogenetics of the HLA system. *Physiol. Rev.* 55:206.

Auffray, C., Lillie, J.W., Korman, A.J., Boss, J.M., Frechin, N., Guillemot, F., Cooper, J., Mulligan, R.C., and Strominger, J.L., (1987). Structure and expression of HLA-DQ $\alpha$  and -DX $\alpha$  genes: interallelic alternate splicing of the HLA-DQ $\alpha$  gene and functional splicing of the HLA-DQ $\alpha$  gene using a retroviral vector. *Immunogenetics.*, 26: 63-73.

Auffray, C., Lillie, J. W., Arnot, D., Grossberger, D., Kappes, D., Strominger, J. L., (1984). cDNA clone for the heavy chain of the human B cell alloantigen DC1: Strong sequence homology to the HLA-DR heavy chain. *Nature*, 308: 327-333.

Binford, C.H., Meyers, W.M., Walsh, G.P., (1982) Leprosy. *J. Amer.Med. Assoc.*, 247:2283-2292.

Blackwell, J.M., (1988). Bacterial infection *In* D.Wakelin, S.M. Blackwell (Ed), Genetics of resistance to bacterial and parasitic infections. Basingstoke.

Boehringer Mannheim Biochemica, (1993). The DIG system user's guide for filter hybridization. Germany.

Brown, J. H., Jardetzky, T., Saper, M. A., Samraoui, B., Bjorkman, P. J., Wiley, D. C., (1988). A hypothetical model of the foreign antigen binding site of class II histocompatibility molecules. *Nature*, 332: 845-850.

Brown J. H., Jardetzky, T. S., Gorga, J. C., Stern, L. J., Urban, R. G., Strominger, J. L., Wiley, D. C., (submitted). The three-dimensional structure of the human class II histocompatibility antigen HLA-DR1.

Bryceson, .A.D.M., Pfaltzraff, R.E., (1979). *Leprosy and edition* Edinburgh: Churchill Livingstone.

Bugawan, T. L., Chang, J. D., Klitz, W., and Erlich, H. A., (1994). PCR/Oligonucleotide probes typing of HLA class II alleles in a Filipino population reveals an unusual distribution of HLA haplotypes. *Am. J. Hum. Genet.*, 54: 331-340.

Cerna, M., Fernandez-Vina, M., Ivaskova, E., Stastny, P., (1992). Comparison of HLA class II alleles in Gypsy and Czech populations by DNA typing with oligonucleotide probes. *Tissue Antigens.*, 39: 111-116.

Chalkravartti, M.R., Vogel, F.A., (1973). A twin study of leprosy. *Top. Hum. Genet.*, 1:1-123.

Chan, S. H., Oon, B. B., Kamarudin, A., Wee, G. B., (1979). HLA and leprosy in Chinese. *Tissue Antigens.*, 13: 73-74.

Clara, G., Flores, J., Arevalo, N., Le Castro., Silva, A., Rodriguez, O., (1987). Tuberculoid leprosy in Mexicans is associated with HLA-DR3. *Lepr. Rev.* 58:401-406.

Dasgupta, A., Mehra, N.K., Ghei, S.K., Vaidya, M.C., (1975). Histocompatibility antigen (HL-A) in leprosy. *Tissue Antigens.*, 5: 85-87.

De Vries, R.R.P., Serjeantson, S. W., Layrisse, Z., (1984). Leprosy, p.362-367. In E. D.Albaert, M. P. Baur, W. R. Mayr (Eds), Histocompatibility testing. Berlin-Verlag.

De Vries, R.R.P., Van Eden, W., Van Rood, J.J., (1981). HLA-linked control of the course of *M. leprae* infection. *Lepr Rev.*, 52 (Suppl): 109-19.

De Vries, R.R.P., (1991). Genetic control of immunopathology induced by *Mycobacterium leprae*. *Amer. J. Trop. Med. Hyg.*, 44: 90-128.

De Vries, R.R.P., Ottenhoff, T.H.M., Van Schooten, W.C.A., (1988). Human leukocyte antigen (HLA) and Mycobacterial disease. Springer. *Semin. Immunopathol.*, 10: 305-318.

De Vries, R.R.P., Van Eden, W., Van Rood, J.J., (1981). HLA-linked control of the course of *M. leprae* infection. *Lepr Rev.*, 52(Suppl): 109-19.

De Vries, R.R., Mehra, N.K., Vaidya, M.C., Gupte, M.D., Meera, Khan, P., Van Rood, J.J., (1980). HLA-linked control of susceptibility to tuberculoid leprosy and association with HLA-DR types. *Tissue Antigens.*, 16: 294-301.

Djoulah, S., Sanchez-Mazas, A., Khalil, I., Benhamamouch, S., Degos, L., Deschamp, I., Hors, J., (1994). HLA-DRB1, DQA1, and DQB1 DNA polymorphisms in healthy Algerian and genetic relationships with other populations. *Tissue Antigens.*, 43: 102-109

Doherty, D.G., Vaughan, R.W., Donaldson, P.T., Mowat, A.P., (1992). HLA DQA, DQB, and DRB genotyping by oligonucleotide analysis: distribution of alleles and haplotypes in British caucasoids. *Human Immunol.*, 34: 53-63.

Escobar-Gutierrez, A., Gorodezky, C., Salazar-Mallen, M., (1973). Distribution of some of HL-A system leukocyte antigens in Mexicans. II. Studies in atopics and in lepers. *Vox Sang.*, 25: 151-155.

Fine, P.E.M., (1988). Implication of genetics for the epidemiology and control of leprosy. *Phil. Trans. R. Soc. Lond. B.*, 321:365-376.

Gao, X., Sun, An, J., Fernandez-Vina, M., Qou, J., Lin, L., Stastany, P., (1991). DNA typing for HLA-DR, and -DP alleles in a Chinese population using the polymerase chain reaction (PCR) and oligonucleotide probes. *Tissue Antigens.*, 38: 24-30.

Godal, T., Myrvang, B., Stanford, J-L., Samuel, D.R., (1974). Recent advances in the immunology of leprosy with special references to new approaches in immunoprophylaxis. *Bull. Inst. Pasteur.*, 72: 273.

Goding, J.W., Walker, I.D., (1980). Allelic forms of  $\beta 2$  microglobulin in the mouse. *Proc. Natl. Acad. Sci. U.S.A.* 77: 7395-7399.

Gorodezky, C., Flores, J., Arevalo, N., Castro, Le., Silva, A., Rodriguez, O., (1987). Tuberculoid leprosy in Mexicans is associated with HLA-DR3. *Lepr. Rev.* 58: 401-406.

Greiner, J., Schleiermacher, E., Smith, T., Lenhard, V., Vogel, F., (1978). The HLA-system and leprosy in Thailand. *Hum Genet.*, 42: 201-13.

Haldane, J.B.S., (1955). The estimation and significant of the logarithm of a ratio of frequencies. *Ann. Hum. Genet.*, 20: 309-311.

Hansen, G.A., (1874). Undersogelser angaaende spedalskhedens arsagar. *Norsk. May. Laegeridensk.*, 4: 1-88.

Hansen T.H., Carreno, B.M., and Sachs, D.H., (1992). The major histocompatibility complex, p578-675. *In* Paul, W. E. (Ed.), *Fundamental Immunology*. Raven press, New York.

Harris, H.W., Gill, T.J. III., (1986). Expression of class I transplantation antigens. *Transplantation.* 42:109-115.

Hasting, R. C. (Ed), (1985). Leprosy. Edinburgh: Churchill Livingstone.

Hill, A.V.S., Allsopp, E.M., Kwiatkowski, D., Anstey, N.M., Twumasi, P., Rowe, P.A., Bennett, S., Brewster, D., McMichael, A.J., and Greenwood, B.M., (1991). Common west African HLA antigens are associated with protection from severe malaria. *Nature.*, 352: 595-600.

Hopkin, R., Denny, O.E., (1929). Leprosy in the United State. *J. Amer. Med. Assoc.*, 92: 191-198.

Izumi, S., Sugiyama, K., Matsumoto, Y., Ohkawa, S., (1982). Analysis of the immunogenetic background of Japanese leprosy patients by the HLA-system. *Vox Sang.*, 42: 243-47.

Jonsson, A. -K., Hyldig-Nielsen, J. -J., Servenius, B., Larhammar, D., Anderson, G., Jorgensen, F., Peterson, P. A., Rask, L., (1987). Class I genes of the human major histocompatibility complex. *J. Biol. Chem.*, 262:8767-8777.

Kanjanahaleuthai, A., Smith, T., Sittisombut, N. *TaqI* restriction fragment length polymorphism of HLA-DQA1 and -DQA2 genes in northern Thailand (in press).

Kikuchi, I., Okawa, T., Hirayama, K., Sasazuki, T., (1986). An HLA-linked gene controls susceptibility to lepromatous leprosy through T cell recognition. *Lepr. Rev.*, 57, Suppl. 2: 139-142.

Kirchheimer, W.F. and Storis, E.E., (1971). *International Journal of Leprosy and other Mycobacterium disease*, 39: 693-702.

Korman, A.J., Boss, J.M., Spies, T., Sorentino, R., Okada, K., Strominger, J.L. (1985). Genetic complexity and expression of human class II histocompatibility antigen. *Immunol. Rev.*, 85: 45-86.

Kreisler, M., Arnaiz, A., Perez, B., Cruz, E.F., Bootello, A., (1974). HL-A antigens in leprosy. *Tissue Antigens.*, 4: 179-201.

Lango, A. and Lindblom, B., (1993). HLA DQA-DQB haplotypes in a swedish population. *Eur. J. Immunogenet.*, 20; 453-460.

Li, S.G., De Vries, R.R.P., (1989). HLA-DQ molecules may be products of an immune suppressive gene responsible for *Mycobacterium leprae*-specific nonresponsiveness. *Int. J. Lepr.*, 57: 554-556.

Long, E.D., (1989). Intracellular traffic and antigen processing. *Immunol. Today.*, 10: 232-234.

Mac Mahon, B. and Pugh, T. F., (1970). *Epidemiology: Principles and methods.* Little, Brown and Co., Boston.

Mach, B., Tiercy, J.-M., (1991). Genotyping of HLA Class II: from the bench to the bedside. *Hum. Immunol.*, 30: 278-284.

Marsh, S.G.E., Bodmer, J.G., (1991). HLA class II nucleotide sequence. *Hum Immunol.*, 31: 207.

Mat, M.C., Yazici, H., Ozbakir, F., Tuzun, Y., (1988). The HLA association of lepromatous leprosy and borderline lepromatous leprosy in Turkey. *Vox Sang.*, 27: 246-247.7

Mehra, N.K., Daskupta, A., Ghei, S.K., Nilakanta Rao, M.S., Vaidya, M.C., (1976). HLA antigens and leprosy. *Microbios. Lett.*, 3: 79-83.

Miettinen, O.S., (1974). Propotion of disease caused or prevented by a given exposure, trait or invention. *Am. J. Med.*, 52:1-8.

Miller, S.A., Dykes, D.D., Polesky, H.F., (1988). A simple salting-out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Research.*, 16: 1215.

Miyataka, K., Juju, T., Maeda, H., Nakajima, S., Kobayashi, S., (1981). Tuberculoid leprosy and HLA in Japanese. *Tissue Antigens.*, 18: 331-334.



Mohamed, A.P., Ramanujam, K., (1966). Leprosy in twin. *Int. J. Lepr.*, 34:405-407.

Nepom, G.T. and Erlich, H., (1991). MHC class II molecules and autoimmunity. *Annual review of immunology.*, 9: 493-526.

Ottenhoff, T.H., Gonzalez, N.M., de Vries, R.R., Convit, J., Van Rood, J.J., (1984). Association of HLA specificity LB-E12 (MB1, DC1, MT1) with lepromatous leprosy in a Venezuelan population. *Tissue Antigens.* 24: 25-29.

Ottenhoff, T.H., de Vries, R.R., (1987). HLA class II immune response and suppression genes in leprosy. *Int. J. Lepr.*, 55: 521-534.

Rani, R., Zaheer, S.A., Mukherjee, R., (1992). Do human leukocyte antigens have a role to play in differential manifestation of multibacillary leprosy; a study on multibacillary leprosy patients from north India. *Tissue Antigens.*, 40: 124-127.

Rani, R., Fernandez-Vina, M.A., Zaheer, S.A., Beena, K.R., Stastny, P., (1993). Study of HLA class II alleles by PCR oligotyping in leprosy patients from north India. *Tissue Antigens.*, 42: 133-137.

Rea, T. H., Levan, N. E., Terasaki, P. I., (1976). Histocompatibility antigens in patients with leprosy. *J Infect Dis.*, 134: 615-618.

Rea, T. H., Terasaki, P.I., (1980). HLA-DR antigens in tuberculoid and lepromatous leprosy. *Lepr. Rev.*, 51: 117-23.

Reference protocol, (1991). 11th International Histocompatibility Workshop, Yokohama Japan. November 6-13.

Reis, A.P., Maia, F., Reis, V. F., Andreade, I.M., Campos, A.A.B., (1974). HL-A antigens in leprosy. *lancet.*, 1384.

Ridley, D.S., Jopling, W.H., (1966). Classification of leprosy according to immunity : A five group system. *Inter. J. Leprosy.*, 34: 255-273.

Robinson, M. A., Kindt, T. J., (1989). Major histocompatibility complex antigens and genes, p.489-539. *In* Paul, W. E. (Ed.), *Fundamental Immunology*. Raven Press.

Ronninggen, K. S., Spurkland, A., Iwe, T., Vartdal, F., Thorsby, E., (1991). Distribution of HLA-DRB1, -DQA1 and DQB1 alleles and DQA1-DQB1 genotypes among Norwegian patients with insulin-dependent diabetes mellitus. *Tissue Antigens.*, 37: 105-111.

Sachs, D.H., (1984). The major histocompatibility complex, p.446-484. *In* W.E. Paul (ed), *Fundamental immunology*. Raven Press.

Salazar, M., Yunis, J.J., Delgado, M.B., Bing, D. and Yunis, E. ., (1992) HLA-DQB1 allele typing by a new PCR-RFLP method: Correlation with PCR-SSO method. *Tissue Antigen.*, 40: 116-123.

Sanger, F., Nicklen, S., and Coulson, A.R., (1977). DNA sequencing with chain terminating inhibitors. *Proc Natl Acad Sci USA.*, 74: 5463-5467.

Schauf, V., Ryan, S., Scollard, D., Jonasson, O., Brown, A., Nelson, K., Smith, T. and Vithayasai, V., (1985). Leprosy associated with HLA-DR2 and DQw1 in the population of northern Thailand. *Tissue Antigens*, 26:243-247.

Schurr, E., Morgan, K., Gres, P. and Skamene E., (1991). Genetic of leprosy. *Am. J. Trop. Med. Hyg.*, 44 Suppl: 4-11.

Skinsnes, O.K., and Chang, P.H.C., (1985). Understanding of leprosy in ancient China. *Int. J. Lepr.*, 53: 289-307.

Smith, G.S., Walford, R. L., Shepard, C.C., Payne, R., Prochazka, G.J., (1975). Histocompatibility antigens in leprosy. *Vox Sang.*, 28: 42-49.



Sveiguard, A., Hauge, M., Jersild, C., Platz, P., Ryder, L. P., Staub Nielsen, L., Thomsen, M., (1975). The HLA system. Karger, Basel.

Takata, H., Sada, M., Ozawa, S., Sekiguchi, S., (1978). HLA and mycobacterial infection: Increased frequency of B8 in Japanese leprosy. *Tissue Antigens.*, 11: 61-64.

Teodorica, L.B., Chang, J.D., William, K., and Erlich, A., (1994). PCR/oligonucleotide probe typing of HLA class II alleles in Filipino population reveals an unusual distribution of HLA haplotypes. *Am. Hum. Genet.*, 54: 331-340.

Thompson, M. W., McInnes, R. R., Willard, H. F., (1986). Genes in population, p. 143-165. In Thompson, M. W., McInnes, R. R., Willard, H. F. (Ed), *Genetics in Medicine*. W. B. Saunders Company.

Thorsby, E., Godal, T., Myrvang, B., (1973). HL-A antigens and susceptibility to disease. II. Leprosy. *Tissue Antigens.*, 3: 373-377.

Tiercy, J.-M., Grundschober, C., Jeannet, M. and Mach, B., (1993). A comprehensive HLA-DRB, -DQB, and -DPB oligotyping procedure by hybridization with sequence-specific oligonucleotide probes, p. 117-148. In Hui, K.M. and Bidwell, J.L. (Ed), *Handbook of HLA typing techniques*. CRC Press.

Tomolo, A., Murphy, O.J., Nguyen, Q., Voyta, J.C., Witney, F., Bronstein, I., (1992). Detection of DNA on membrane with alkaline phosphatase-labeled probes and chemiluminescent AMPPD substrate, p. 128-144. In Kricka, L. J. (Ed), *Nonisotopic DNA probe techniques*. Academic Press.

Trowsdale, J., Ragoussis, J., Campbell, R.D., (1991) Map of the human MHC. *Immunol Today.*, 12: 445-448.

Trowsdale, J., Young, J.A., Kelly, A.P., Austin, P.J., Carson, S., Meunier, H., So, A., Erlich, H.A., Spielman, R.S., Bodmer, J., et al., (1985). Structure, sequence and polymorphism in the HLA-D region. *Immunol. Rev.*, 85:5-43.

Van Eden, W., Gonzalez, N.M., de Vries, r.r., Convit, J., Van Rood, J.J., (1985). HLA-linked control of predisposition to lepromatous leprosy. *J. Infect. Dis.*, 151:9-14.

Van Eden, W., Mehra N.K., Vaidya, M.C., D' Amaro, J. Schreuder, G.M.Th, Van Rood, J.J., (1981). A population study in Maharashtra, India. *Tissue Antigens.*, 18: 189-93.

Vogel, F.. Motulsky., (1986). Formal genetics of man, p. 111-226. *In* Vogel, F.. Motulsky (Ed), *Human genetics*. Springer-Verlag Berlin Heidelberg New York, Tokyo.

Wang, F. Q., Semana, G., Fauchet, R., and Genetet, B., (1993). HLA-DR and -DQ genotyping by PCR-SSO in Shanghai Chinese. *Tissue Antigens.*, 41: 223-226.

WHO Nomenclature Committee., (1992). Nomenclature for factor of the HLA system. *Immunogenet.*, 36: 135.


Wood, W. I., Gitschier, J., Lasky, L. A., and Lawn, R. M., (1985). Base composition independent hybridization in tetramethylammonium chloride; A method for oligonucleotide screening of highly complex gene libraries. *Proc. Natl. Acad. Sci. USA.*, 82: 1585-1590.

Woof, B., (1955). On estimating the relation between blood group and disease. *Ann. Hum. Genet.*, 19:251-253.

World Health Organization, (1978). Nomenclature for factors of the HLA system. *Bull. WHO.*, 54: 46.

Yanagawa, T., Manglabruks, A., Chang, Y.-B., Okamoto, Y., Fisfalen, M.-E., Curran, P.G. and Degroot, L.J.,(1993). Human histocompatibility leukocyte antigen-DQA1\*0501 allele associated with genetic susceptibility to Graves' disease in a Caucasian population. *J. Clin. Endocrinol. Metab.*, 76: 1569-1574.

Youngchaiyud, U., Chandanayingyong, D., and Vibhatavanija, T., (1977). The incidence of HLA antigens in leprosy. *Vox. Sang.*, 32: 342-345.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่

Copyright© by Chiang Mai University

All rights reserved