

## REFERENCES

- Andersen, P. (1975). Capillary density in skeletal muscle of man. Acta Physiol. Scand., 95, 203-205.
- Aranda, J., Montoya, E., Herrera, E. (1972). Effects of hypo-and hyper-thyroidism on liver composition, blood glucose, ketone bodies and insulin in the male rat. Biochem. J., 128, 597-604.
- Argov, Z., Renshaw, P.F., Boden, B., Winokur, A, and Bank, W.J. (1988). Effects of thyroid hormones on skeletal muscle bioenergetics. J. Clin. Invest., 81, 1695-1701.
- Baldwin, K.M., Ernst, S.B., Herrick, R.E., Hooker, A.M., and Mullin, W.J. (1980). Exercise capacity and cardiac function in trained and untrained thyroid-deficient rats. J. Appl. Physiol., 49, 1022-1026.
- Baldwin, K.M., Hooker, A.M., Herrick, R.E., and Schrader, L.F. (1980). Respiratory capacity and glycogen depletion in thyroid-deficient muscle. J. Appl. Physiol., 49, 102-106.
- Balsam, A., and Leppo, L.E. (1975). Effect of physical training on the metabolism of thyroid hormones in man. J. Appl. Physiol., 38, 212.
- Ba'ra'ny, M. (1967). ATPase activity of myosin correlated with speed of muscle shortening. J. Gen. Physiol., 50, 197-216.
- Barnard, R.J., Edgerton, V.R., and Peter, J.B. (1970). Effect of exercise on skeletal muscle. II. Contractile properties. J. Appl. Physiol., 28(6), 767-770.
- Bernardis, L.L., and Skelton, F.R. (1967). Growth and obesity in male rats after placement of ventromedial hypothalamic lesions at four different ages. J. Endocrinal., 38, 351-352.

- Borensztajn, R., Rone, M.S., Babirak, S.P., McGarr, J.A., and Oscai, L.B. (1975).  
Effect of exercise on protein lipase activity in rat heart and skeletal muscle.  
Am.J. physiol., 229, 394-397.
- Boyden, T.W., Pamentor, R.W., Stanforth, P., Rotkis, T., and Wilmore, J.H. (1982).  
Evidence for mild thyroidal impairment in women undergoing endurance  
training. J. Clin. Endocrinol. Metab., 54, 53-56.
- Brodal, P., Ingjer, F., and Hermansen, L. (1977). Capillary supply of skeletal muscle  
fibres in untrained and endurance trained men. Am. J. Physiol. 232, 705-  
712.
- Burini, R., Santidrian, S., and Moreyra, M., et al., (1981). Interaction of thyroid status  
and diet on muscle protein breakdown in the rat, as measured by  $N^1$  -  
methylhistidine excretion. Metabolism, 30, 679.
- Burke, R.E., Levine D.N., Tsairis, P., and Zajac, F.E. (1973). Physiological types and  
histochemical profiles of motor units in the cat gastrocnemius. J. Physiol.  
(Lond.), 234, 723-748.
- Burke, R.E., Levine, D.N., Zajac, F.E., Tsairis, P., and Engel, W.K. (1971).  
Mammalian motor units : physiological-histochemical correlation in three  
types in cat gastrocnemius. Science Wash. DC, 174, 709-712.
- Caiozzo, V.J., Herrick, R.E., and Baldwin, K.M. (1991). The influence of  
hyperthyroidism on the maximal shortening velocity and myosin isoform  
distribution in slow and fast skeletal muscles, Am. J. Physiol., 261, C285-  
C295.
- Caiozzo, V.J., Herrick, R.E., and Baldwin, K.M. (1992). Response of slow and  
fast muscle to hypothyroidism : maximal shortening velocity and myosin  
isoforms. Am. J. Physiol., 263, C86-C94.

- Chernausek, S.D., Underwood, L.E., Utiger, R.D. (1983). Growth hormone secretion and plasma somatomedin-C in primary hypothyroidism. Clin. Endocrinol., 19, 337-344.
- Chi, M.Y., Hintz, C.S., Coyle, E.F., and Lowry, O.H. (1983). Effects of detraining on enzymes of energy metabolism in individual human muscle fibers. Am. J. Physiol., 244, C276-C287.
- Close, R. (1964). Dynamic properties of fast and slow skeletal muscles of the rat during development. J. Physiol. (Lond.), 173, 74-95.
- Close, R. (1972). Dynamic properties of mammalian skeletal muscles. Physiol. Rev., 52, 129-197.
- Cooper, D.S. (1984). Antithyroid drugs. New Engl. Med., 311, 1353-1362.
- Cooper, S., and Eccles, J.C. (1930). The isometric responses of mammalian muscle. J. Physiol. (Lond.), 69, 377-385.
- Crews, E.L., Fuge, K.W., Oscari, L.B., Holloszy, J.O., and Shank, R.E. (1969). Weight, food intake, and body composition : effects of exercise and of protein deficiency. Am. J. Physiol., 216(2), 359-363.
- Crispell, K.R., Parson, W., and Hollifield, G. (1956). A study of the rate of protein synthesis before and during the administration of L-triiodothyronine to patients with myxedema and healthy volunteers using N-15 glycine. J. Clin. Invest., 35, 164-169.
- Delia, A.J., and Thompson, E. B. (1988). A time-course study of hypothyroidism-induced hypotension : its relation to food deprivation. Arch. Int. Pharmacodyn., 296, 210-223.
- Denys, E.H., and Hofmann, W.W. (1972). An in vivo study of the biochemical changes produced by hypermetabolism and hypometabolism of skeletal muscle. Neurology., 22, 22.

- Eddinger, T.J., and Moss, R.L. (1987). Mechanical properties of skinned single fibers of identified types from rat diaphragm. *Am. J. Physiol.*, 253, C210-C218.
- Edstrom, L., and Grimby, L. (1986). Effect of exercise on the motor unit. *Muscle & Nerve*, 9, 104-126.
- Edwards, R.H. T., Jones, D.A., and Newham, D.J. (1982). Low frequency stimulation and changes in human muscle contractile properties. *J. Physiol. (Lond.)*, 328, 29.
- Everts, M.E., and Hardeveld, C.V. (1987). Effects of dantrolene on force development in slow and fast-twitch muscle of euthyroid, hypothyroid, and hyperthyroid rats. *Br. J. Pharmacol.*, 92, 47-54.
- Everts, M.E., Hardeveld, C.V., Ter Keurs, H.E.D.J., and Kassenaar, A.A.H. (1981). Force development and metabolism in skeletal muscle of euthyroid and hypothyroid rats. *Acta Endocrinol.*, 97, 221-225.
- Fangburg, B.L. (1968). Calcium transport by skeletal muscle sarcoplasmic reticulum in the hypothyroid rat. *J. Clin. Invest.*, 4, 2499-2506.
- Fisher, D.A., and Klein, A.H. (1981). Thyroid development and disorders of thyroid function in newborn. *N. Engl. J. Med.*, 304, 702-712.
- Fitts, R.H., Costill, D.L., and Gardetto, P.R. (1989). Effect of swim exercise training on human muscle fibre function. *J. Appl. Physiol.*, 66, 465-475.
- Fitts, R.H., and Holloszy, J.O. (1977). Contractile properties of rat soleus muscle : effects of training and fatigue. *Am. J. Physiol.*, 2(2), C86-C91.
- Fitts, R.H., Winder, W.W., Brooke, M.H., Kaiser, K.K., and Holloszy, J.O. (1980). Contractile, biochemical, and histochemical properties of thyrotoxic rat soleus muscle. *Am. J. Physiol.*, 238, C15-C220.

- Fitzsimons, D.P., Diffie, G.M., Herrick, R.E., and Baldwin, K.M. (1990). Effects of endurance exercise on isomyosin patterns in fast- and slow- twitch skeletal muscles. J. Appl. Physiol., 68(5), 1950-1955.
- Fitzsimons, D.P., Herrick, R.E., and Baldwin, K.M. (1990). Isomyosin distributions in rodent muscles : effects of altered thyroid state. J. Appl. Physiol., 69(1), 321-327.
- Flaim, K.E., Li, J.B., and Jefferson, L.S. (1978). Effects of thyroxine on protein turnover in rat skeletal muscle. Am. J. Physiol., 235, E231.
- Friden, J., and Sjostrom, M. (1983). Muscle fibre characteristics of endurance trained and untrained individuals. Umea University Medical Dissertations., 105, 55-62.
- Gardiner, P. F., Hibel, B., Simpson, D.R., Roy, R.R., and Edgerton, V.R. (1980). Effects of a mid weigh-lifting program on the progress of glucocorticoid induced atrophy in rat hindlimb muscle. Muscle & Nerve. 385, 147-153.
- Gold, H.K., Spann, J.F., and Braunwald, E. (1970). Effect of alterations in the thyroid state on the intrinsic contractile properties of isolated rat skeletal muscle. J. Clin. Invest., 49, 849-854.
- Gollnick, P.D., and Ianuzzo, C.D. (1972). Hormonal deficiencies and the metabolic adaptations of rats to training. Am. J. Physiol., 223, 273-282.
- Gollnick, P.D., and King, D.W. (1969). Effect of exercise and training on mitochondria of rat skeletal muscle. Am. J. Physiol., 216, 1502-1509.
- Green, H.J., Klug, G.A., Reichmann, H., Seedorf, U., Wiehrer, W., and Pette, D. (1984). Exercise- induced fiber type transitions with regard to myosin, parvalbumin, and sarcoplasmic reticulum in muscles of the rat. Pfluegers Arch., 400, 432-438.

- Gregory, P., Low, R.B., and Stirewalt, W.S. (1986). Changes in skeletal-muscle myosin isozymes with hypertrophy and exercise. Biochem. J. 238, 55-63.
- Grossie, J. (1978). Properties of thyroidectomized rat extensor muscle. Am. J. Physiol., 3(2), C90-C95.
- Halverson, P.B., Kozin, F., Ryan, L.M., and Sulaiman, A.R. (1979). Rhabdomyolysis and renal failure in hypothyroidism. Ann. Intern. Med., 91, 57.
- Hardeveld, C.v., and Kassenaar, A.A.H. (1978). Effects of experimental hypothyroidism on skeletal muscle metabolism in the rat. Acta endocrinol., 87, 114-124.
- Heilmann, C., and Pette, D. (1979). Molecular transformations in sarcoplasmic reticulum of fast-twitch muscle by electrostimulation. Eur. J. Biochem., 93, 437-446.
- Helander, E.A.S. (1961). Influence of exercise and restricted activity on the protein composition of skeletal muscle. Biochem. J., 78, 478-482.
- Hickson, R.C., Kurowski, T.T., Capaccio, J.A., and Chartier, R.T. (1984). Androgen cytosol binding in exercise-induced sparing of muscle atrophy. Am. J. Physiol., 10, E597-E602.
- Holloszy, J.O., and Coyle, E.F. (1984). Adaptations of skeletal muscle to endurance exercise and their metabolic consequences. J. Appl. Physiol., 56, 831-838.
- Houston, M.E., Benzen, H., and Larsen, H. (1979). Interrelationships between skeletal muscle adaptations and performance as studied by detraining and retraining. Acta. Physiol. Scand., 105, 163-170.
- Ianuzzo, D., Patel, P., Chen, V., and O'Brien, P. (1980). A possible thyroidal trophic influence of fast and slow skeletal muscle myosin. In : Plasticity of Muscle, edited by Pette, D. New York, de Gruyter, 593-605.

- Irvine, C.H.G. (1968). Effect of exercise on thyroxine degradation in athletes and non-athletes. J. Clin. Endocrinol. Metab., 28, 942-948.
- Izumo, S., Nadal-Ginard, B.M., and Mahdavi, V. (1986). All members of the MHC multigene family respond to thyroid hormone in a highly tissue-specific manner. Science Wash. DC, 231, 597-600.
- Janssen, J.W., Hardeveld, C.V., and Kassenaar, A.A.H. (1978). Evidence for a different response of red and white skeletal muscle of the rat in different thyroid states. Acta Endocrinol., 87, 768-775.
- Jansson, E., and Kaijser, L. (1977). Muscle adaptation to extreme endurance training in man. Acta Physiol. scand., 100, 315-324.
- Johnson, M.A., Mastaglia, F.L., Montgomery, A. G., Pope, B., and Weeds, A.G. (1980). Changes in myosin light chains in the rat soleus after thyroidectomy. FEBS Lett., 110, 230-235.
- Jones, N.L. (1988). Physiology of exercise. In : Clinical Exercise Testing. W.B. Saunders Company, Philadelphia, 58.
- Katzeff, H.L., Bovbjerg, D., and Mark, D.A. (1988). Exercise regulation of triiodothyronine metabolism. Am. J. Physiol., 255, E824-E828.
- Katzeff, H.L., O'Connell, M., Horton, E.S., Danforth, E., Young, J.B., and Landsberg, L. (1986). The impact of over-and underfeeding on energy expenditure in lean and obese man : response to graded infusions of norepinephrine. Metabolism, 35, 166-175.
- Krotkiewski, M., Sjostrom, L., Sullivan, L., Lundberg, P.A., Lindstedt G., Wetterqvist, H., and Bjorntorp, P. (1984). The effect of acute and chronic exercise on thyroid hormones in obesity. Acta Med. Scand. 216(3), 269-275.
- Lambert, E.H., Underdahl, L.O., Beckett, S., and Mederos, L.O. (1951). A study of the ankle jerk in myxedema. J. Clin. Endocrinol., 11, 1186-1951.

- Larsen, P.R., and Ingbar, S.H. (1992). The thyroid gland. In : Williams Textbook of Endocrinology, edited by Wilson, J.D., and Foster, D.W., W.B. Saunders Co., Philadelphia, 357-488.
- Laughlin, M.H., and Armstrong, R.B. (1982). Muscular blood flow distribution patterns as a function of running speed in rats. Am. J. Physiol., 243, H296-H306.
- Levy, L., Adesman, J., and Spergel, G. (1970). Studies on the carbohydrate and lipid metabolism in thyroid disease. Effect of glucagon. J. Clin. Endocrinol. Metab., 30, 372-379.
- Loucks, A.B., and Callister, R. (1993). Induction and prevention of low-T<sub>3</sub> syndrome in exercising women. Am. J. Physiol., 264, R924-R930.
- Lynch, G.S., and Williams, D.A. (1994). The effect of exercise on the contractile properties of single skinned fast and slow-twitch skeletal muscle fibres from the adult rat. Acta Physiol. Scand., 150, 141-150.
- Mai, J.V., Edgerton, V.R., and Barnard, R.J. (1970). Capillary of red, white and intermediate muscle fibers in trained and untrained guinea pigs. Experientia., 26, 1222-1223.
- Mazzeo, R.S., and Horvath, S.M. (1986). Effects of training on weight, food intake, and body composition in aging rats. Am. J. Clin. Nutr., 44, 732-738.
- McAllister, R.M., Ogivie, R.W., and Terjung, R.L. (1991). Functional and metabolic consequences of skeletal muscle remodeling in hypothyroidism. Am. J. Physiol., 260, E272-E279.
- McDaniel, H.G., Pittman, C.S., Oh, S.J., and Dimauro, S. (1977). Carbohydrate metabolism in hypothyroid myopathy. Metabolism., 26(8), 867-873.
- Mckeran, R.O., Slavin, G., Ward, P., Paul, E., and Mair, W.G.P. (1980). Hypothyroid myopathy : A clinical and pathological study. J. Pathol., 132, 35-54.



- Nicod, P., Burger, A., Staehelli, V. and Vallotton, M.B. (1976). A radioimmunoassay for 3, 3', 5'-triiodo- L- thyronine in unextracted serum : method and clinical results. J. Clin. Endocrinol. Metab., 42 : 823.
- Nihoye, L., and Mommaerts, W.F. (1981). The effects of thyroid status on some properties of rat fast-twitch muscle. J. Muscle Res. Cell Motil., 2: 307, 1981.
- Nishikawa, M., Inada, M., Naito, K., Ishii, H., Tanaka, K., Mashio, Y., and Imura, H. (1981). Age-related changes of serum 3, 3'-diiodothyronine, 3',5'-diiodothyronine, and 3, 5-diiodothyronine concentrations in man. J. Clin. Endocrinol. Metab., 52, 517-522.
- Nwoye, L., Momarets, W.F.H.M., Seraydarian, S.K., and Marusich M. (1982). Evidence for a direct action of thyroid hormone in specifying muscle properties. Am. J. Physiol., 242, R401-R408.
- Okajima, F. and Ui, M. (1979). Metabolism of glucose in hyper-and hypo-thyroid rats (in vivo) : Glucose-turnover values and futile-cycle activities obtained with <sup>14</sup>C and <sup>3</sup>H-labeld glucose. Biochem. J., 182,565.
- Oscai. L.B. (1973). The role of exercise in weight control. Exerc. Sport Sci. Rev., 1, 103-123.
- Oscai, L.B., Mole', P.A., and Holloszy, J.O. (1971). Effects off exercise on cardiac weight and mitochondria in male and female rats. Am. J. Physiol., 220, 1944-1948.
- Paul, P. (1971). Uptake and oxidation of substrates in the intact animal during exercise. In : Muscle Metabolism During Exercise, edited by Pernow, B., and Saltin, B. New York : Plenum, 225-247.
- Petrof, B. J., Kelly, A.M., Rubinstein, N.A., and Pack, A. I. (1992). Effect of hypothyroidism on myosin heavy chain expression in rat pharyngeal dilator muscles. J. Appl. Physiol., 73(1), 179-187.

- Pette, D., Smith, M.E., Staudte, H.W., and Vrbova, G. (1973). Effects of long-term electrical stimulation on some contractile and metabolic characteristic of fast rabbit muscle. Pfluegers Arch., 338, 257-272.
- Poehlman E.T., Tremblay, A., Nadeau, A., Dussault, J., The'riault, G., and Bouchard, C. (1986). Heredity and changes in hormones and metabolic rates with short-term training. Am. J. Physiol., 250(13), E711-E717.
- Rabin. D., and Mckenna, T.J. (1982). The thyroid. In : Clinical Endocrinology and Metabolism. Grune and Stratton, Inc., New York. 254-315.
- Ruff, R.L. (1986). Endocrine myopathies. In : Myology : Basic and Clinical, edited by Engel A. G., and Banker, B.Q., McGraw-Hill Book Co., New york, 1871-1906.
- Rupp, H., Felbier, H.R., Bukhari, A.R., and Jacob, R. (1984). Modulation of myosin isoenzyme populations and activities of monoamine oxidase and phenylethanolamine-N-methyltransferase in pressure loaded and normal rat heart by swimming exercise and stress arising from electrostimulation in pairs. Can. J. Physiol. Pharmacol., 62, 1209-1218.
- Rupp, H., and Wahl, R. (1990). Influence of thyroid hormones and catecholamines on myosin of swim-exercised rats. J. Appl. Physiol., 68(3), 973-978.
- Russek, M., and Pina, S. (1962). Conditioning of adrenalin anorexia. Nature, 193, 1296-1297.
- Schersten, T. (1982). Morphometric analysis of human muscle fiber types. Muscle & Nerve, 5, 538-553.
- Scow, R.O. (1959). Effect of growth hormone and thyroxine on growth and chemical composition of muscle, bone and other tissue in thyroidectomized-hypophysectomized rats. Am. J. Physiol., 196, 859-865.

- Seiden, D.A. (1976). A quantitative analysis of muscle cell changes in compensatory hypertrophy and work-induced hypertrophy. Am. J. Anat., 145, 459-468.
- Sjodin, B. (1976). Lactate dehydrogenase in human skeletal muscle. Acta Physiol. Scand. Suppl., 436, 9-18.
- Song, M. K., Ianuzzo, C.D., Saubert, I.V., and Gollnick (1973). The mode of adrenal gland enlargement in the rat in response to exercise training. Pfluegers Arch., 339, 59-68.
- Spiro, A.J., Hirano, A., Beilin R.L., and Finkelstein, J.W. (1970). Critinism with muscular hypertrophy (Kocher-Debre-Semelaigne Snyderome). Arch. Neurol., 23, 340.
- Tata, J.R., Ernster, L., Lindberg, O., Arrhenius, E., Pedersen, S., and Hedman, R. (1963). The action of thyroid hormones at the cell level. Biochem. J., 86, 408-427.
- Terjung, R.L., and Koerner, J.E. (1976). Biochemical adaptation in skeletal muscle of trained thyroidectomized rats. Am. J. Physiol., 230, 1194-1197.
- Tipton, C.M., Terjung, R.L., and Barnard, R.J. (1968). Response of thyroidectomized rats to training. Am. J. Physiol., 215, 1137-1142.
- Tsika, P.W., Herrick, R.E., and Baldwin, K.M. (1987). Subunit composition of rodent isomyosins and their distribution in hindlimb skeletal muscles. J. Appl. Physiol., 63(5), 2102-2110.
- Utiger, R.D. (1989). Hypothyroidism. In : Endocrinology, edited by DeGroot, L.J., et al., Grune and Stratton, Inc., New York, 471-488.
- Van Lunteren, E., Salomone, R.J., Manubary, P., Supinski, G.S., and Dick, T.E. (1990). Contractile and endurance properties of geniohyoid and diaphragm muscles. J. Appl. Physiol., 69, 1992-1997.

- Vendsalu, A. (1960). Plasma concentrations of adrenaline and noradrenaline during muscular work. Acta. Physiol. Scand. Suppl., 173, 57-69.
- Westgren, U., Burger, A., Ingemansson, S., Melander, A., Tibblin, S., and Wahlin, E. (1976). Blood levels of 3, 5, 3'-triiodothyronin and thyroxine : differences between children, adults, and elderly subjects. Acta Med. Scand., 200, 493.
- Wiles, C.M., Young, A., Jones, D.A., and Edwards, R.H.T. (1979). Muscle relaxation rate, fibre-type composition, and energy turnover in hyper-and hypo- thyroid patients. Clin. Sci. Lond., 57, 375-384.
- Winder, W.W., and Heninger, R.W. (1973). Effect of exercise on degradation of thyroxine in the rat. Am. J. Physiol., 224(3), 572-575.