

REFERENCES

Ball, J. 1982, Stream Classification Guidelines for Wisconsin. Wisconsin

Department of Natural Resources. Technical Bulletin. Wisconsin Department
of Natural Resources, Madison, Wisconsin.

Bishop K. A., Pidgeon R. W. J. and Walden D. J. 1995. Studies on fish

movement dynamics in a tropical floodplain river: Pre-requisites for a

procedure to monitor the impacts of mining. Aust. J. Ecol.(20), 81-107.

Butcher, R.W., Pentelow, F.T.K. and Woodley, J.W.A. 1931. An investigation of

the River Lark and the effect of beet-sugar pollution. Fish Invest, Lond. Series

1,(3),No.3.

Carpenter, K.E. 1926. The lead mine as an active agent in river pollution. Ann. appl. Biol.(13) 395-401.

Chandler, J.R. 1970. A biological approach to water quality management. Water Pollut. Control,(69), 415-422.

Chapman D. 1992. Water Quality Assessments. Chapman&Hall, London. 585 pp.

Chessman B. C. 1995. Rapid assessment of rivers using macroinvertebrates: A procedure based on habitat-specific sampling, family level identification and a biotic index. Aust.J. Ecol.(20),122-9.

Dowing, J.A. and Rigler, F. H.(eds). 1984. A manual of methods for the assessment of secondary productivity on fresh waters. Backwell, Oxford.

DIN 38410 T2. 1990. German Standard Methods for Examination of Water, Wastewater and Sludge. Biologischökologische Gewässerynlersuchung

Beslimmung des Saprobie-nindex. Berlin.

Elliott, J.M. 1979. Some methods for the statistical analysis of benthic invertebrates. Scientific Publication No.25. Fershwater Biological Association, Windermere.

Flannagan, J. F. 1970. Efficiencies of various grabs and corers in sampling freshwater benthos, J. Fish. Res. Bd. Can.,(27),1691-1700.

Green, R.H. 1979. Sampling design and sttistical methods for environmental biologists. Wiley, New York.

Hagebro, C., Bang,S.and Somer, E. 1983. Nitrate/load discharge relationships and nitrate laod trends in Danish rivers. IAHS Publ. No. 141, 377-386.

Harris J.H. 1995. The use of fish in ecological assessments. Aust. J. Ecol.(20),

65-80.

Haslam S.M. 1990. River Pollution: An Ecological Perspective. Belhaven Press,
London. P. 133-138 .

Hawkes, H.A. 1959. The biological assessment of pollution in Birmingham
Streams. J. Instn. Munic. Engrs. 82452, Surveyor, LO nd. 115, 129-131.

Hellawell, J.M. 1986^a. Biological indicators of freshwater pollution and
environmental management. Applied Science Publishers, London.

Hellawell, J.M. 1986^b. Biological Indicators of Feshwater Pollution and
Environmental Management. Pollution Monitoring Series, K. Mellangy(Ed.),
El-sevier Applied Sciences Publishers, London, 546 pp.

Hughes, B.D. 1975. A comparison of four samplers for benthic
macroinvertebrates inhabiting coarse river deposits. Water Res.,(9),61-69.

Humphy C.L.,P.L. Dostine.1994. Development of biological monitoring

programmes to detect mining-waste impacts upon aquatic ecosystem of the

Alligator River Region Northern Territory, Australia Mitt. Internat. Verein.

Limno.(24)293-314.

Hynes, H.B.N. 1959. The use of invertebrates as indicators of river pollution.

Proc.Linn.Soc.Lond.(2) 165-169.

Hynes, H.B.N. 1960. The Biology of Polluted Waters, Liv. Un. Press. 202 pp.

ISO- BMWP. 1979. Assessment of the Biological Quality of Rivers by a

Macroinvertebrate Score. ISO/TC147/SC5/WG6/N5, International Standards

Organization, 18 pp.

Kinzie, R. A., J. I. Ford, A. R. Yuen and Chow, S. J. L. 1986. Habitat

Modelling of Hawaiian Streams. Water Resources Research Center.

University of Hawaii at Manoa. Honolulu, Hawaii.

Kolkwitz, R. & Marsson, M. 1908. Ökologie der pflanzlichen Saprobien. Ber.

dt. bot. Ges.(26),505-519

Kolkwitz, R. & Marsson, M. 1909. Ökologie der tierischen Saprobien. Int. Rev.

ges. Hydrobiol.(2), 126-152.

Lehmkuhl D. M. 1979. Aquatic Insect. Wm. C. Brown Company Publishers,

Dubuque, Iowa, USA.

Ludwig, J. A. and Reynolds, J.F. 1988. Statistical Ecology. A Primer on Methods

and Computing. John Wiley and Sons, New York, USA. 339pp.

McCafferty, W. P. 1981. Aquatic Entomology. The Fishermen's Illustrated

Guide to Insects and Their Relatives. Science Books International, Boston.

Mellanby, M. 1963. 6th ed. Animal life in fresh water. A Guide to Fresh-Water

Invertebrates. Methuen and Co. Ltd., London.

Needham, P. R., 1969. 5th ed. A Guide to the study of Fresh-Water Biology.

Holden-Day, Inc. San Francisco.

Peckarsky, B.L. 1984. Sampling the stream benthos. In Doving, J.A. and Rigler,

F.H.(eds). A manual on methods for the assessment of assessment of

secondary productivity in freshwaters, pp. 131- 160. Backwell, Oxford.

Pennak, R.W. 1978. Fresh-Water Invertebrates of the United States.

John Wiley & Son, Inc. New York. 803 pp.

Pinder, L.C.V. and Farr, I.S. 1987. Biological surveillance of water quality-2

temporal and spatial variation in the macroinvertebrate fauna of the River

Frome, a Dorset Chalk Stream. Arch. Hydrobio., 109(3), 321-331.

Plafkin, J.L., M.T. Barbour, K.D. Porter, S.K. Gross and Hughes, R.M. 1989.

Rapid Bioassessment Protocols for Use in Streams and Rivers.

- Benthic Macroinvertebrates and Fish. EPA/440/4-89/001. Office of Water Regulations and Standards U.S. EPA, Washington, D.C.
- Platts, W.S., W.F. Megahan, and Minshall, G.W. 1983.** Method for Evaluating Stream, Riparian, and Biotic Conditions. General Technical Report INT-138. U.S. Department of Agriculture, U.S. Forest Service, Ogden, Utah.
- Rajchapakdee, M. 1992.** Benthic Invertebrates Relative to Altitude and Water Quality in Streams in Doi Suthep. M.S. Thesis, Chiang Mai University.
- Reid M. A., Tibby J. C., Penny D. and Gell P.A. 1995.** The use of diatoms to assess past and present water quality. *Aust. J. Ecol.* (20), 57-64.
- Resh V. H. and Rosenberg, D.M. 1993.** Freshwater Biomonitoring and Benthic Macroinvertebrates. Chapman & Hall, London.
- Resh V. H. 1979.** Sampling variability and life history features: Basic considerations

in the design of aquatic insect studies. J. Fish. Res. Board Can.(36),290-311.

Richardson,R.E. 1928. The bottom fauna of the Middle Illinois River 1913-1925:

its distribution, abundance, variation and index value in the study of stream

pollution. Bull.Nat. Hist. Surv., Chicago,(17), 387-475.

Roberts, G. and March, T. 1987. The effects of agricultural practices on the nitrate

concentrations in the surface water domestic supply sources of western Europe.

Water for the future : Hydrology in Perspective, LAHS Publ.(164), 365-380.

Rutter, A. J., 1958. The effects of afforestation on rainfall and run-off. J. Instn

Publ.Heth Engrs, 119-38. 2,3.

Sannarm, G. 1993. Biological surveillance using macroinvertebrate of the quality

of the Mae Kwaung River the Northen Region Industrial Estat. M.S. Thesis,

Chiang Mai , University.

Storey A. W., Edward D. H. D. and Gazey P. 1991. Syrber and kick sampling: A

comparison for the assessment of macroinvertebrate community structure in streams of southwestern Australia. *Hydrobiologia*(211), 111-21.

Stalnaker, C. B. 1979. The use of habitat structure preferenda for

eastablishing flow regimes necessary for maintenance of fish habitat.

In *The Ecology of Regulated Streams*, ed. J. C. Ward and J. A.

Stanford, pp. 321-37.

Thaweeburus, P. 1994. Biomonitoring techniques to assess water loss at the Old

Mae Ping Dam and water quality of the Mae Ping River. M.S. Thesis,

Chiang Mai University.

Thuedam, Y., 1994. Use of Macroinvertebrate Communities to Assess Water

Quality Class in Doi Inthanon Stream and Ping River by Biotic and Saprobic

Indices. MS Thesis, Dept. of Biol., Chiang Mai Univ.

Tolkamp, H.H. 1985. Using several indices for biological assessment of water quality in running water. *Verh. Internat. Verein. Limnol.*, (22), 2281-2286.

Tuffery, G. and Verneaux, J. 1967. Methode de determination de la qualite biologique des eaux courantes. Exploitation codifiee des inventaires de la faune de fond. *Trav. Sect. P. et P., Cerafer, Paris*, 23 pp.

Welch, E.B. 1992. Ecological Effects to Wastewater. Chapman & Hall, London. 425 pp.

Wetzel, R.G. and Linkens, G.E. 1979. Limnology analyses. W.B. Saunders Company, London. 357 pp.

Whitton B. A. and Kelly M. G. 1995. Use of algae and other plants for monitoring rivers. *Aust. J. Ecol.* (20), 45-56.

Williams W. D., 1991. Australian Freshwater Life. Set in Bembo by Savage & Co.,

Brisbane, Hong Kong. 321 pp.

Wright J. F., Armitage P. D. and Furse M. T. 1989. Prediction of invertebrate

communities using stream measurements. *Reg. Riv.*(4), 147-55.

Wright J. F. 1995. Development and use of a system for predicting the

macroinvertebrate fauna in flowing waters. *Aust. J. Ecol.*(20), 181-97.