

## VII. REFERENCES

- Abbas, A.K., Lichtman, A.H. and Pober, J.S. (1994). The major histocompatibility complex. In Abbas, A.K., Lichtman, A.H. and Pober, J.S. (Ed.), Cellular and Molecular Immunology (pp.97-114), W.B.Saunders.
- Adams, D.D. and Purves, H.D. (1956). Abnormal response in the assay of thyrothophin. Proceeding of the University of Otago medical school 34: 11-12.
- Anderson, G., Anderson, L., Larhammar, D., Rask, L. and Sigurdardóttir, S. (1994). Simplifying genetic locus assignment of HLA-DRB genes. Immunol Today 15(2): 58-62.
- Austyn, J.M. and Wood, K.J. (1994). The major histocompatibility complex. In Austyn, J.M. and Wood, K.J. (Ed.), Principles of Cellular and Molecular Immunology (pp.63-114), Oxford University Press.
- Auwerda, B., Schuit, F., Lyaruu, I., Falorni, A., Svanholm, S., Vandewalle, C.L. and Goris, F.K., (1995). Genetic susceptibility for insulin-dependent diabetes mellitus in Caucasians revisited: the importance of diabetes registries in disclosing interaction between HLA-DQ and insulin gene-linked risk. Belgian Diabetes registry. J. Clin. Endocrinol. Metab. 80(9): 2567-2573.
- Badenhoop, K., Walfish, P.G., Rau, H., Fischer, S., Nicolay, A., Bogner, U., Schleusener, H. and Usadel, K.H. (1995). Susceptibility and resistance alleles of human leukocyte antigen (HLA) DQA1 and HLA DQB1 are shared in endocrine autoimmune disease. J. Clin. Endocrinol. Metab. 80(7): 2112-2117.
- Baldini, M., Pappalettera, M., Lecchi, L., Orsatti, A., Meroni, L., Tozzi, R., Scalamogna, M. and Cantalamessa, L. (1995). Human lymphocyte antigens in Graves' disease: Correlation with persistent course of disease. Am. J. Med. Sci. 309(1): 43-48.
- Begovich, A.B., Vu, T.H. and Jones, P.P. (1990). Characterization of the molecule defects in the mouse E beta f and E beta q genes. Implication for the origin of MHC polymorphism. J. Immunol. 144: 1957-1964.

- Bidwell, J. (1994). Advances in DNA-based HLA-typing methods. *Immunol Today* 15(7): 303-307.
- Bidwell, J.L., Bidwell, E.A. and Bradley, L.A. (1990). HLA class II gene: typing by DNA analysis. *Baillieres Clin. Haematol.* 3(2): 355-384.
- Bidwell, J.L. (1988). DNA-RFLP analysis and genotyping of HLA-DR and DQ antigens. *Immunol. Today* 9(1): 18-23.
- Bidwell, J. and Jarrold, E.A. (1986). HLA-DR allogenotyping using exon-specific cDNA probes and application of rapid minigel methods. *Molecular Immunology* 23: 1111-1116
- Bodmer, J.G., Marsh, S.G.E., Albert, E.D., Bodmer, W.F., Dupont, B., Erlich, H.A., Mach, B., Mayr, W.R., Parham, P., Sasazuki, T., Schreder, G.M.Th., Strominger, J.L., Svejgaard, A., and Terasaki, P.I. (1994). Nomenclature for factors of the HLA system, 1994. *Tissue Antigens* 44: 1-18.
- Bottazzo, G.F., Borrell, R.P. and Hanafusa, T. (1983). Role of aberrant HLA-DR expression and antigen presentation in induction of endocrine autoimmunity. *Lancet* 12: 1115-1119.
- Bottazzo, G.F. and Doniach, D. (1986). Autoimmune thyroid disease. *Annu. Rev. Med.* 37: 353-359.
- Brown, J.H., Jardetzky, T.S., Saper, M.A., Samraoui, B., Bjorkman, P.J. and Wiley, D.C. (1988). A hypothetical model of the foreign antigen binding site of class II histocompatibility molecules. *Nature* 332: 845-850.
- Brown, J.H., Jardetzky, T.S., Gorga, J.C., Stern, L.J., Urban, R.G., Strominger, J.L. and Wiley, D.C. (1993). Three-dimensional structure of the human class II histocompatibility antigen HLA-DR1. *Nature* 364: 33-39.
- Carson, D.A. (1992). Genetic factors in the etiology and pathogenesis of autoimmunity. *The FASEB J.* 6: 2800-2805.

- Caven, D. A., Penny, M.A., Jacob, K.H., Kelly, M.A., Jenkins, D., Mjovic, C., Chow, C., Oock, C.S., Hawkin, B.R. and Bennett, A.H. (1994). The HLA association with Graves' disease is sex specific in Hong Kong Chinese subjects. *Clin. Endocrinol.* 40: 63-66.
- Cerna, M., Falco, M., Friedman, H., Raimondi, E., Maccagna, A., Fernandez-Viña, M. and Stastny, P. (1993). Differences in HLA class II alleles of isolated South American Indian populations from Brazil and Argentina. *Hum. Immunol.* 37(4): 213-220.
- Chan, S.H., Lin, Y.N., Wee, G.B., Ren, E.C., Lui, K.F. and Cheah, J.S. (1993). Human Leucocyte antigen DNA typing in Singaporean Chinese patients with Graves' disease. *Ann. Acad. Med. Singapore* 22(4):867-879.
- Charron, D. (1990). Molecular basis of human leukocyte antigen class II disease associations. *Adv. Immunol.* 48: 107-159.
- Chicz, R.M., Urban, R.G., Gorga, J.C., VIGNALI, D.A.A., Lane, W.S. and Strominger, J.L. (1993). Specificity and promiscuity among naturally processed peptides bound to HLA-DR alleles. *J. Exp. Med.* 178: 27-47.
- Cho, B.Y., Rhee, B.D., Lee, D.S. Lee, M.S., Lee, H.K., Koh, C., Min, H.K. and Lee, M. (1986). HLA and Graves' disease in Korean. In Vichayanrat, A., Nitiyanant, W., Eastman, C. and Nagataki, S. (Ed.). Recent progress in thyroidology (pp.188-192), Crystall House press.
- Creagh, F., Teece, M., Williums, S., Didcote, S., Perkins, W., Hashim, F. and Smith, R. (1985). An analysis of thyrotropin receptor binding and thyroid stimulating activities in a series of Graves' sera. *Clin. Endocrinol. (Oxf.)* 23 (4): 395-404.
- Cuddihy, R.M. and Bahn, R.S. (1996). Lack of an independent association between the human leukocyte antigen allele DQA1\*0501 and Graves' disease. *J. Clin. Endocrinol. Metab.* 81(2): 847-849.

Dahlberg, P.A., Holmlund, G., Karlsson, F.A. and Säfwenberg, J. (1981). HLA-A, -B, -C and -DR antigens in patients with Graves' disease and their correlation with signs and clinical course. *Acta Endocrinologica* 97: 42-47.

DeGroot, L.J. (1996). Autoimmunity to the thyroid gland. In DeGroot, L.J., Larsen, P.R., Hennemann, (Ed.). *The thyroid and its disease* (pp.255-306), Churchill livingstone.

Doherty, D.G., Vaughan , R.W., Donaldson, P.T. and Mowat, A.P. (1992). HLA DQA, DQB, and DRB genotyping by oligonucleotide analysis: distribution of alleles and haplotypes in British Caucasoids. *Hum. Immunol.* 34(1): 53-63.

Dong, R., Kimura, A., Okubo, R., Shinagawa, H., Tamai, H., Nishimura, Y. and Sasazuki, T. (1992). HLA-A and DPB1 loci confer susceptibility to Graves' disease. *Hum. Immunol.* 35: 165-172.

Dyer, P. and Middleton, D. (1993). Introduction. In Dyer, P. and Middleton, D. (Ed.), *Histocompatibility testing a practical approach* (pp.1-13), IRL press.

Eisenstein, Z., Engelman, E., Weiso, M., Kalechman, Y. and Sredni, B. (1994). Modulation of the IL-2 production defect in vitro in Graves' disease. *Clin. Exp. Immunol.* 96(2): 323-328.

Farid, N.R., Sampson, L., Noel, E.P., Bernard, J.M., Mandeville, R., Larsen, B. and Marshall, W.H. (1979). A study of human leukocyte D locus related antigen in Graves' disease. *J. Clin. Invest.* 63: 108-113.

Feliciano, D.V. (1992). Everything you wanted to know about Graves' disease. *Am. J. Surg.* 164: 404-411.

Festenstein, H., Awad, J., Hitman, G.A., Cutbush, S., Groves, A.V., Cassell, P., Ollier, W. and Sachs, J.A. (1986). New HLA DNA polymorphisms associated with autoimmune disease. *Nature* 322: 64-67.

- Gao, X., Sun, Y., An, J., Fernandez-Viña, M., Qou, J., Lin., L. and Stastny, P. (1991). DNA typing for HLA-DR and -DP alleles in a Chinese population using the polymerase chain reaction (PCR) and oligonucleotide probes. *Tissue Antigens* 38: 24-30.
- George, A.J.T., Ritter M.A. and Lechner, R.I. (1995). Disease susceptibility, transplantation and the MHC. *Immunol. Today* 16(5): 209-211.
- Geng, L., Imanishi, T., Tokunaga, K., Zhu, D., Mizuki, N., Xu, S., Geng, Z., Gojobori, T., Tsuji, K. and Inoko, H. (1995). Determination of HLA class II alleles by genotyping in a Manchu population in the northern part of China and its relationship with Han and Japanese population. *Tissue Antigens* 46(32): 111-116.
- Gergely, P. (1992). The pathogenesis of autoimmune disease. *Immunol. Today* 13(8): A21-A23.
- Ghosh, P., Amaya, M., Mellins, E. and Wiley, D.C. (1995). The structure of an intermediate in class II MHC maturation: CLIP bound to HLA-DR3. *Nature* 378: 457-462.
- Gossage, A.A.R. and Munro, D.S. (1985). The pathogenesis of Graves' disease. *Clin. Endocrinol. Metab.* 14(2): 299-331.
- Gray, J. and Hoffenberg, R. (1985). Thyrotoxicosis and stress. *Quarterly J. Med.* 54(214): 153-160.
- Green, A. (1982a) The epidemiologic approach to studies of association between HLA and disease I. The basic measures, concepts and estimation procedures. *Tissue Antigens* 19: 245-258.
- Green, A. (1982b) The epidemiologic approach to studies of association between HLA and disease II. Estimation of absolute risks, etiologic and preventive fraction. *Tissue Antigens* 19: 259-268.

Grumet, F.C., Payne, R.O., Konishi, J. and Kriss, J.P. (1974). HL-A antigens as markers for disease susceptibility and autoimmunity in Graves' disease. *J. Clin. Endocrinol. Metab.* 39(6): 1115-1119.

Haldane, J.B.S. (1955). The estimation and significant of the logarithm of a ratio of frequencies. *Ann. Hum. Genet.* 20:309-311.

Hammer, J., Valsasnini, P., Tolba, K., Bolin, D., Higelin, J., Takacs, B. and Sinigaglia, F. (1993). Promiscuous and allele-specific anchors in HLA-DR binding peptides. *Cell* 74: 197-203.

Hanafusa, T., Pujol-Borrell, R., Chiovato, L., Rossell, R.C.G., Doniach, D. and Bottazzo, G.F. (1983). Aberrant expression of HLA-DR antigen on thyrocytes in Graves' disease: Relevance for autoimmunity. *Lancet* 12: 1111-1114.

Hansen, T.H., Carreno, B.M. and Sachs, D.H. (1993). The major histocompatibility complex. In Paul, W.E. (Ed.), *Fundamental Immunology* (pp.577-628), Raven Press.

Hashim, F.A., Creagh, F.M., El Hawrani, A., Parkes, A.B., Buchland, P.R. and Smith, R. (1986). Characterization of TSH antagonist activity in the serum of patients with thyroid disease. *Clin. Endocrinol. (Oxf.)* 25: 275.

Herold, K.C. and Quintans, J. (1995). Immunological mechanisms causing autoimmune endocrine disease. In Quintans, J. (Ed.), *Endocrinology* (pp.2990-3012), W.B. Saunders.

Hill, A.V.S., Yates, S.N.R., Allsopp, C.M., Gupta, S., Gilbert, S., Lalvani, A., Aidoo, M., Davenport, M. and Plebanski, M. (1994). Human leukocyte antigens and natural selection by malaria. *Phil. Trans. R. Soc. Lond. B* 346: 379-385.

Holmdahl, R., Karlsson, M., Anderson, M.E., Rask, L. and Anderson, L. (1989). Localized of a critical restriction site on the I-A beta chain that determines susceptibility to collagen-induced arthritis in mice. *Proc. Natl. Acad. Sci. USA* 86 (23): 9475-9479.

Huang, H., Peng., J., She, J.Y., Zhang, L., Chao, C.C.K., Liu, K. and She, J. (1995). HLA-encoded susceptibility to insulin dependent diabetes mellitus is determined by DR and DQ genes as well as their linkage disequilibria in a Chinese population. *Hum. Immunol.* 44: 21-219.

Imanishi, T., Akaza, T., Kimura, A., Tokugawa, K. and Gojobori, T. (1992). Allele frequencies and haplotype frequencies for HLA and complement loci in various ethnic group. *HLA 1991*. In Tsuji, K., et al. (Ed.), vol 1. Oxford University press.

Inoue, D., Sato, K., Enomoto, T., Sugawa, H., Maeda, M., Inoko, H., Tsuji, K., Mori, T. and Imura, H. (1992). Correlation of HLA types and clinical finding in Japanese patients with hyperthyroid Graves' disease: evidence indicating the existence of four subpopulation. *Clin. Endocrinol.* 36: 75-82.

Jacob, K., Ruderhof, R., Neill, S.D., Dougherty, J.P., Brown, E.L. and Fritsch, E.F. (1988). The thermal stability of oligonucleotide duplexes is sequence independent in Tetraalkylammonium solution: application to identifying recombinant DNA clones. *Nucleic Acids Res.* 16(10): 4637-4650.

Kendell, E., Todd, J.A. and Campbell, R.D. (1991). Molecular analysis of the MHC class II region in DR4, DR7 and DR9 haplotypes. *Immunogenetics* 34: 349-357.

Kimura, A. and Sasazuki, T. (1991). Eleventh international histocompatibility workshop reference protocol for the HLA DNA-typing technique, W 3.3, Yokohamo Japan. November 6-13.

Koopman, G., and Pals, S.P. (1992). Cellular interaction in the germinal center: Role of adhesion receptors and significance for pathogenesis of AIDS and malignant lymphoma. *Immunol. Rev.* 126-: 21-45.

Kunachiwa, W. (1995). HLA class II in leprosy patients in northern Thailand. M.S. Thesis, Mahidol University.

Kwok, W.W., Nepom, G.T. and Raymond, F.C. (1995). HLA-DQ polymorphisms are highly selective for peptide binding interactions. *J Immunol.* 155: 2468-2476.

Lawley, T.J., Hall, R.P., Fauci, A.S. Katz, S., HamBurger, M.I. and Frank, M.M. (1981). Defective Fc receptor function associated with the HLA-B8/DRw3 haplotype. *N. Eng. J. Med.* 304(4): 185-192.

Liblau, R.S., Singer, S.M. and McDevitt, H.O. (1995). Th1 and Th2 CD4+ T cell in the pathogenesis of organ-specific autoimmune diseases. *Immunol. Today* 16(1): 34-38.

Linsley, P.S., Wallace, P.M., Johnson, J., Gibson, M.G., Greene, J.L., Lebbetter, J.A., Singh, C. and Tepper, M.A. (1992). Immunosuppression in vivo by a soluble form of the CTLA-4 T cell activation molecule. *Science* 257: 792-795.

Lou, G., Lin, S., Chen, X. and Pan, C. (1994). Genetic analysis of HLA-DR alleles in Graves' disease by PCR-amplified DNA with sequence specific of oligonucleotide probes. In the 10<sup>th</sup> Asian-Oceania congress of endocrinology, Beijimg China. 30th October-3rd November

Ludgate, M.E. and Vassart, G. (1995). The thyrotropin receptor as a model to illustrate receptor and receptor antibody diseases. *Baillière Clin. Endocrinol. Metab.* 9(1): 95-113.

Macchia, E., Concetti, R., Carone, G., Borgoni, F., Fenzi, G.F. and Pinchera, A. (1988). Demonstration of blocking immunoglobulins G, having a heterogenous behaviour, in sera of patients with Graves' disease: possible coexistence of different autoantibodies directed to the TSH receptor. *Clin. Endocrinol. (Oxf).* 28: 147-156.

Mangklabruks, A., Cox, C. and DeGroot, L.J. (1991). Genetic factors in autoimmune thyroid disease analyzed by restriction fragment length polymorphisms of candidate genes. *J. Clin. Endocrinol. Metab.* 73(2): 236-244.

Marazuela, M., Vargas, J.A., Durantez, A. and Alvarez-Mon, M. (1994). T lymphocyte activation in Graves' disease. *Endocrine Research* 20(3): 291-306.

Marsh, S.G.E. and Bodmer, J.G. (1995). HLA class II region nucleotide sequences, 1995. *Tissue Antigens* 45:258-280.

Matsumura, M., Fremont, D.H., Peterson, P.A. and Wilson, I.A. (1992). Emerging principles for the recognition of peptide antigens by MHC class I molecules. *Science* 257: 927-934

McCloskey, D.J., Brown, J. and Navarrete, C. (1993). Serological typing of HLA-A, -B and -C antigens. In Hui, K.M. and Bidwell, J. (Ed.), *Hand book of HLA typing techniques* (pp.175-247), CRC Press.

McDougall , I.R. (1991). Graves' disease current concepts. in Greenspan, P.S. (Ed.), *The medical clinics of north America* (pp.79-95), W.B. Saunders company.

McGregor, A., Rees Smith, B., Hall, R., Peterson, M.M., Miller, M. and Dewar, P.J. (1980). Prediction of relapse in hyperthyroid Graves' disease. *Lancet* i: 1101-1103.

Middleton, D., Bodmer, J., Heyes, J. and Marsh, S. (1993). HLA typing reagents: alloantisera and monoclonal antibodies. In Dyer, P. and Middleton, D. (Ed.), *Histocompatibility Testing A Practical Approach* (pp.13-50), IRL Press.

Miguel, A., Gonzalez-Gay, M.A., Nabozny, G.H., Bull, M.J., Zanelli, E., Douhan III, J., Griffiths, M.M., Glimcher, L.H., Luthra, H.S., and David, C.S. (1994). Protective role of major histocompatibility complex class II Eb<sup>d</sup> transgene on collagen-induced arthritis. *J. Exp. Med.* 180: 1559-1564.

Miller, S.A., Dykes, D.D. and Polesky, H.F. (1988). A simple salting-out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res.* 16: 1215.

Nagashima, M., Eguchi, K., Ida, H., Yamashita, L., Sakai, M., Origuchi, T., Kawabe, Y., Ishikawa, N., Ito, K. and Nagataki, S. (1994). The Expression of adhesion molecules in thyroid gland from patients with Graves' disease. *Thyroid* 4(1): 19-25.

- Nagayama, Y. and Rapoport, B. (1992). Thyroid stimulatory autoantibodies in different patients with autoimmune thyroid disease do not all recognize the same components of the human thyrotropin receptor: selective role of receptor amino acids Ser<sub>25</sub>-Glu<sub>30</sub>. *J. Clin. Endocrinol. Metab.* 75(6): 1425-1430.
- Navarrete, C., McClosky, D.J. and Brown, J. (1993). Definition of HLA-Dw and HLA-DPw determinations by the primed lymphocyte test. In Hui, K.M. and Bidwell, J. (Ed.), *Hand book of HLA typing techniques* (pp.351-372), CRC Press.
- Ober, K.P. (1992). Thyrotoxic periodic paralysis in the United States report of 7 cases and review of the literature. *Medicine* 71(3): 109-120.
- Onuma, H., Ota, M., Sugenoya, A. and Inoko, H. (1994). Association of HLA-DPB1\*0501 with early-onset Graves' disease in Japanese. *Hum. Immunol.* 39: 195-201.
- Orhan, Y., Azezli, A., carin, M., Aral, F., Sencer, E. and MolValilar, S. (1993). Human Leukocyte antigens (HLA) and Graves' disease in Turkey. *J Clin. Immunol.* 13(5): 339-343.
- O'Sullivan, D., Sidney, J., del Guercio, M., Colon, S.M. and Sette, A.J. (1991). Truncation analysis of several DR binding epitopes. *J. Immunol.* 146(4): 1240-1246.
- Paulsen, G., Lundin, K.E.A., Gjertsen, H.A., Hansen, T., Sollid, L.M. and Thorsby, E. (1995). HLA-DQ2-restricted T-cell recognition of gluten-derived peptides in celiac disease influence of amino acid substitutions in the membrane distal domain of DQ $\beta$  1\*0201. *Hum. Immunol.* 42: 145-153.
- Payami, H., Joe, S., Farid, N., Stenszky, V., Chan, S.H., Yeo, P.P.B., Cheah, J.S. and Thomson, G. (1989). Relative predispositional effects (RPEs) of marker alleles with disease: HLA-DR alleles and Graves' disease. *Am. J. Hum. Genet.* 45: 541-546.
- Powis, S.H. and Geraghty, D.E. (1995). What is the MHC?. *Immunol. Today* 16(10): 466-468.

Ratanachaiyavong, S., Gunn, C.A., Bidwell, E.A., Darke, C., Hall, R. and McGregor, A.M. (1990). DQA2 U allele: a genetic marker for relapse of Graves' disease. *Clin. Endocrinol.* 32: 241-251.

Ratanachaiyavong, S., Bidwell, J.L., Bidwell, E.A. and McGregor, A.M. (1991). Linkage disequilibrium between the HLA-DQA2 alleles and the HLA-DR/DQ complex. *Human Immunol.* 30: 136-139.

Ratanachaiyavong, S. and McGregor, A.M. (1994). HLA-DPB1 polymorphisms on the MHC-extended haplotypes of families of patients with Graves' disease: two distinct HLA-DR17 haplotypes. *Eur. J. Clin. Invest.* 24: 309-315.

Rønningten, K.S., Spurkland, A., Markussen, G., Iwe, T., Vartdal, F. and Thorsby, E. (1990). Distribution of HLA class II alleles among Norwegian Caucasian. *Hum. Immunol.* 29: 275-281.

Ropars, A., Marion, S., Takorabet, L., Braun, J. and Charreire, J. (1994). Antibodies specific for human thyrotropin receptor induce MHC antigen expression in thyroid cells. *J. Immunol.* 153: 3345-3352.

Saiki, R.K., Walsh, P.S., Levenson, C.H. and Erlich, H.A. (1989). Genetic analysis of amplified DNA with immobilized sequence-specific oligonucleotide probes. *Proc. Natl. Acad. Sci. USA* 86(16): 6230-6234.

Saper, M.A., Bjorkman, P.J. and Wiley, D.C. (1991). Refined structure of the histocompatibility antigen HLA-A2 at a 2.6 Å° resolution. *J. Mol. Biol.* 219(2): 277-319

Schleusener, H., Schleusener, G., Mayr, W.R., Kotulia, P., Bogner, U., Finke, R., Meinholt, H., Kopenhagen, K. and Wenzel, K.W. (1983). HLA-DR3 and HLA-DR5 associated thyrotoxicosis-two different types of toxic diffuse goiter. *J Clin. Endocrinol. Metab.* 56(4): 781-785.

Segall, M. (1988). HLA and genetics of IDDM holism & reductionism?. *Diabetes* 37: 1005-1008.

Sette, A., Sidney, J., Gaeta, F.C.A., Appella, E., Colón, S.M., del Guercio, M., Guéry, J. and Adorini, L. (1992). MHC class II molecules bind indiscriminate self and non-self peptide homologs: effect on the immunogenicity of non-self peptides. *Inter. Immunol.* 5(6) : 631-636.

Shaw, G. and Kamen, R. (1986). A conserved AU sequence from the 3' untranslated region of GM-CSF mRNA mediates selective mRNA degradation. *Cell* 46(5): 659-667.

Smith, B.R. and Hall, R. (1974). Binding of thyroid stimulators to thyroid membranes. *The FEBS Lett* 42: 301-304.

Soliman, M., Kaplan, E., Yanagawa, T., Hidaka, Y., Fisfalen, M. DeGroot, L.J. (1995). T-cells recognize multiple epitopes in the human thyrotropin receptor extracellular domain. *J. Clin. Endocrinol. Metab.* 80(3): 905-914.

Song, Y., Li, Y. and Maclaren, N.K. (1996). The nature of autoantigens targeted in autoimmune endocrine diseases. *Immunol. Today* 17(5): 232-238.

Stastny, P., Ball, E.J., Dry, P.J. and Nunez, G. (1983). The human immune response region (HLA-D) and disease susceptibility. *Immunological Rev.* 70: 111-154.

Stern, L.J., Brown, J.H., Jardetzky, T.S., Gorga, J.C., Urban, R.G., Strominger, J.L. and Wiley, D.C. (1994). Crystal structure of the human class II NHC protein HLA-DR1 complexed with an influenza virus peptide. *Nature* 368: 215-221.

Stenszky, V., Kozma, L., Baláza C.S., Rochlitz, S.Z., Bear, J.C. and Farid, N.R. (1985). The genetics of Graves' disease: HLA and disease susceptibility. *J. Clin. Endocrinol. Metab.* 61(4): 735-740.

Sujirachato, K., Chiewsilp, P., Tsuji, K., Inoko, H., Panyim, S., Mongkolsuk, T. and Leetrakool, N. (1994). HLA class II polymorphism in Thais detected by PCR-SSO and PCR-RFLP. *Tissue Antigens* 43: 224-228.

Svejgaard, A., Hauge, M., Jersild, C., Platz, P., Ryder, L.P., Staub Nielson, L. and Thomsen, M. (1975). The HLA system. An introductory survey. *Monogr. Hum. Genet.* 7

Tamai, H., Kimura, A., Dong, R., Matsubayashi, S., Kuma, K., Nagataki, S. and Sasazuki, T. (1994). Resistance to autoimmune thyroid disease is associated with HLA-DQ. *J. Clin. Endocrinol. Metab.* 78(1): 94-97.

Tamai, H., Matsubayashi, S., Fukata, S., Nagai, K., Mori, K., Hirota, Y., Kuma, K., Kumagai, L.F. and Nagataki, S. (1986). Major gene linked to HLA controls susceptibility to Japanese thyrotoxic periodic paralysis. In Vichayanrat, A., Nitayanant, W., Eastman, C. and Nagataki, S. (Ed.). *Recent progress in thyroidology* (pp.183-187), Crystall House press.

Tandon, N., Mehra, N.K., Taneja, V., Vaidya, M.C. and Kochupillai, N. (1990). HLA antigens in Asian Indian patients with Graves' disease. *Clin. Endocrinol.* 33: 21-26.

Theofilopoulos, A. (1995). The basis of autoimmunity: part II genetic predisposition *Immunol. Today* 16(3): 150-159.

Thorsby, E and Rønningten, K.S. (1992). Role of HLA genes in predisposition to develop insulin diabetes mellitus. *Ann. Med.* 24(6): 523-531.

Tiwari, J.L. and Terasaki, P.I (1985). The HLA complex. In Tiwari, J.L. and Terasaki, P. I (Ed.), *HLA and disease associations* (pp. 4-15). Springer-Verlag.

Todd, J. A., Bell, J.I. and McDevitt, H.O. (1987). HLA-DQ $\beta$  gene contributes to susceptibility and resistance to insulin-dependent diabetes mellitus. *Nature* 329(15): 599-604.

Trowsdale, J., Ragoussis, J. and Campbell, R.D. (1991). Map of the human MHC. *Immunol. Today* 12: 445.

Volpé, R. (1991). The pathophysiology of autoimmune thyroid disease. *Endocrine Regulations* 25: 187-192.

Vogel, F. and Motulsky, A.G. (1982). Major histocompatibility complex (MHC). In Vogel, F. and Motulsky, A.G., *Human genetic* (pp.131-137), Springer-Verlag.

- Wang, F.Q., Semana, G., Fauchet, R. and Genetet, B. (1993). HLA-DR and -DQ genotyping by PCR-SSO in Shanghai Chinese. *Tissue Antigens* 41: 223-226.
- Weetman, A.P. (1994). The potential Immunological role of the thyroid cell in autoimmune thyroid disease. *Thyroid* 4(4): 493-499.
- Winchester, R. (1994). The molecular basis of susceptibility to rheumatoid arthritis. *Adv. Immunol.* 56: 389-466.
- Woolf, B. (1955). On estimating the relation between blood group and disease. *Ann. Hum. Genet.* 19: 251-253.
- Wongkuttiya, D. (1994). Comparison of frequency of HLA-DQA1 and HLA-DQB1 alleles between leprosy patients and normal controls in northern Thailand. M.S. Thesis, Chiang Mai University.
- Yanagawa, T., Hidaka, Y., Guimaraes, V., Soliman, M. and DeGroot, L.J. (1995). CTLA-4 gene polymorphism associated with Graves' disease in a Caucasian population. *J. Clin. Endocrinol. Metab.* 80(1): 41-45.
- Yanagawa, T., Mangklabruks, A., Chang, Y., Okamoto, Y., Fisfalen, M., Curran, P.G. and DeGroot, L.J. (1993). Human histocompatibility leukocyte antigen-DQA1\*0501 allele associated with genetic susceptibility to Graves' disease in a Caucasian population. *J. Clin. Endocrinol. Metab.* 76(6): 1569-1574.
- Yanagawa, T., Mangklabruks, A., DeGroot, L.J. (1994). Strong association between HLA-DQA1\*0501 and Grave's disease in a male Caucasian population. *J. Clin. Endocrinol. Metab.* 79: 227-229.
- Yeo, P.P.B., Chan, S.H., Lui, K.F., Wee, G.B., Lim, P. and Cheah, J.S. (1978). HLA and thyrotoxic periodic paralysis. *Br. Med. J.* 2(6142): 930.

Yeo, P.P.B., Chan, S.H., Thai, A.C., Ng, W.Y., Lui, K.F., Wee, G.B., Tan, S.H., Lee, B.W., Wong, H.B. and Cheah, J.S. (1989). HLA-Bw46 and DR9 associations in Graves' disease of Chinese patients are age- and sex-related. *Tissue Antigens* 34: 179-184.

Yoshigawa, N., Morita, T., Resetkova, E., Arreanza, G., Carayon, P. and Volpé, R. (1992). Reduced activation of suppressor T lymphocytes by specific antigens in autoimmune thyroid disease. *J. Endocrinol. Invest.* 15: 609-617.

Zanelli, E., Gonzalez-Gay, M. and David, C.S. (1995). Could HLA-DRB1 be the protective locus in rheumatoid arthritis? *Immunol. Today* 16(6): 274-278.