

## REFERENCES

- Abel, P.D. (1989). Water pollution biology. Ellis Horwood, Chichester, England
- American Public Health Association (APHA). (1975). Ch. 1005. Benthic macroinvertebrates, in standard methods for the examination of water and waste water. Washington DC. American Public Health Association. pp. 1060-1075.
- Armitage, P.D. (1979). A flooding artificial substratum sampler for use in standing water. *Hydrobiologia*. 66: 245-248.
- Arthur, J.W., Horning, H.B. (1969). The use of artificial substrates in pollution surveys. *Am. Midi. Nat.* 82, 83-89.
- Barber, W.E., Greenwood, J.E., Crocos, P. (1979). Artificial Seagrass- A new technique for sampling the community. *Hydrobiologia* 65, 135-140.
- Beak, T.W., Griffing, T.C., Appleby, A.G. (1973). Use of artificial substrate samplers to assess water pollution. In biological methods for the assessment of water quality. ASTM spec. tech. publ. eds. J. Cairns, Jr. and Dickson, K.L. pp. 227-241.
- Bergersen, E.P., Galat, D.L. (1975). Coniferous tree bark: A lightweight substitute for limestone rock in barbeque basket macroinvertebrate samplers. *Water Res.* 9: 729-731.
- Boobar, L.R., Granett, D.L. (1978). Evaluation of polyethylene samplers for black fly larvae. *Can. J. Zool.* 56, 2245-2248.

- Chessmen, B.C. (1995). Rapid assessment of rivers using macroinvertebrates. A procedure based on habitat-specific sampling, family level identification and a biotic index. *Australian journal of Ecology*. 20, 122-129.
- Cover, E.C., Harrel, R.C. (1978). Sequences of colonization, diversity, biomass and productivity of macroinvertebrates on artificial substrates in a fresh water canal. *Hydrobiologia* 59, 81-95.
- Crossman, J.S., Cairns Jr, J. (1974). A comparative study between two different artificial substrate samplers and regular sampling techniques. *Hydrobiologia* 44, 517-522.
- Crowe, J.- A.M.E. (1974). Test of artificial samplers for collecting stream macroinvertebrates in Manitoba. *Manitoba Entomol.* 8, 19-31.
- Dickson, K.L., Cairns jr., J. (1972). The relationship of freshwater macroinvertebrate communities collected by floating artificial substrates to the MacArthur-Wilson equilibrium model. *Am.Midi. Nat.* 88: 68-75.
- Disney, R.H. (1972). Observation on sampling pre-imaginal population of blackflies (Dipt., Simuliidae) in West Cameroon. *Bull, Entomol. Res.* 61, 485-503.
- Dudgeon, D. (1984). Seasonal and long term changes in the hydrobiology of the Lam Tsuen River, New territories, Hong Kong, with special reference to benthic macroinvertebrate distribution and abundance. *Arch. Hydrobiologia. Suppl.* 69, 56-129.
- Elliott, J. (1977). Some methods for the statistical analysis of samples of benthic invertebrates. 2nd ed. *Freshwater Biol. Assoc. Sci. Publ.* 25, 1-160.

- Elliott, J.M., Dreak, C.M. (1981b). A comparative study of four dredges used for sampling benthic macroinvertebrates in rivers. *Freshwat. Biol.* 11: 245 - 261
- Flannagan, J.F., Rosenberg, D.M. (1982). Types of artificial substrates used for sampling freshwater benthic macroinvertebrates. In: Artificial Substrates. Ed. Cairns JR., J. pp. 237-265.
- Flannagan, J.F., Townsend, B.E., De March, B.G.E. (1980). Acute and long-term effects of Methoxychlor larviciding on the aquatic invertebrates of the Athabasca River, Alberta. Technical report, Alberta Department Environment, Edmonton, Alberta. pp. 151-168.
- Fredeen, F.J.H., Spurr, D.T. (1978). Collecting semi-quantitative samples of black fly larvae and other aquatic insects from large rivers with the aid of artificial substrates. *Quaest. Entomol.* 14, 411-431.
- Fullner, R.W. (1971). A comparison of macroinvertebrates collected by basket and modified multiple- plate samplers. *J. Water Poll. Control Fed.* 43, 494-499.
- Furse, M.T., Wright, J.F., Armitage, P.D., Moss, D. (1981). An appraisal of pond net samples for biological monitoring of lotic macro-invertebrates. *Water Research.* 15, 679-689.
- Gersabeck Jr, E.F., Merritt, R.W. (1979). The effect of physical factors on the colonization and relocation behavior of immature black flies. *Environ. Entomol.* 8, 34-39.
- Growns, J.E., Chessmen, B.C., McEvoy, P.K., Wright, I.A. (1995). Rapid assessment of rivers using macroinvertebrates case studies in the Nepean River and Blue Mountains, New South Wales. *Australian Journal of Ecology.* 20, 130-141.

- Hawkes, H.A (1979). Invertebrates as indicators of river water quality. In: Biological indicators of water quality, eds. James and L. Evison, Chap. 2. Jown Wiley, Chichester, England.
- Hellawell, J.M. (1986). Biological indicators of fresh water pollution and environment management. Elsevier Applied Science Publishers, London, England.
- Hellawell, J.M. (1978). Biological surveillance of rivers. A biological monitoring hand book. Water Research Center, Medmenham, England.
- Hester, F.E., Dendy, J.S. (1962). A multiple-plate sampler for aquatic macroinvertebrates. *Trans. Am. Fish. Soc.* 91, 420-421.
- Hilsenhoff, W.L. (1969). An artificial substrate device for sampling benthic stream invertebrates. *Limnol. Oceanog.* 14, 465-471.
- Hoar, M.A., Miller, M.C. (1972). Midsummer studies of a new year -round artificial substrate for macroinvertebrate population in the upper Mississippi River. M.Sc Thesis, St. Mary's College, Winona MN.
- Konstantinov, A.S. (1977). Ability of chironomid larvae to colonize substrates variously positioned in a lake. *Hydrobiol. J.* 13, 11-14
- Lehmkuhl, D.M. (1979). How to know aquatic insects. Wm.C. Brown Company, Dubuque, Iowa.
- Macan, T.T. (1977). The fauna in the vegetation of a Morland fishpond as revealed by different methods of collecting . *Hydrobiologia* 55, 3-15.
- MacCafferty, W.P. (1981). Aquatic entomology. Jones and Bartlett Publishers, Inc. Portola valley, CA.

- Markosova, R. (1980). Effect of submerged substrates quality on the periphytic macrofauna in ponds. *Vestn. Cesk. Spol. Zool.* 44, 46-52.
- Mason Jr., W.T., Weber, C.I., Lewis, P.A, Julian, E.C. (1973). Factors affecting the performance of basket and Multiplate macroinvertebrates samplers. *Freshwater Biol.* 3: 409-436.
- McBride, G.B. (1985). The role of monitoring in the management of water resources. In biological monitoring in freshwater. eds. R.D. Pridmore and A.B. Cooper, pp7-16.
- Meier, P.G., Penrose, D.L., Polak, L. (1979). The rate of colonization by macroinvertebrates on artificial substrate samplers. *Freshwater Biol.* 9, 381-392.
- Merritt, R. W., Cummins, K.W. (Eds) ( 1984). An introduction to the aquatic insects of North America, 2nd Ed. Kendall Hunt Publishing Co., Dubuque, Iowa.
- Metcalf, J.L.(1989). Biological water quality assessment of running waters based on macroinvertebrates communities. History and present status in Europe. *Environmental pollution* 60, 101-139.
- Minshall, G.W., Minshall, J. N. (1977). Microdistribution of benthic invertebrates in a rocky mountain (U.S.A) stream . *Hydrobiologia* 55, 231-249.
- Moon, H.P. (1935b). Flood movements of the littoral fauna of Windermere. *J. Animal Ecol.* 4, 216-228.
- Needham, J.G. (1962). A Guide to the study of fresh water biology. Holden Day Inc. San Francisco, CA.

- Norris, R.H., Norris, K.R. (1995). The need for biological assessment of water quality. Australian perspectives. *Australian Journal of Ecology*. 20, 1-6.
- Parsons, D.S., Tatum, J.W. (1974). A New shallow water multiple-plate sampler. *Prog.Fish.Cult.* 36, 179-180.
- Pearson, R.G., Jones, N.V. (1975). The colonization of artificial substrata by stream macroinvertebrates. *Prog. Water Technol.* 7: 497-504.
- Pennak, M. (1978). Fresh-water invertebrates of the United States. Jown Wiley and Son, Inc. New York.
- Penny, S.F. (1985). The use of macroinvertebrates in the assessment of point source pollution. In biological monitoring in freshwaters. R.D. Pridmore and A.B. Cooper, pp. 205-15.
- Peterson, R.C., Cummins, K.W. (1974). Leaf processing in a Woodland stream. *Freshwater Biol.* 4, 343-368.
- Pinder, L.C. V., Farr, I.S. (1987). Biological surveillance of water quality -3. The influence of organic enrichment on the invertebrate fauna of small Chalk streams. *Arch. Hydrobiologia.* 109, 207-226.
- Pinder, L.C.V., Ladle, M., Gledhill, T., Bass, J.A.B., Matthews, A.M. (1987). Biological surveillance of water quality -1. A comparison of macroinvertebrates surveillance methods in relation to assessment of water quality in a Chalk Stream. 109, 207-226.
- Plafkin, J.L., Barbour M.T., Porter K.D., Gross S.K., Hughes R.M. (1989). Rapid bioassessment protocols for use in streams and rivers. Benthic macroinvertebrates and fish. EPA/440/4-89/001. Office of water regulation and standards, US Environmental Protection Agency, Washington, DC.

- Rabeni, C.F., Gibbs, K.E. (1978). Comparison of two methods used by divers for sampling benthic invertebrates in deep rivers. *J. Fish. Res. Board Can.* 35, 332-336.
- Rajchapakdee, M. (1992). Benthic invertebrates relative to altitude and water quality in streams on Doi Suthep. M.S Thesis Chiang Mai University, Thailand.
- Resh, V.H.(1995). Design and implementation of rapid assessment approaches for water resources monitoring using benthic macroinvertebrates. *Australian Journal of Ecology.* 20, 108-121.
- Roby, K.B., Newbold, J.D., Erman, D.C. (1978). Effectiveness of a artificial substrate for sampling macroinvertebrates in small streams. *Freshwater Biol.* 8, 1-8.
- Rosenberg, D.M., Resh, R.H. (1993). Fresh water biomonitorng and benthic macroinvertebrates. Chapman & Hall. New York
- Rosenberg, D.M., Resh, R.H. (1982). The use of artificial substrates in the study of freshwater benthic macroinvertebrates. in: *Artificial Substrates.* ed. Cairns Jr., J. pp. 175-235.
- Rosenberg, D.M., Wiens, A.P. (1976). Community and species responses of Chironomidae (Diptera) to contamination of freshwater by crude oil and petroleum products, with special reference to the river Trail, Northwest Territories. *J. Fish. Res. Bd. Can.* 33: 1955-1963
- Sannarm, G.(1993). Biological surveillance using macroinvertebrates communities of the quality of the Mae Kwang River The Northern Region Industrial Estate. M.S. Thesis Chiang Mai University, Thailand.

- Shaw, D.W., Minshall, G.W. (1980). Colonization of an introduced substrate by stream macroinvertebrates . *Oikos* 34, 259-271.
- Suwanrat, P. (1996). Qualitative survey of macroinvertebrates in running waters to test rapid assessment technique for water quality. M.S Thesis Chiang Mai University, Thailand.
- Thaweeburus, P. (1994). Biomonitoring techniques to assess water loss at the old Mae Ping dam and water quality of the Mae Ping River. M.S Thesis Chiang Mai University, Thailand.
- Throne, R. (1993). The development and testing of a biological monitoring protocol for incorporation into GEMS/WATER system. King's College. London.
- Tuyor, J.B. (1994). Impact of highland agriculture on streams macroinvertebrates communities. M.S. Thesis Chiang Mai University, Thailand.
- Watchawong, O. (1996). Quantitative survey and habitat assessment of macroinvertebrates to assess water quality in different running waters. M.S Thesis Chiang Mai University, Thailand.
- Weber, C.I. (1973). Ed. Biological field and laboratory methods for measuring the quality of surface waters and effluents. *U.S Environmental Protection Agency, Environmental Monitoring Series*, EPA- 670/4-73-001. pp. 1-186.
- Welton, J.S., Cooling, D.A., Ladle, M. (1982). A comparison of two colonization samplers with a conventional techniques for quantitative sampling of benthic macroinvertebrates in the gravel substrates of an experimental recirculating stream. *Hydrobiologia* 67, 901-906.

Williams, D.D., Hynes, H.B.N. (1976a). The Recolonization mechanisms of Stream Benthos. *Oikos* 27. 265-272.

Wright, I.A., Chessmen, B.C., Fairweather, P.G., Benson, L. J. (1995). Measuring the impact of sewage effluent on the macroinvertebrate community of an upland stream. The effect of different levels of taxonomic resolution and quantification. *Australian Journal of Ecology*. 20, 142-149.

มหาวิทยาลัยเชียงใหม่  
Chiang Mai University