

| | | |
|------------|-------------------------------------------------------------------------------------------------------|----|
| 4.3 | Sorting and identification | 41 |
| 4.4 | Colonization curve | 42 |
| 4.5 | Data analysis | 43 |
| Chapter 5. | Results | 51 |
| 5.1 | Missing data | 51 |
| 5.2 | Available colonization area of ASS | 51 |
| 5.3 | Abundance and taxon richness of macroinvertebrates in sampling methods | 51 |
| 5.4 | Abundance of animals per site using different sampling methods.. | 52 |
| 5.5 | Coefficient of variance in ASS | 55 |
| 5.6 | Data transformation | 56 |
| 5.7 | Taxa richness in ASS | 64 |
| 5.8 | Taxonomic composition of macroinvertebrates in ASS | 66 |
| 5.9 | Taxonomic composition of macroinvertebrates in ASS According to sites..... | 70 |
| 5.10 | Artificial substrate preference by macroinvertebrates | 74 |
| 5.11 | Colonization sequence in ASS..... | 74 |
| 5.12 | Effect of increased water level and velocity on abundance of macroinvertebrates in ASS | 78 |
| 5.13 | Natural substrate preference by macroinvertebrates | 78 |
| 5.14 | Macroinvertebrates found in different substrate types and seasons according to sampling sites..... | 81 |
| 5.15 | Taxa richness according to substrate type and season | 81 |

| | |
|----------------------------------------------------------------------|-----|
| 5.16 Composition of macroinvertebrates in different substrates | 85 |
| 5.17 Evaluation of water quality using Rapid Bioassessment | |
| Protocol II (RBP II) based on macroinvertebrates found | |
| in WB sampler and conventional methods | 86 |
| 5.17.1 Application of RBP (II) for macroinvertebrates | |
| Collected from the ASS and conventional methods .. | 86 |
| 5.18 Clustering of macroinvertebrate families found in WB sampler | |
| and conventional methods | 92 |
| Chapter 6. Discussion | 99 |
| Chapter 7. Conclusions and Recommendations | 109 |
| References | 111 |
| Appendices | |
| Appendix A | 120 |
| Appendix B | 122 |
| Appendix C | 124 |
| Appendix D | 135 |
| Appendix E | 143 |
| Curriculum Vitae..... | 147 |

List of Tables

| | Page |
|----------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Table 1 | Sampling sites, altitude and corresponding codes..... 25 |
| Table 2 | Physical and chemical parameters of the study sites..... 27 |
| Table 3 | Types of sampling methods and their relevant codes 34 |
| Table 4 | Quality points range for each score..... 49 |
| Table 5 | Quality points range for the final classification of sites..... 50 |
| Table 6 | Mean number of animals per m ² , total number of families per unit sampler in each sampling methods and families common to all sampling sites 52 |
| Table 7 | Coefficient of variance in artificial substrate sampler and conventional methods for two seasons..... 55 |
| Table 8 | Values of Levene test for homogeneity of variance for untransformed data and different transformation methods calculated for independent variables..... 57 |
| Table 9 | Mean number of animals /m ² , and 95 % upper and lower confidence limits per sampling methods irrespective of sites and seasons..... 63 |
| Table 10 | Mean number of animals /unit sampler, and 95 % upper and lower confidence limits per sampling methods irrespective of sites and seasons..... 63 |
| Table 11 | Mean number of families per unit sampler, and 95 % confidence limits recorded from study sites irrespective to sampling methods and season. 66 |
| Table 12 | Mean number of animals/m ² in different sites, substrates and seasons... 80 |
| Table 13 | Results of the various indices for the wooden box sampler (dry season)..... 86 |
| Table 14 | Results of the various indices for the wooden box sampler (wet season)..... 87 |
| Table 15 | Results of the various indices for the conventional methods (dry season)..... 88 |

| | | |
|----------|-------------------------------------------------------------------------------|----|
| Table 16 | Results of the various indices for the conventional methods (wet season)..... | 88 |
| Table 17 | Water quality scores for the Wooden box sampler (dry season)..... | 89 |
| Table 18 | Water quality scores for the Wooden box sampler (wet season)..... | 90 |
| Table 19 | Water quality scores for the conventional methods (dry season)..... | 91 |
| Table 20 | Water quality scores for the conventional methods (wet season)..... | 91 |

List of Figures

| | | |
|-----------|-----------------------------------------------------------------------------------------------------------------------|----|
| Figure 1 | Map of the study sites | 26 |
| Figure 2 | Study site ST 1 in stream | 32 |
| Figure 3 | River site R 1 in the Ping River | 32 |
| Figure 4 | Sewage canal site SC1 | 33 |
| Figure 5 | Sewage canal site SC2 | 33 |
| Figure 6 | Artificial substrate samplers used in the research..... | 37 |
| Figure 7 | Assembling artificial substrate samplers as one unit..... | 38 |
| Figure 8 | Processing of retrieved artificial substrate sampler | 39 |
| Figure 9 | Abundance of macroinvertebrates in WMC sampler per site | 53 |
| Figure 10 | Abundance of macroinvertebrates in WB sampler | 53 |
| Figure 11 | Abundance of macroinvertebrates in MP sampler per site | 54 |
| Figure 12 | Abundance of macroinvertebrates in conventional methods /site .. | 54 |
| Figure 13 | Normal distribution curve for row data and log n transformed data (abundance/m ²) | 58 |
| Figure 14 | Normal distribution curve for log 10 and square root transformed data (abundance/m ²) | 59 |
| Figure 15 | Normal distribution curve for row data and log n transformed data (No: of families /unit area) | 60 |
| Figure 16 | Normal distribution curve for log 10 and square root transformed data (No: of families /unit area) | 61 |
| Figure 17 | Mean No: of animals/ m ² , upper and lower limits of 95 % confident levels, upper and lower range | 62 |
| Figure 18 | Comparison of mean number of animals /m ² in each sites in both season | 65 |
| Figure 19 | Number of families (per unit sampler) colonized in sampling methods irrespective of sites and seasons | 67 |
| Figure 20 | Taxonomic composition of macroinvertebrates in ASS in dry season | 68 |
| Figure 21 | Taxonomic composition of macroinvertebrates in ASS in wet season | 69 |

| | | |
|-----------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| Figure 22 | Taxonomic composition of macroinvertebrates in WMC sampler per site in dry and wet season | 71 |
| Figure 23 | Taxonomic composition of macroinvertebrates in WB sampler per site in dry and wet season | 72 |
| Figure 24 | Taxonomic composition of macroinvertebrates in MP sampler per site in dry and wet season | 73 |
| Figure 25 | Colonization sequence for artificial substrate samplers | 75 |
| Figure 26 | Colonization sequences in replicate WMC samplers | 76 |
| Figure 27 | Colonization sequences in replicate WB samplers | 77 |
| Figure 28 | Colonization sequences in replicate MP samplers | 77 |
| Figure 29 | Effect of water level and velocity on colonization of macroinvertebrates in WMC and WB sampler | 79 |
| Figure 30 | Effect of water level and velocity on colonization of macroinvertebrates in MP sampler | 80 |
| Figure 31 | Abundance of macroinvertebrates in different substrate types in study sites, mean value, upper and lower limits of 95 % confident limits , upper and lower limits | 82 |
| Figure 32 | Taxa richness in different substrate types | 83 |
| Figure 33 | Taxa richness in different seasons | 83 |
| Figure 34 | Taxonomic composition of macroinvertebrates in different substrates according to seasons | 84 |
| Figure 35 | Taxonomic composition of macroinvertebrates in different in cool season | 85 |
| Figure 36 | Clustering of macroinvertebrates families found in WB sampler in dry season | 93 |
| Figure 37 | Clustering of macroinvertebrates families found in WB sampler in wet season | 94 |
| Figure 38 | Clustering of macroinvertebrates families found in conventional methods in dry season | 95 |
| Figure 39 | Clustering of macroinvertebrates families found in conventional methods in wet season | 96 |

| | | |
|-----------|---------------------------------------------------------------------------------------------------------------------------------|----|
| Figure 40 | Clustering of study sites according to similarity of macroinvertebrates families found in WB sampler- dry season | 97 |
| Figure 41 | Clustering of study sites according to similarity of macroinvertebrates families found in WB sampler- wet season | 98 |
| Figure 42 | Clustering of study sites according to similarity of macroinvertebrates families found in conventional methods dry season | 98 |
| Figure 43 | Clustering of study sites according to similarity of macroinvertebrates families found in conventional methods wet season | 98 |

มหาวิทยาลัยเชียงใหม่
Chiang Mai University