

REFERENCES

- Ausubel, F.M., Brent, R., Kingston, R.E., Moore, D.D., Seidman, J.G., Smith, J.A. and Struhl, K. (1995). *Current Protocols in Molecular Biology*, volume 1. John Wiley & Sons. Inc., Massachusetts, pp. 2.10.1.
- Bagnarelli, P., Menzo, S., Manzin, A., Giacca, M., Varaldo, P.E. and Clementi, M. (1991). Detection of human immunodeficiency virus type 1 genomic RNA in plasma samples by reverse transcription polymerase chain reaction. *J. Med. Virol.* 34: 89-95.
- Blok, J. (1985). Genetic relationships of the dengue virus serotypes. *J. Gen. Virol.* 66: 1323-1325.
- Blok, J., Gibbs, A.J., McWilliam, S.M. and Vitarana, U.T. (1991). NS1 gene sequences from eight dengue-2 viruses and their evolutionary relationships with other dengue-2 viruses. *Arch. Virol.* 118: 209-223.
- Blok, J., McWilliam, S.M., Butler, H.C., Gibbs, A.J., Weiller, G., Herring, B.L., Hemsley, A.C., Aaskov, J.G., Yoksan, S. and Bhamarapravati, N. (1992). Comparison of a dengue-2 virus and its candidate vaccine derivative: sequence relationships with the flaviviruses and other viruses. *Virology.* 187: 573-590.

- Blok, J., Samuel, S., Gibbs, A.J. and Vitaratana, U.T. (1989). Variation of the nucleotide and encoded amino acid sequences of the envelope gene from eight dengue-2 viruses. *Arch. Virol.* 105: 39-53.
- Brinton, M.A., Fernandez, A.V. and Dispoto, J.H. (1986). The 3'-nucleotides of flavivirus genomic RNA form a conserved secondary structure. *Virology.* 153: 113-121.
- Burke, D.S., Nisalak, A., Johnson, D.E. and Scott, R.M. (1988). A prospective study of dengue infections in Bangkok. *Am. J. Trop. Med Hyg.* 38: 172-180.
- Cardiff, R.D. and Lund, J.K. (1976). Distribution of dengue 2 antigens by electron immunocytochemistry. *Infect. Immun.* 3: 1699-1709.
- Chungue, E., Deubel, V., Cassar, O., Laille, M. and Martin, P.M.V. (1993). Molecular epidemiology of dengue 3 viruses and genetic relatedness among dengue 3 strains isolated from patients with mild or severe form of dengue fever in French Polynesia. *J. Gen. Virol.* 74: 2765-2770.
- Duangchanda, S., Tanaka, M., Morita, K., Rojanasuphot, S. and Igarashi, A. (1994). Comparative nucleotide and deduced amino acid sequence of the envelope glycoprotein gene among three dengue virus type 2 strains isolated from patients with different disease severities in Maha Sarakham, Northeast Thailand. *Southeast Asian J. Trop. Med. Pub. Hlth.* 25: 243-251.

Feighny, R., Burrous, J., McCown, J., Hoke, C. and Putnak, R. (1992). Purification of native dengue-2 viral proteins and the ability of purified proteins to protect mice. *Am. J. Trop. Med. Hyg.* 47(4): 405-412.

Fu, J., Tan, B.H., Yap, E.H., Chan, Y.C. and Tan, Y.H. (1992). Full-length cDNA sequence of dengue type 1 virus (Singapore strain S275/90). *Virology*. 188: 953-958.

Gubler, D.J. and Trent, D.W. (1994). Emergence of epidemic dengue/dengue hemorrhagic fever as a public health problem in the Americas. *Infectious Agent and Disease*. 2 (6): 383-393.

Hahn, Y.S., Galler, R., Hunkapiller, T., Dalrymple, J.M., Strauss, J.H. and Strauss, E.G. (1988). Nucleotides sequence of dengue 2 RNA and comparison of the encoded proteins with those of other flaviviruses. *Virology*. 162: 167-180.

Haishi, S., Tanaka, M. and Igarashi, A. (1990). Comparative amino acid sequences of dengue viruses. *Trop. Med.* 32(2): 81-87.

Halstead, S.B. (1988). Pathogenesis of dengue: Challenges to molecular biology. *Science*. 239: 476-481.

Halstead, S.B. (1992). The XXth century dengue pandemic: need for surveillance and research. *World Health Stat Q.* 45: 292-298.

Heinz, F.X., Mandl, C., Holzmann, H., Guirakhoo, F., Tuma, W. and Kunz, C. (1990). Antigenic structure and function of the flavivirus envelope protein E. In New aspects of positive-strand RNA viruses. American Society for Microbiology. Washington, D.C.: 294-300.

Henchal, E.A., Gentry, M.K., McCown, J.M. and Brandt, W.E. (1982). Dengue virus-specific and flavivirus group determinants identified with monoclonal antibodies by indirect immunofluorescence. *Am. J. Trop. Med. Hyg.* 31(4): 830-836.

Innis, M.A., Gelfand, D.H., Sninsky, J.J. and White, T.J. (1990). PCR Protocols: A guide to methods and applications. Academic Press, Inc., San Diego, California: 9-10.

Jacob, K., Ruderdorf, R., Neil, S.D., Dougherty, J.P., Brown, E.L. and Fritsch, E.F. (1988). The thermal stability of oligonucleotide duplex is sequence independent in Tetraalkylammonium solution: application to identifying recombinant DNA clones. *Nucleic Acids Res.* 16(10): 4637-4650.

Jatanasen, S. and Thongcharoen, P. (1993). Dengue haemorrhagic fever in South-East Asia countries. In Monograph on dengue/dengue haemorrhagic fever. World Health Organization: 23-30.

Kaufman, B.M., Summers, P.L., Dubois, D.R. and Eckels, K.H. (1987). Monoclonal antibodies against dengue 2 virus E-glycoprotein protect mice against lethal dengue infection. *Am. J. Trop. Med. Hyg.* 36(2): 427-434.

Kerschner, J.A.H., Vorndam, A.V., Monath, T.P. and Trent, D.W. (1986). Genetic and epidemiological studies of dengue type 2 viruses by hybridization using synthetic deoxyoligonucleotides as probes. *J. Gen. Virol.* 67: 2645-2661.

Kimura, A. and Sasazuki, T. (1991). Eleventh International Histocompatibility Workshop reference protocol for the HLA DNA-typing technique, W3.3, Yokohama Japan. November: 6-13.

Kliks, S.C., Nisalak, A., Brandt, W.E., Wahl, L. and Burke, D.S. (1989). Antibody-dependent enhancement of dengue virus growth in human monocytes as a risk factor for dengue hemorrhagic fever. *Am. J. Trop. Med. Hyg.* 40(4): 444-451.

Lanciotti, R.S., Lewis, J.G., Gubler, D.J. and Trent, D.W. (1994). Molecular evolution and epidemiology of dengue-3 viruses. *J. Gen. Virol.* 75: 65-75.

Lewis, J.G., Chang, G.J., Lanciotti, R.S. and Trent, D.W. (1992). Direct sequencing of large flavivirus PCR products for analysis of genome variation and molecular epidemiological investigations. *J. Virol. Methods.* 38: 11-24.

Lewis, J.A., Chang, G.J., Lanciotti, R.S., Kinney, R.M., Mayer, L.W. and Trent, D.W. (1993). Phylogenetic relationships of dengue-2 viruses. *Virology.* 197: 216-224.

Mackenzie, J.M., Jones, M.K. and Young, P.R. (1996). Immunolocalization of the dengue virus nonstructural glycoprotein NS1 suggests a role in viral RNA replication. *Virology.* 220: 232-240.

- Mackow, E., Makino, Y., Zhao, B., Zhang, Y., Markoff, L., Buckler-White, A., Guiler, M., Chanock, R. and Lai, C. (1987). The nucleotide sequence of dengue type 4 virus: Analysis of gene coding for non-structural proteins. *Virology*. 159: 217-228.
- Mandl, C.W., Guirakhoo, F., Holzmann, H., Heinz, F.X. and Kunz, C. (1989). Antigenic structure of the flavivirus envelope protein E at the molecular level, using tick-borne encephalitis virus as a model. *J. Virol.* 63: 564-571.
- Monath, T.P., Wands, J.R., Hill, L.J., Brown, N.V., Marciniak, R.A., Wong, M.A., Gentry, M.K., Burke, D.S., Grant, J.A. and Trent, D.W. (1986). Geographic classification of dengue-2 virus strains by antigen signature analysis. *Virology*. 154: 313-324.
- Nimmannitya, S. (1987). Dengue haemorrhagic fever in Thailand. *Southeast Asian J. Trop. Med. Pub. Hlth.* 18(3): 291-294.
- Nimmannitya, S. (1993). Clinical manifestations of dengue/dengue haemorrhagic fever. In *Monograph on dengue/dengue haemorrhagic fever*. World Health Organization: 48-54.
- Nisalak, A., Nimmannitaya, S. and Innis, B. (1990). Severe dengue infections among children in metropolitan Bangkok, 1973-1989: description of epidemics and characterization of virus serotypes associated with them. *Southeast Asian J. Trop. Med. Hyg. Pub. Hlth.* 21: 678-679.

Osatomi, K. and Sumiyoshi, H. (1990). Complete nucleotide sequence of dengue type 3 virus genome RNA. *Virology*. 176: 643-647.

Pang, T. (1987). Pathogenesis of dengue haemorrhagic fever: towards a more balanced view. *Southeast Asian J. Trop. Med. Pub. Hlth*. 18(3): 321-325.

Preugschat, F., Yao, C.W. and Strauss, J.H. (1990). In Vitro processing of dengue virus type 2 nonstructural proteins NS2A, NS2B, and NS3. *J. Virol*. 64: 4364-4374.

Pryor, M.J. and Wright, P.J. (1993). The effects of site-directed mutagenesis on the dimerization and secretion of the NS1 protein specified by dengue virus. *Virology*. 194: 769-780.

Putnak, J.R. and Henchal, E.A. (1990). The dengue viruses. *Clin. Micro. Rev*. 3: 376-396.

Quintos, F.N., Lim, L.E., Juliano, L., Reyes, A. and Lacson, P. (1954). Haemorrhagic fever observed among children in the Philippines. *Phil. J. Pedriat*. 3: 1-19.

Repik, P.M., Dalrymple, J.M., Brandt, W.E., McCown, J.M. and Russell, P.K. (1983). RNA fingerprinting as a method for distinguishing dengue 1 virus strains. *Am. J. Trop. Med. Hyg*. 32(3): 577-589.

Rey, F.A., Heinz, F.X., Mandl, C., Kunz, C. and Harrison, S.C. (1995). The envelope glycoprotein from tick-borne encephalitis virus at 2 Å^o resolution. *Nature*. 375: 291-298.

Rice, C.M. (1996). Flaviviridae: The viruses and their replication. In: Fields, D.M., Knipe, D.M., Howley, P.M., eds. *Fields Virology*. Lippincott-Raven Publishers, Philadelphia: 931-959.

Rico-Hesse, R. (1990). Molecular evolution and distribution of dengue viruses type 1 and 2 in nature. *Virology*. 174: 479-493.

Rosen, L. (1977). The emperor's new clothes revisited, or reflections on the pathogenesis of dengue hemorrhagic fever. *Am. J. Trop. Med. Hyg.* 26: 337-343.

Russell, P.K. and Nisalak, A. (1967). Dengue virus identification by the plaque reduction neutralization test. *J. Immunol.* 99: 291-296.

Sambrook J, Fritsch E.F. and Maniatis T. (1989). *Molecular Cloning: A Laboratory Manual*. Second edition. Cold Spring Harbor Laboratory Press. New York: 11.7-10, 11.45-55.

Samuel, S., Koh, C.L., Blok, J., Pang, T. and Lam, S.K. (1989). Nucleotide and amino acid sequences of the envelope protein gene of a Malaysian dengue-2 virus isolated from a patient with dengue fever or DSS (two separate reports). *Nucleic. Acids. Res.* 17: 8887 and 8888.

Sistayanarain, A. (1993). Base sequence variation of the envelope gene of dengue viruses isolated from dengue hemorrhagic fever patients with different grades of severity. M.S. Thesis, Chiang Mai University.

Sistayanarain, A., Maneekarn, N., Polprasert, B., Sirisanthana, V., Makino, Y., Fukunaga, T. and Sittisombut, N. (1996). Primary sequence of the envelope glycoprotein of a dengue, type 2 virus isolated from patient with dengue hemorrhagic fever and encephalopathy. *Southeast Asian J. Trop. Med. Pub. Hlth.* 27(2): 221-227.

Sittisombut, N., Sistayanarain, A., Cardoso, M.J., Salminen, M., Damrongdachakul, S., Kalayanaroj, S., Rojanasuphot, S., Supawadee, J. and Maneekarn, N. (1997). Possible occurrence of a genetic bottleneck in dengue serotype 2 viruses between the 1980 and 1987 epidemic seasons in Bangkok, Thailand. *Am. J. Trop. Med. Hyg.* 56(6): 100-108.

Tan, B.H., Fu, J., Sugrue, R.J., Yap, E.H., Chan, Y.C. and Tan, Y.H. (1996). Recombinant dengue type 1 virus NS5 protein expressed in *Escherichia coli* exhibit RNA-dependent RNA polymerase activity. *Virology.* 216: 317-325.

Thant, K.Z., Morita, K. and Igarashi, A. (1995). Sequences of E/NS1 gene junction from four dengue-2 viruses of Northeastern Thailand and their evolutionary relationships with other dengue-2 viruses. *Microbiol. Immunol.* 39(8): 581-590.

Thant, K.Z., Morita, K. and Igarashi, A. (1996). Detection of the disease severity-related molecular differences among new Thai dengue-2 isolates in 1993, based on their structural proteins and major non-structural protein NS1 sequences. *Microbiol. Immunol.* 40(3): 205-216.

Thongcharoen, P. and Jatanasen, S. (1993). Epidemiology of dengue and dengue haemorrhagic fever. In Monograph on dengue/dengue haemorrhagic fever. World Health Organization: 1-8.

Trent, D.W., Grant, J.A., Monath, T.P., Manske, C.L., Corina, M. and Fox, G.E. (1989). Genetic variation and microevolution of dengue 2 virus in Southeast Asia. *Virology*. 172: 523-535.

Trent, D.W., Grant, J.A., Rosen, L. and Monath, T.P. (1983). Genetic variation among dengue 2 viruses of different geographic origin. *Virology*. 128: 271-284.

Ungchusak, K. (1987). Review of DHF in Thailand. *Weekly Epidemiological Surveillance Report*. 19(23): 127-129.

Ungchusak, K. and Kunasol, P. (1988). DHF in Thailand, 1987: The worsening situation in 30 years. Division of Epidemiology, Ministry of Public health.

Vaughn, D.W., Green, S., Kalayanarooi, S., Innis, B.L., Nimmannitya, S., Suntayakorn, S., Rothman, A.L., Ennis, F.A. and Nisalak, A. (1996). Peak dengue 2 virus titer early in secondary infection correlates with plasma leakage. In program and abstracts of the 45th annual meeting of the American Society of Tropical Medicine and Hygiene. *Am. J. Trop. Med. Hyg.* 55(2): 139-140.

Vezza, A.C., Rosen, L., Repik, P. Dalrymple, J. and Bishop, H.L. (1980). Characterization of the viral RNA species of prototype dengue viruses. *Am. J. Trop. Med. Hyg.* 29(4): 643-652.

Viputtigul, K. (1996). Determination of the association between HLA-DQA1, HLA-DQB1 and HLA-DRB1 loci with graves' disease in the Northern Thai population. M.S. Thesis, Chiang Mai University.

Walker, P.J., Henchal, E.A., Blok, J., Repik, P.M., Henchal, L.S., Burke, D.S., Robbins, S.J. and Gorman, B.M. (1988). Variation in dengue type 2 viruses isolated in Bangkok during 1980. *J. Gen. Virol.* 69: 591-602.

Wengler, G. and Wengler, G. (1991). The carboxy- terminal part of the NS 3 protein of the West Nile Flavivirus can be isolated as a soluble protein after proteolytic cleavage and represents an RNA- stimulated NTPase. *Virology.* 184: 707-715.

Winkler, G., Maxwell, S.E., Ruemmler, C. and Stollar, V. (1989). Newly synthesized dengue-2 virus nonstructural protein NS1 is a soluble protein but becomes partially hydrophobic and membrane-associated after dimerization. *Virology.* 171: 302-305.

Winkler, G., Randolph, V.B., Cleaves, G.R. Ryan, T.E. and Stollar, V. (1988). Evidence that the mature form of the flavivirus nonstructural protein NS1 is a dimer. *Virology.* 162: 187-196.

Wood, W.I., Gitschier, J., Lasky, L.A. and Lawn, R.M. (1985). Base composition-independent hybridization in tetramethylammonium chloride: A method for oligonucleotide screening of highly complex gene libraries. *Proc. Natl. Acad. Sci. USA.* 82: 1585-1588.

World Health Organization. (1986). Dengue haemorrhagic fever: diagnosis, treatment and control. WHO Press. Geneva: 7-15.

Yang, P.Y., Kautner, I., Koh, C.L. and Lam, S.K. (1994). Nucleotide and deduced amino acid sequences of genes encoding the structural and nonstructural NS1 proteins of a dengue-2 virus isolated in China. *Virus Gene.* 8: 71-74.

Yenchitsomanus, P., Sricharoen, P., Jaruthasana, I., Pattanakitsakul, S., Nitayaphan, S., Mongkolsapaya, J. and Malasit, P. (1996). Rapid detection and identification of dengue viruses by polymerase chain reaction (PCR). *Southeast Asian J. Trop. Med. Pub. Hlth.* 27(2): 228-236.

Zhao, B., Mackow, E., Buckler-White, A., Markoff, L., Chanock, R.M., Lai, C. and Makino, Y. (1986). Cloning full-length dengue type 4 viral DNA sequences: Analysis of gene coding for structural proteins. *Virology.* 155: 77-88.