

REFERENCES

- Abe, J.J., Forrester, T., Nakahara, J., Lafferty, A., Kotzin, B.L., and Leung, D.Y.M.(1991). Selective stimulation of human T cells with streptococcal erythrogenic toxin A and B. **J. Immunol.**, 146: 3747-3750.
- Agarwal, B.L.(1988). Rheumatic fever and rheumatic heart disease. Bangalore, India: India Arnold Publishers.
- Arbeit, R.D. (1995). Laboratory procedures for the epidemiologic analysis of microorganisms. in Murray, P.R., Baron, E.J., Tenover, F.C., and Tenover, F.C., and Tenover, F.C. (eds), *Manual of clinical microbiology*, 6th ed (pp. 190-208),Washington: American Society for Microbiology.
- Beachey, E.H., (1981). Bacterial adherence: adhesin-receptor interactions mediating the attachment of bacteria to mucosal surface. **J. Infect. Dis.**, 143: 325-345.
- Beachey, E.H., and Ofek, I. (1976). Epithelium cell binding of group A streptococci by lipoteichoic acid on fimbriae denuded of M protein. **J. Exp. Med.**, 43: 759-771.
- Beachey, E.H., Campbell, G.L., and Ofek, I. (1974). Peptic digestion of streptococcal M protein. II. Extraction of M antigen from group A streptococci with pepsin. **Infect. Immun.**, 9: 891-896.
- Beachey, E.H., Simpson, W.A., Ofek, I., Hasty, D.L., Dale, J.B., and Whitnack, E.(1983). Attachment of *Streptococcus pyogenes* to mammalian cells. **Review. Infect. Dis.**, 5: 5670-5677.
- Beachey, E.H., Tartar, A., Seyer, J.M., and Chedid, L. (1984). Epitope-specific protective immunogenicity of chemically synthesised 13-, 18-, and 23-residue peptide fragments of streptococcal M protein. **Proc. Natl. Acad. Sci.USA.**, 81: 2203-2207.

- Belani, K., Schlievert, P.M., Kaplan, E.L., and Ferrieri, P. (1991). Association of exotoxin-producing group A streptococci and severe disease in children. **Ped. Infect. Dis. J.**, 10: 351-354.
- Bernheimer, A.W. (1972). Hemolysin of streptococci: Characterization and effects on biological membrane, in L.W. Wannamaker and J.M. Matsen, (eds), *Streptococci and streptococcal disease* (pp. 19-29), New York and London: Academic Press.
- Bert, F., Picard, B., Lambert-Zechovsky, N., and Goullet, P. (1995). Identification and typing of pyogenic streptococci by enzyme electrophoretic polymorphism. **J. Clin. Microbiol.**, 42: 442-451.
- Bessen, D.E., and Fischetti, V.A. (1990). A human IgG receptor of group A streptococci is associated with tissue site of infection and streptococcal class. **J. Infect. Dis.**, 161: 747-754.
- Bessen, D.E., Jones, K.E., and Fischetti, V.A. (1989). Evidence for two distinct classes of streptococcal M protein and their relationship to rheumatic fever. **J. Exp. Med.**, 169: 269-283.
- Billi, D.C., Auckenthaler, R., Paunier, L., and Ferrier, C.E. (1984). Throat cultures for group A β -hemolytic streptococcus. **Am. J. Dis. Child.**, 138:274
- Bingen, E., Denamur, E., Lambert-Zechovsky, N., Braimi, N., Ellakany, M., and Elion, J. (1992a). DNA restriction fragment length polymorphism differentiates recurrence from relapse in treatment failures of *Streptococcus pyogenes* pharyngitis. **J. Med. Microbiol.**, 37: 162-164.
- Bingen, E., Denamur, E., Lambert-Zechovsky, N., Boissinot, C., Brahimi, N., Aujard, Y., Blot, P., and Elion, J. (1992b). Mother-to-infant vertical transmission and cross-colonization of *Streptococcus pyogenes* confirmed by DNA restriction fragment length polymorphism analysis. **J. Infect. Dis.**, 165: 147-150.

- Bisno, A.L. (1979). Alternative complement pathway activation by group A streptococci: role of M protein. **Infect. Immun.**, 26: 1172-1176.
- Bisno, A.L. (1991). Group A streptococcal infections and acute rheumatic fever. **New. Eng. J. Med.**, 325: 763-793.
- Bjorvatn, B., Lund, V., Kristiansen, B., Korsnes, L., Spanne, O., Lindqvist, B. (1984). Applications of restriction endonuclease fingerprinting of chromosomal DNA of *Neisseria meningitidis*. **J. Clin. Microbiol.**, 19: 763-765.
- Blumberg, H.M., Stephens, D.S., Licitra, C., Pigott, N., Facklam, R., Swaminathan, B., and Wachsmuth, I.K. (1992). Molecular epidemiology of group B streptococcal infections: use of restriction endonuclease analysis of chromosomal DNA and DNA restriction fragment length polymorphism of ribosomal RNA genes (Ribotyping). **J. Infect. Dis.**, 166: 574-579.
- Bouvet, A., Geslin, P., Kriz-Kuzemenska, P., Blanc, V., Devine, C., and Grimont, F. (1994). Restricted association between biotypes and serotypes within group A streptococci. **J. Clin. Microbiol.**, 32: 1312-1317.
- Briston, C.C. (1965). Structural, function, synthetic and genetic control of bacterial pili and molecular model for DNA and RNA transports in Gram negative bacteria. **Trans. N.Y. Acad. Sci.**, 27: 1003.
- Bronze, M.S., and Dale, J.B. (1993). Epitopes of streptococcal M protein that evoked antibodies that cross-react with human brain. **J. Immunol.**, 151: 2820-2828.
- Bronze, M.S., Beachey, E.H., and Dale, J.B. (1987). Protective and heart cross-reactive epitopes located within the NH₂-terminus of type 19 M protein. **J. Exp. Med.**, 163: 1191-1202.

- Bruneau, S., DeMontclos, H., Drouet, E., and Denoyel, G.A. (1994). rDNA gene restriction patterns of *Streptococcus pyogenes*: Epidemiological applications and relation to serotypes. **J. Clin. Microbiol.**, 32: 2953-2958.
- Chandler, K.F., Razin, S., Stephens, E.B., Harasawa, R., Barile, M.F. (1982). Genomic and phenotypic analyses of *Mycoplasma pneumoniae* strains. **Infect. Immun.**, 38: 604-9.
- Chen, C., and Cleary, P. (1990). Complete nucleotide sequence of the streptococcal C5a peptidase gene of *Streptococcus pyogenes*. **J. Biol. Chem.**, 265: 3161-3167.
- Christensen, P., and Oxelius, V.A. (1975). A reaction between some streptococci and IgA myeloma proteins. **Acta. Patho. Microbiol. Scand. Sect.**, 83:184-188.
- Cleary, P., Johnson, D., Wannamaker, L. (1979). Genetic variation in the M antigen of group A streptococci reassortment of type-specific markers and possible antigenic drift. **J. Infect. Dis.**, 140:747-57.
- Cleary, P.P., Kaplan, E.L., Handley, J.P. et al. (1992). Clonal basis for resurgence of serious *Streptococcus pyogenes* disease in the 1980s. **Lancet.**, 339: 518-521.
- Cleary, P.P., Kaplan, E.L., Lindahl, C., and Skjold, S. (1988). DNA fingerprints of *Streptococcus pyogenes* are M type specific. **J. Infect. Dis.**, 158: 1317-1322.
- Cleary, P.P., Peterson, J., Chen, C., and Nelson, C. (1991). Virulent human strains of group G streptococci express a C5a peptidase enzyme similar to that produced by group A streptococci. **Infect. Immun.**, 59: 2305-2310.
- Cunningham, M.W., and Beachey, E.H. (1974). Peptic digestion of streptococcal M protein. I. Effect of digestion at suboptimal pH upon the biology and immunological properties of purified M protein extract. **Infect. Immun.**, 9: 244-248.

- Cunningham, M.W., McCormack, J.M., Fenderson, P.G., Ho, M.K., Beachey, E.H., and Dale, J.B.(1989). Human and murine antibodies cross-reactive with streptococcal M protein and myosin recognize the sequence Gln-Lys-Ser-Lys-Gln in M protein. **J. Immunol.**, 143: 2677-2683.
- Dale, J.B. and Beachey, E.H. (1982). Protective antigenic determinant of streptococcal M protein shared with sarcolemmal membrane protein of human heart. **J. Exp. Med.**, 156: 1165-1176.
- Dale, J.B., and Beachey, E.H. (1985). Epitopes of streptococcal M proteins shared with cardiac myosin. **J. Exp. Med.**, 162: 583-591.
- Dale, J.B., and Beachey, E.H.(1986). Sequence of myosin-cross reactive epitopes of streptococcal M proteins. **J. Exp. Med.**, 164: 1785-1790.
- Denning, D.W., Baker, C.J., Troup, N.T., and Tompkins, L.S. (1989). Restriction endonuclease analysis of human and bovine group B streptococci for epidemiologic study. **J. Clin. Microbiol.**, 27: 1352-1356.
- Efstratiou, A. (1980). Preparation of *Streptococcus pyogenes* suspensions for typing by the agglutination method. **Med. Lab. Sci.**, 37: 361-363.
- Ellen, R.P., and Gibbons, R.J. (1972). M protein associated adherence of *Streptococcus pyogenes* to epithelial surfaces: prerequisite for virulence. **Infect. Immun.**, 5: 826-830.
- Facklam, R.R.(1976). A review of the microbiological techniques for isolation and identification of streptococci. **Crit. Rev. Clin. Lab. Sci.**, 6:287.
- Facklam, R.R., Padula, S.F., Wortham, E.C., Cooksey, C.C., and Rountee, H.A. (1979). Presumptive identification of group A, B and D streptococci on agar plate media. **J. Clin. Microbiol.**, 9: 665
- Falk, E.S., Danielsson, D., Bjorvatn, B., Melby, K., Sorensen, B., Kristiansen, B-E, Lund, S., Sandstrom, E.(1985). Phenotypic and genotypic characterization of penicillinase-producing strains of *Neisseria gonorrhoeae*. **Acta. Pathol. Microbiol. Immun. Scand. (B)**, 93: 91-97.

- Fast, O.J., Schlievert, P.M., and Nelson, R.D. (1989). Toxic shock syndrome-associated staphylococcal and streptococcal pyrogenic exotoxins are potent inducers of tumor necrosis factor production. **Infect. Immun.**, 57: 291-294.
- Fischetti, V.A. (1977). Streptococcal M protein extracted by nonionic detergent. II. Analysis of the antibody response to the multiple antigenic determinants of the M-protein molecule. **J. Exp. Med.**, 146: 1108-1123.
- Fischetti, V.A. (1989). Streptococcal M protein. **Clin. Microbiol. Rev.**, 2: 286-309.
- Fischetti, V.A. (1991). Streptococcal M protein. **Scientific American**, June: 32-39.
- Fischetti, V.A., and Manjula, B.N. (1982). Biologic and immunologic implications of the structural relationship between streptococcal M protein and mammalian tropomyosin. in J. B. Robins, J.C. Hill., and J.C. Sadoff (Ed.), *Seminars in infectious disease*, vol. 4. Bacterial vaccines. Thieme (pp. 411-418), New York: Stratton.
- Fischetti, V.A., Jones, K.F., and Scott, J.R. (1985). Size variation of the M protein in group A streptococci. **J. Exp. Med.**, 161: 1384-1401.
- Fischetti, V.A., Gotschlich, E.C., Siviglia, G., and Zabriskie, J.B. (1976). Streptococcal M protein extracted by non-ionic detergent. I. Properties of the antiphagocytic and type specific molecules. **J. Exp. Med.**, 144: 32-53.
- Fischetti, V.A., Jarymowycz, M., Jones, K.F. and Scott, J.R. (1986). Streptococcal M protein size mutants occur at high frequency within single strain. **J. Exp. Med.**, 164: 971-980.
- Fleischer, B., Gerardy-Schahn, R., Mextroth, B., Carrel, S., Gerlach, D., and Kohler, W. (1991). An evolutionary conserved mechanism of T cells activation by microbial toxins: evidence for different affinities of T cell receptor-toxin interaction. **J. Immunol.**, 146: 11-17.

- Fox, E.N. (1974). M proteins of group A streptococci. **Bacteriol Rev.**, 38: 57-86.
- Fox, E.N., and Wittner, M.K. (1969). New observations on the structure and antigenicity of the M proteins of the group A streptococcus. **Immunochemistry.**, 6:11-24.
- Friedman, J.M., Van de-Rijn, I., Ohkuni, H., Fischetti, V.A., and Zabriskie, J.B.(1984). Immunological studies of post-streptococcal sequelae: evidence for presence of streptococcal antigen in circulating immune complex. **J. Clin. Invest.**, 74: 1027-1034.
- Gardiner, D., Hartas, J., Currie, B., Mathews, J.D., Kemp, D.J., and Sriprakash, K.S. (1995). *Vir*-Typing: A long-PCR typing method for group A streptococci. **PCR. Method. App.**, 4: 288-293.
- Gomi, H., Hozumi, T., Hattori, S., Tagawa, C., Kishimoto, F., and Bjorek, L. (1990). The gene sequence and some properties of protein H: a novel IgG binding protein. **J. Immunol.**, 144:4046-4052.
- Gordillo, M.E., Singh, K.V., Baker, C.J., and Murray, B.E. (1993). Typing of group B streptococci: comparison of pulse field gel electrophoresis and conventional electrophoresis. **J. Clin. Microbiol.**, 31: 1430-1434.
- Gordis, L. (1985). The virtual disappearance of rheumatic fever in the United States: lessons in the rise and fall of disease. **Circulation.**, 72: 1155-1162.
- Goroncy-Bernes, P., Dale, J.B., Beachey, E.H., and Opferkuch, W. (1987). Monoclonal antibody to human renal glomeruli cross-react with streptococcal M protein. **Infect. Immun.**, 55: 2416-2419.
- Goshorn, S.C., and Schlievert, P.M.(1989). Bacteriophage association of streptococcal pyrogenic exotoxin type C. **J. Bacteriol.**, 171: 3968-3073.
- Guyer, RL., and Koshland, Jr., D.F. (1989). The molecule of the year. **Science.**, 246:1543-1544

- Haanes, E.J., and Cleary, P.P. (1989). Identification of divergent M protein gene and M protein related gene family in *Streptococcus pyogenes* serotype 49. **J. Bacteriol.**, 171: 6397-6408.
- Haanes, E.J., Heath, D.G., Cleary, P.P. (1992). Architecture of the *Vir* regulons of group A streptococci parallels opacity factor phenotype and the M protein class. **J. Bacteriol.**, 174: 4967-4976.
- Haanes-Fritz, E., Kraus, W., Burdett, V., Dale, J.B., and Beachey, E.H. (1988). Comparison of the leader sequence of four group A streptococcal M protein genes. **Nucleic. Acid. Res.**, 16: 4667-4677.
- Haase, A.M., Melder, A., Mathews, J.D., Kemp, D.J., and Adams, M. (1994). Clonal diversity of *Streptococcus pyogenes*. Associated with acute poststreptococcal glomerulonephritis and rheumatic fever. **Epidemiol. Infect.**, 113: 455-462.
- Hallas, G., and Widdowson, J.P. (1979). The nature of streptococcal opacity factor and its relationship with M protein. in Parker, M.T.(ed.), Pathogenic streptococci (pp 36-37), England: Reedbook Ltd Surrey.
- Harbaugh, M.P., Podbielski, A., Hugl, S., and Cleary, P.P.(1993). Nucleotide substitutions and small-scale insertion produce size and antigenic variation in group A streptococcal M1 protein. **Molecular Microbiology.**, 8: 981-991.
- Hauser, A.R., Goshom, S.C., Kaplan, E.L., Stevens, D.L., and Schlievert, P.M. (1991). Molecular analysis of the streptococcal pyogenic exotoxin. in. Dunny, G.M. Cleary, P.P., and McKay, L.L.(eds.), Genetics and molecular biology of the streptococci, lactococci, and enterococci (pp 195-200), Washington, D.C: England American Society for Microbiology.
- Heath, D.G., and Cleary, P.P. (1989). Fc-receptor and M protein of group A streptococci are products of gene duplication. **Proc. Natl. Acad. Sci. USA.**, 89: 4791-4745.

- Hensler, T., Koller, M., Geoffroy, C., Alouf, J., and Koning, W. (1991). *Staphylococcus aureus* Toxic shock syndrome toxin I and *Streptococcus pyogenes* erythrogenic toxin A modulate inflammatory mediator release from human neutrophils. **Infect. Immun.**, 61: 1055-1061.
- Herman, A., Kappler, J.W., Marrack, P., and Pullen, A.M. (1991). Superantigens: mechanism of T-cell stimulation and role in immune response. **Ann. Rev. Immunol.**, 9: 745-772.
- Hollingshead, S.K., Fischetti, V.A., and Scott, J.R. (1986). Complete nucleotide sequence of type 6 M protein of the group A streptococcus: repetitive structure and membrane anchor. **J. Biol. Chem.**, 261: 1677-1686.
- Hollingshead, S.K., Fischetti, V.A., and Scott, J.R. (1987). Size variation in group A streptococcal M protein is generated by homologous recombination between intragenic repeat. **Mol. Gen. Genet.**, 207: 196-203.
- Hollingshead, S.K., Readdy, T.L., Young, D.L., and Bessen, D.E. (1993). Structural heterogeneity of the *emm* gene cluster in group A streptococci. **Mol. Microbiol.**, 8: 707-717.
- Hong, K., Kinoshita, T., Takeda, J. et al., (1990). Inhibition of the alternative C3 convertase and classical C5 convertase of complement by group A streptococcal M protein. **Infect. Immun.**, 58: 2535-2541.
- Horstmann, R.D., Sievertsen, H.J., Knobloch, J., and Fischetti, V.A. (1988). Antiphagocytosis activity of streptococcal M protein: selective binding of complement control protein factor. **Proc. Natl. Acad. Sci. USA.**, 85: 1657-1661.
- Huang, T.T., Malke, H., and Ferretti, J.J. (1989). Heterogenicity of the streptokinase gene in group A streptococci. **Infect. Immun.**, 57: 502-506.
- Huxley, H.E., and Perutz, M.F. (1953). New X-ray evidence on the configuration of polypeptide chain. **Nature (London)**, 167: 1053-1054.

- Imanishi, K., Igarashi, H., and Uchiyama, T. (1990). Activation of murine T cells by streptococcal pyrogenic exotoxin type A. Requirement of MHC class II molecule on accessory cells and identification of $\nu\beta$ - elements in T cells receptor of toxin-reactive T cell. **J. Immunol.**, 145: 3170-3176.
- Innes, M.A., and Gelfand, D.H. (1990). Optimization of PCR. in Innes, M.A., Gelfand, D.H., Sninky, J.J., and White, J.J.(eds), PCR protocols: a guide to methods and applications (pp. 3-12), America: Academic Press.
- Isselbacher, K.J., Braunwald, F., Wilson, J.D., Martin, J.B., Fauci, A.S., and Kasper, D.L. (1994). Harrison's principles of internal medicine. Vol 14 13rd ed, McGraw-Hill, Inc. Health Profession Division.
- Jacks-Weis, I., Kim, Y., and Cleary, P.P. (1982). Restricted deposition of C3 on M+ group A streptococci: correlation with resistance of phagocytosis. **J. Immunol.**, 128: 1897-1902.
- Jamal, F., Pit, S., Johnson, D.R., and Kaplan, E.L. (1995). Characterization of group A streptococci isolated in Kuala Lumpur, Malaysia. **J. Trop. Med. Hyg.**, 98: 343-346.
- Jawetz, F., Melnick, J.L., Adelberg, E.A. (eds.) (1995). The Streptococci. in: Medical Microbiology, 20th ed.(pp. 192), U.S.A.: Appleton & Lange. A Simon Schuster Company.
- Johnson, D.R., and Kaplan, E.L. (1988). Microtechnique for serum opacity factor characterization of group A streptococci adaptable to the use of human sera. **J. Clin. Microbiol.**, 26: 2025-2030.
- Johnson, D.R., and Kaplan, E.L. (1993). A review of the correlation of T-agglutination patterns and M-protein typing and opacity factor production in the identification of group A streptococci. **J. Med. Microbiol.**, 38: 311-315.

- Johnson, D.R., Stevens, D.L., Kaplan, E.L. (1992). Epidemiology analysis of group A streptococcal serotypes associated with severe systemic infection, rheumatic fever, or uncomplicated pharyngitis. **J. Infect.Dis.** 166: 374-382.
- Johnson, I.P., and Schievert, P.M. (1984). Group A streptococcal phage T12 carries the structural gene for pyrogenic exotoxin A. **Mol. Gen.Genet.**, 195: 52-56.
- Jupin, C., Anderson, S., Damias, C., Alouf, J.E., and Parant, M. (1988). Toxic shock syndrome toxin 1 as an inducer of tumor necrosis factors and gamma interferon. **J. Exp. Med.**, 167: 752-761.
- Kaper, J.B., Bradford, H.B., Roberts, N.C., Falkow, S. (1982). Molecular epidemiology of *Vibrio cholerae* in the U.S. gulf coast. **J. Clin. Microbiol.**, 16: 129-34.
- Kaplan, E.L., Johnson, D.J., and Cleary, P.P. (1989). Group A streptococcal serotype isolate from patients and sibling contacts during the resurgence of rheumatic fever in the United States in the mid -1980s. **J. Infect. Dis.**, 159;101-103
- Kaplan, E.L., Johnson, D.R., Nanthapisud, P., Sililertpanrana, S., and Chumdermpadetsuk, S. (1992). A Comparison of group A streptococcal serotypes isolates from the upper respiratory tract in the USA and Thailand: Implications. **Bull. of WHO.**, 70: 433-437.
- Katsukawa, C. (1994). Cloning and nucleotide sequence of type 3 M protein gene (*emm3*) consisting of an N-terminal variable portion and C-terminal conserved C repeat regions: relation to other genes of *Streptococcus pyogenes*. **Kansenshogaku Zasshi.**, 68: 698-705.
- Kaufhold, A., and Ferrieri, P. (1993). The Microbiologic aspects, including diagnosis, of β -hemolytic streptococcal and enterococcal infections. **Infectious Disease Clinics of North America.**, 7: 235-256.

- Kaufhold, A., Podbielski, A., Johnson, D.R., Kaplan, E.L., and Lütticken, R. (1992). M protein gene typing of *Streptococcus pyogenes* by nonradioactively labeled oligonucleotide probes. **J. Clin. Microbiol.**, 30: 2391-2397.
- Kaufhold, A., Podbielski, A., Baumgarten, G., Blokpoel, M., Top., J., and Schouls, L. (1994). Rapid typing of group A streptococci by the use of DNA amplification and nonradioactive allele-specific oligonucleotide probes. **FEMS. Microbiol. Lett.**, 119: 19-26.
- Kehoe, M.A. (1994). Cell-wall associated protein in Gram-positive bacteria. **New. Compr. Biochem.**, 27: 217-261.
- Kotb, M. (1992). Role of superantigens in the pathogenesis of infectious disease and sequelae. **Curr. Opin. Infect. Dis.**, 5: 304-374.
- Kotb, M., Ohnishi, H., Majumdar, G., Hackett, S., Bryant, A., Higgins, G., and Steven, D. (1993). Temporal relationship of cytokine release by peripheral blood mononuclear cells stimulated by the streptococcal superantigen pep M5. **Infect. Immun.**, 61: 1194-1201.
- Kristiansen, B.E., Sorensen, B., Simonsen, T., Spanne, O., Lund, V., and Bjorvatn, B. (1984). Isolates of *Neisseria meningitidis* from different site in the same patients: phenotypic and genotypic study, with special reference to adherence, Palliation and DNA restriction endonuclease pattern. **J. Infect. Dis.**, 150: 389-396.
- Kronvall, G. (1973). A surface component in group A, C and G streptococci with non-immune reactivity for immunoglobulin G. **J. Immunol.**, 111: 1401-1406.
- Land, M.A., and Bisno, A.L. (1983). Acute rheumatic fever: a vanishing disease in suburbia. **JAMA.**, 249: 894-898.
- Lancefield, R.C. (1928). The antigenic complex of *Streptococcus hemolyticus*. I. Demonstration of a type-specific substance in extracts of *Streptococcus hemolyticus*. **J. Exp. Med.**, 47: 91-103.

- Lancefield, R.C. (1940). Specific relationship of cell composition to biological activity of hemolytic streptococci. **Harvey Lect.**, 36: 251.
- Lancefield, R.C. (1962). Current knowledge of type specific M antigens of group A streptococci. **J. Immunol.**, 89: 307-313.
- Lange, K., Ahmed, U., Kleinberger, H. and Treser, G. (1976). A neither to unknown Streptococcal antigen and its probable relation to acute poststreptococcal glomerulonephritis. **Clin. Exp. Immunol.**, 34: 318-325.
- Le Febvre, R.B., Foley, J.W., Thiermann, A.B. (1985). Rapid and simplified protocol for isolation and characterization of leptospiral chromosomal DNA for taxonomy and diagnosis. **J. Clin. Microbiol.**, 22: 606-8
- Lennon, D., and Martin, D., Wong, E., Anizmit, T.(1988). Longitudinal study of post streptococcal diseases in Auckland: rheumatic fever, glomerulonephritis, epidemiology and M typing 1881-1886. **NZ. J. Med.**, 101: 396-398.
- Majeed, H. A., Khuffash, F.A., Yousof, A.M., Farwana, S.S., Chung, T.Y., Rotta, j. (1980). The rheumatogenic and nephitogenic strains of the group A streptococcus; the Kuwait experience. **NZ. J. Med.**, 8:398-401.
- Manjula, B.N., Acharya, A.S., Mische, S.M ., Fairwell, T., Trus, B.L., and Fischetti V.A. (1985). The complete amino acid sequence of streptococcal Pep M5 protein: presence of two structurally distinct domains and implications of biological functions. in Y. Kimura, S. Kotani, and Y. Shiokawa. (ed.), Recent advances in streptococci and streptococcal disease (pp 185-187), England: Reedbooks Ltd., Chertsey.
- Manjula, B.N., and Fischetti, V.A. (1980). Studies on group A streptococcal M proteins: purification of type 5 M-protein and comparison of its amino terminal sequence with two immunologically unrelated M-protein molecules. **J. Immunol.**, 124: 261-267.

- Manjula, B.N., Acharya, A.S., Mische, S.M., Fairwell, T., and Fischetti, V.A. (1984). The complete amino acid sequence of a biologically active 197-residue fragment of M protein from type 5 group A streptococci. **J. Biol. Chem.**, 259: 3686-3693.
- Manjula, B.N., Khandke, K.M., Fairwell, T., Relf, W.A., and Sriprakash, K.S. (1991). Heptad motifs within the distal subdomain of the coiled-coil rod region of M-protein from rheumatic fever and nephritis associated serotypes of group A streptococci are distinct from each other: nucleotides sequence of the M57 gene and relation to the deduce amino acid sequence of other M proteins. **J. Prot. Chem.**, 10: 369-384.
- Markowitz, M. (1985). The decline of rheumatic fever: role of medical intervention. **J. Pediatr.**, 106: 545-550
- Markowitz, M. (1991). Streptococcal disease in developing countries. **Pediatr. Infect. Dis.**, 10: 511-514.
- Markowitz, M. (1994). Changing epidemiology of group A streptococcal infections. **Pediatr. Infect. Dis. J.**, 13: 557-560.
- Martin, D.R., and Single, L.A. (1992). Molecular epidemiology of group A streptococcus M protein 1 infection. **J. Inf. Dis.**, 167: 1112-1117.
- Martin, D.R., Hoby, E.A. (1990). Streptococcal group A epidemic in Norway 1987-1988. Scand. **J. Infect. Dis.**, 22: 421-429.
- Maslow, J.N., Slutsky, A.M., and Arbeit, R.D. (1993). The application of pulsed field gel electrophoresis to molecular epidemiology. in Persing D.H. Tenover, F.C. (pp563-572), Washington. DC: American Society for Microbiology..
- Massel, B.F., Chute, C.G., Walker, A.M., Kurland, G.S. (1988). Penicillin and the marked decrease in morbidity and mortality from rheumatic fever in the United States. **N. Engl. J. Med.**, 31: 280-286.
- Matsuno, T., and Slade, H.D., (1970). Composition and properties of group A streptococcal teichoic acid. **J. Bacteriol.**, 102: 747-752.

- Maxted, W.R., Widdowson, J.P. (1972). The protein antigens of group A streptococci. in Wannamaker L.W. Matsen J.M. (eds.), Streptococci and streptococcal disease (pp. 251-266), New York: Academic Press.
- McCarty, M. (1990). Streptococcus. in Davis, B.D., Dulbecco, B., Eisen, H.N., and Ginsberg, H.S. (eds), Microbiology. 4th edition (pp. 528-529), Singapore: Harper and Row, Publisher.
- McLachlan, A.D., and Stewart, M (1975). Tropomyosin coiled-coil interaction: evidence for an unstaggered structure. **J. Mol. Biol.**, 98: 393-398.
- Miller, L., Grey, L., Beachey, E.H., and Kehoe, M.A. (1988). Antigenic variation among group A streptococcal M proteins: nucleotide sequence of serotype 5 M protein gene and its relationship with gene encoding type 6 and type 24 M protein. **J. Biol. Chem.**, 263: 5668-5673.
- Mische, S.M., Manjula, B.N., and Fischetti, V.A. (1987). Relation of streptococcal M protein with human and rabbit tropomyosin: the complete acid sequence of human cardiac alpha tropomyosin, a highly conserved contractile protein. **Biochem. Biophys. Res. Comm.**, 142: 813-818.
- Moody, M.D., Padula, J., Lizana, D., and Hall, C.T. (1965). Epidemiologic characterization of group A streptococci by T-agglutination and M-precipitation tests in the public health laboratory. **HLth. Lab. Sci.**, 2: 149-162.
- Mouw, A.R., Beachey, E.H., and Burdett, V. (1988). Molecular evolution of streptococcal M protein: cloning and nucleotide sequence of type 24 M protein gene and relation to other genes of *Streptococcus pyogenes*. **J. Bacteriol.**, 170: 676-684.

- Murray, P.R., Kobayashi, G.S., Pfaller, M.A., Rosenthal, K.S. (eds.) (1994). *Streptococcus and Related Gram-positive bacteria*. in Medical Microbiology 2 nd ed. International Student Edition Published, Wolfe inprint.
- Murray, P.R., Wold, D.R., Marsha, M.H., and Washington II, J.A. (1976). Bacitracin differentiation of presumptive identification of group A beta-hemolytic streptococci. **J. Pediatr.**, 89: 576
- Musser, J.M., Grey, B.M., Schlivert, P.M., and Pichichero, M.E. (1992). *Streptococcus pyogenes* pharyngitis: characterization of strains by multilocus enzyme genotype, M and T protein serotype and pyogenic exotoxin gene probing. **J. Clin. Microbiol.**, 30: 600-603.
- Musser, J.M., Hauser, A.R., Kim, M.H., Schlievert, P. M., Nelson, K., and Selander, R.K. (1991). *Streptococcus pyogenes* causing toxin-shock-like syndrome and other invasive disease: Clonal diversity and pyrogenic exotoxin expression. **Proc. Natl. Acad. Sci. USA.**, 88: 2668-2672.
- Musser, J.M., Kapur, V., Szeto, J., Pan, X., Swanson, D.S., and Martin, D.R. (1995). Genetic diversity and relationships among *Streptococcus pyogenes*. Strains expressing serotype M1 protein: Recent intercontinental spread of a subclone causing episodes of invasive disease. **Infect. Immun.**, 63: 994-1003.
- Mylvaganam, H., Bjorvatn, B., Hofstad, T., Hjetland, R., Arne Hoiby, E., and Holms, S.E. (1994). Small-fragment restriction endonuclease analysis in epidemiological mapping of group A streptococci. **J. Med. Microbiol.**, 40: 256-260.
- Norrby-Teglund, A., Newton, A.D., Kotb, M., Holm, S.E., and Norgren, M. (1994). Superantigenic properties of group A streptococcal exotoxin, SPEF (MF). **Infect. Immun.**, 62: 5227-5233.

- O'Toole, P., Stenberg, L., Rissler, M., and Lindahl, G. (1992). Two major classes in the M protein family in group A streptococci. **Proc. Natl. Acad. Sci. USA.**, 89:8661-8665.
- Ofek, I., Beachey, E.H., Jefferson, W., and Campbell, G.L., (1975). Cell membrane binding properties of group A streptococcal lipoteichoic acid. **J. Exp. Med.**, 141: 990-1003.
- Okada, N., Geist, R.T., and Caperon, M.G. (1993). Positive transcriptional control of *mry* regulates virulence in the group A streptococcus. **Mol. Microbiol.**, 7: 893-903.
- Parry, D.A.D., and Fraser, R.D.B. (1985). Intermediate filament structure. I. analysis of 1F protein sequence data. **Int. J. Biol. Macromol.**, 7: 203-213.
- Pappenheimer, A.M.Jr, Murphy, J.R. (1983). Studies on the molecular epidemiology of diphtheria. **Lancet**, 2: 923-6.
- Parsonnet, J., Hickman, R.K., Eardley, D.P., and Pier, G. (1985). Induction of human interleukin 1 by toxic shock syndrome toxin 1. **Infect. Immun.** 55: 1070-1076.
- Penney, T.J., Martin, D.R., Williams, L.C., DeMalmanche, S.A., and Bergquist, P.L. (1995). A single *emm* gene-specific oligonucleotide probe does not recognise all members of the *Streptococcus pyogenes* M type 1. **FEMS. Microbiol. Lett.**, 130: 145-150.
- Peterson, P.K., Schmeling, D., Cleary, P.P., Wilkinson, B.J., Kim, Y., and Quie, P.G.(1979). Inhibition of alternative complement pathway opsonization by group A streptococci M protein. **J. Infect. Dis.**, 139: 575-585.
- Philips, G.N., Flicker, P.F., Cohen, C., Manjula, B.N., and Fischetti, V.A. (1981). Streptococcal M protein: alpha-helical coiled-coil structure and arrangement on the cell surface. **Proc. Natl. Acad. Sci. USA.**, 78: 4689-4693.

- Podbielski, A. (1993). Tree different types of organization of the *vir*- regulon in group A streptococci. **Mol. Gen. Genet.**, 237: 287-300.
- Podbielski, A., Kaufhold, A., and Cleary, P.P.(1993). PCR-method amplification of group A streptococcal genes encoding immunoglobulin-binding protein., **Immuno. Meth.** 2: 55-64.
- Podbielski, A., Melzer, B., and Lüttichen, R.(1991). Application of the polymerase chain reaction to study the M protein (-like) gene family in β -hemolytic streptococci. **Med. Microbiol. Immunol.**, 180: 213-227.
- Podbielski, A., Weber-Heynemann, J., and Cleary, P.P.(1994). Immunoglobulin-binding FcrA and Enn proteins and M proteins of group A streptococci evolved independently from a common ancestral protein. **Med. Microbiol. Immunol.**, 183: 33-42.
- Prakash, K., and Dutta, S. (1991). Application of serum opacity factor in subtyping of group A streptococci & identification of new M-types. **Indian. J. Med. Res. [A].**, July: 222-224.
- Pruksakorn, S. Unpublished data
- Pruksakorn, S., Phornphutkul, C., Boonchoo, S., Supasert, S., Krairojananan, M., Boonthum, A., Murai, T., Inazumi, Y., Sriburee, P., and Tharavichitkul, P. (1990). Prevalence of group A streptococci from school children and patients in Chiangmai, Thailand. **Chiangmai. Med. Bull.**, 29: 15-26.
- Quinn, R.W. (1982). Streptococcal infections. in Evany, A.S. and Feldman, H.A. (ed.), Bacterial infections of humans Epidemiology and Control, New York and London: Plenum Medical Book Company, .
- Quinn, R.W., Vander-Zwagg, R., and Lowry, A.N. (1985). Acquisition of group A streptococcal M protein antibodies. **Pediatr. Infect. Dis.**, 4: 374-378.

- Rabkin, C., Jarvis, W., Anderson, R., Govan, J., Klinger, J., LiPuma, J., Martone, W., Monteil, H., Richard, C., Shigeta, S., Sossa, A., Stull, J., Swenson, J., and Woods, D. (1989). *Pseudomonas cepacia* Typing systems Collaborative study to assess their potential in epidemiologic investigations. **Rev. Infect. Dis.** 11: 600-607.
- Reader, R., and Boyle, MD.P. (1993a). Association between expression of immunoglobulin-G binding proteins by group A streptococci and virulence in mouse skin infection model. **Infect. Immun.**, 61; 1378-1384.
- Reader, R., and Boyle, MD.P. (1993b). Association of type II Immunoglobulins -G binding protein expression and survival of group A streptococci in human blood. **Infect. Immun.**, 61: 3696-3702.
- Reed, W.P., Matzler, C., and Albright, E. (1994). Streptococcal adherence to langerhans cells: A possible step in the pathogenesis of streptococcal pharyngitis. **Clin. Immun. Immunopathol.**, 70: 28-31.
- Relf, W.A., and Sriprakash, K.S. (1990). Limited repertoire of the C-terminal region of the M-protein in *Streptococcus pyogenes*. **FEMS Microbiology Lett.**, 71: 345-350.
- Relf, W.A., Martin, D.R., and Sriprakash, S. (1992). Identification of sequence type among the M-nontypable group A streptococci. **J. Clin Immunol**, 30: 3190-3194.
- Relf, W.A., Martin, D.R., and Sriprakash, K.S. (1994). Antigenic diversity within a family of M proteins from group A streptococci: Evidence for the role of frameshift and compensatory mutation. **Gene.**, 144: 25-30.
- Rotta, J. (1978). Group and type (group A and B) identification of haemolytic streptococci. in Bergan T, Norris JR (eds), *Methods in Microbiology*, Vol 12 (pp. 177-198), London: Academic Press.
- Rotta, J., and Facklam, R.R. (1980). *Manual of microbiological diagnostic method for streptococcal infection and their sequelae*. World Health Organization.

- Saiki, R.K.(1989). The design and optimization of the PCR. in Erlic, H.A.(ed), PCR technology principle and applications for DNA amplification (pp7-16), America: Stockton press.
- Salvadori, L.G., Blake, M.S., McCarty, M., Tai, J.Y., and Zabriskie, J.B. (1995). Group A streptococcus-liposome ELISA antibody titer to group A polysaccharide and opsonophagocytic capabilities of the antibodies. **J. Infect.Dis.**, 171:593-560.
- Saurnier, P., Bournenx, C., Prevost, G., and Andremont, A. (1993). RAPD analysis assay is less discriminant than pulse-field gel electrophoresis for typing strains of methicillin-resistant *Staphylococcus aureus*. **J.Clin. Microbiol.** 31: 982-985.
- Schalen, C., Truedsson, L., Christensen, K.K., and Christensen, P. (1985). Blocking of antibody complement-dependent effector functions by streptococcal IgG Fc-receptor and staphylococcal protein A. **Acta. Path. Microbiol. Immunol.Scand.**, 93: 395-400.
- Scherer, S., Stevens, D.A. (1987). Application of DNA typing methods to epidemiology and taxonomy of *Candida species*. **J. Clin. Microbiol.**, 25: 675-679.
- Schwartz, B., Facklam, R., and Breiman, R.F. (1990). Changing epidemiology of group A streptococcal infection in the USA. **Lancet.**, 336: 1167-1171.
- Scott, J.R. (1990). The M protein of group A streptococcus: evolution and regulation. in Iglewski, B. (Ed.), *The Bacteria*, Volumn XI.(pp 117-203), New York: Academic Press.
- Selander, R.W., Caugant, D.A., Ochman, H., and Musser, J.E., Gilmour, M.N., and Wittam, T.S. (1986). Methods of multilocus enzyme electrophoresis for bacterial population genetics and systematics. **Appl. Environ. Microbiol.**, 51: 873-884.

- Seppälä, H., He, Q., Osterblad, M., and Huovinen, P. (1994a). Typing of group A streptococci by random amplified polymorphic DNA analysis. **J. Clin. Microbiol.**, 32: 1945-1948.
- Seppälä, H., Vuopio-Varkila, J., Osterblad, M., Jahkola, H., Rummukainen, M., Holm, S.E., and Huovinen, P. (1994b). Evaluation of method for epidemiologic typing of group A streptococci. **J. Inf. Dis.**, 169: 519-525.
- Simpson, W.J., LePenta, D., Chen, C., and Cleary, P. (1990). Coregulation of type 12 M protein and streptococcal C5a peptidase genes in group A streptococci: evidence for a virulence regulon controlled by the vir-R locus. **J. Bacteriol.**, 172: 696-700.
- Skjold, S.A., Quie, P.G., Fries, L.A., Barnham, M., and Cleary, P.P. (1987). DNA fingerprinting of *Streptococcus zooepidemicus* (Lancefield Group C) as an aid to epidemiological study. **J. Infect. Dis.**, 155: 1145-1150.
- Stevens, D.L., Tanner, M.H., Winship, J. et al. (1989). Severe group A streptococcal infection associated with a toxic shock-like syndrome and scarlet fever toxin A. **N. Engl. J. Med.**, 312: 1-7.
- Stokes, E.J. (ed.) (1975). Streptococci. In: Clinical Bacteriology. pp 113. An Enward Arnold Publication.
- Stull, T.L., LiPuma, J.J., and Edlind, T.D. (1988). A broad-spectrum probe for molecular epidemiology of bacteria: ribosomal RNA. **J. Infect. Dis.**, 157: 280-286.
- Swanson, J., Hsu, K.C., and Gotschlich, E.C. (1969). Electron microscopic studies on streptococci, I. M antigen. **J. Exp. Med.**, 130: 1063-1091.
- Swift, H.F, Wilson, A.T., Lancefield, R.C. (1943). Typing group A hemolytic streptococci by M precipitin reactions in capillary pipettes. **J. Exp. Med.**, 78: 127-133.
- Taylor, G.R. (1991). Polymerase chain reaction: basic principles and automation. in McPherson, M.J., Quirke, P., and Taylor, G.R. (eds), PCR a practical approach (pp1-14), New York: Oxford University Press.

- Tenover, F.C., Arbeit, R., Archer, G., Biddle, J., Byrne, S., Goering, R., Hancock, G., Hebert, G.A., Hill, B., Hollis, R., Jarvis, W.R., Kreiswirth, B., Eisner, W., Maslow, J., McDougal, L.K., Miller, J.M., Mulligan, M., and Pfaller, M.A. (1994). Comparison of traditional and molecular methods of typing isolates of *Staphylococcus aureus*. **J. Clin. Microbiol.**, 32 : 407-415.
- Tomai, M.A., Schleivert, P.M., and Kotb, M. (1992). Distinct T cell receptor $\nu\beta$ -gene usage by human T lymphocytes simulated with streptococcal pyogenic exotoxins and pep M 5 protein. **Infect. Immun.**, 60: 701-705.
- Tomai, M., Kotb, M., Manjumdar, G., and Beachey, E.H. (1990). Superantigenicity of streptococcal M protein. **J. Exp. Med.**, 172: 359-362.
- Tomai, M., Aelion, J.A., Dockter, M.E., Manjumdar, G., Spirella, D.G., and Kotb, M. (1991). T cell receptor V gene usage by human T cells stimulated with the superantigen streptococcal M protein. **J. Exp. Med.**, 174: 285-288.
- Top, F.H., and Wannamaker, L.W. (1968). The serum opacity factor of *Streptococcus pyogenes*: frequency of production of streptococcal lipoproteinase by strains of different serological types and the relationship to M protein production. **J. Hyg.**, 66: 49-58.
- Tran, P.O.T., Johnson, D.R. and Kaplan, E.L. (1993). The presence of M protein in nontypable group A streptococcal upper respiratory tract isolates from Southeast Asia. **J. Infect. Dis.**, 169: 658-661.
- Van de Rijn, I., and Fischetti, V.A. (1981). Immunochemical analysis of intact M protein secreted from cell wall-less streptococci. **Infect. Immun.**, 32: 86-91.
- Vashishtha, A., and Fischetti, V.A. (1993). Surface-exposed conserved region of the streptococcal M protein induces antibodies cross-reactive with denatured forms of myosin. **J. Immunol.**, 150: 4693-4701.

- Veasy, L.G., Wiedmeier, S.E., Orsmond, G.S. et al. (1987). Resurgence of acute rheumatic fever in the intermountain area of the United States. **N. Engl. J. Med.**, 316: 421-427.
- Villarreal, H., Fischetti, V.A., Van de-Rijn, I., Zabriskie, J.B. (1979). The occurrence of a protein in the extracellular products of streptococci isolated from patients with acute glomerulonephritis. **J. Exp. Med.**, 149: 459-472.
- Welsh, J., and McClelland, M. (1990). Fingerprinting genomes using PCR with arbitrary primers. **Nucleic. Acids. Res.**, 18: 7213-7218.
- Wessels, M.R. (1994). Streptococcal infection. in Isselbacher, J.H., Braunwald, E., Wilson, J.D., Martin, J.B., Fanci, A.S., and Kasper, D.L. (eds), Harrison's principles of internal medicine Vol 1, 13th ed, United States of America: McGraw-Hill Inc.
- Wexler, D.E., and Cleary, P.P. (1985). Purification and characteristics of the streptococcal chemotactic factor inactivator. **Infect. Immun.**, 50: 775-764.
- Whatmore, A.M., Kapur, V., Musser, J.M., and Kehoe, M.A. (1995). Molecular genetic analysis of the *emm* subdivision of group A streptococcal *emm*-like gene: horizontal gene transfer and restricted variation among *emm* gene. **Mol. Microbiol.**, 15: 1039-1048.
- Whatmore, A.M., Kapur, V., Sullivan, D.J., Musser, J.M., and Kehoe, M.A. (1994a). Noncongruent relationships between variation in *emm* gene sequences and the population genetic structure of group A streptococci. **Molecular Microbiology.**, 14: 619-631.
- Whatmore, A.M., and Kehoe, M.A. (1994b). Horizontal gene transfer in the evolution of group A streptococcal *emm*-like gene; mosaics and variation in *vir* regulon. **Molecular Microbiology.**, 11: 363-374.

- WHO Cardiovascular Disease Unit and principal investigators.(1992). WHO program for the prevention of rheumatic fever/rheumatic heart disease in 16 developing countries: report from phase I (1986-1990). **Bull. WHO.**, 70: 213-218.
- WHO. Rheumatic fever and rheumatic heart disease.(1988). Report of a WHO study group. WHO technical report series. 764, Geneva, p. 5-58.
- Widdowson, J.P., Maxted, W.R., and Grant, D.L. (1970). The production of opacity in serum by group A streptococci and its relationship with the presence of M antigen. **J. Gen. Microbiol.**, 61:343-353.
- Widdowson, J.P., Maxted, W.R., Grant, O.L., and Pinney, A.M. (1971). The relationship between M antigen and opacity factor in group A streptococci. **J. Gen. Microbiol.**, 128: 959-968.
- Willett, H.P. (1992). Streptococcus. in Joklik, W.K., Willett; H.P., Amos, O.B., and Wilfet, C.M.(eds), Zinsser Microbiology. 20th edition,(pp 420), California: Publishing Division of Prentice Hall.
- Williams, J.G.K., Kubelik, A.R., Livak, K.J., Rafalski, J.A., and Tingey, S.V. (1990). DNA polymorphisms amplified by arbitrary primers are useful as genetic marker. **Nucleic. Acids. Res.**, 18: 6531-6535.
- Williams, REO. (1958). Laboratory diagnosis of streptococcal infection. **Bull. Wld. Hlth. Org.**, 19: 153-176.