

TABLE OF CONTENTS

	PAGE
ACKNOWLEDGEMENT	iii
ABSTRACT IN THAI	iv
ABSTRACT IN ENGLISH	vi
LIST OF TABLES	x
LIST OF ILLUSTRATION	xii
CHAPTER 1 INTRODUCTION	1
1.1 General overview	1
1.2 Review of regional geology	2
1.3 Structure framework	5
1.4 Basin formation	8
1.5 Exploration history in the Gulf of Thailand	8
1.6 Fluorescence microscopy history	11
1.7 Purpose of Study	12
CHAPTER 2 REGIONAL GEOLOGY	13
2.1 Location and extent of the study Area	13
2.2 Type of sample and data source	15
2.3 Development of the Pattani Basin	15
2.4 Stratigraphy and sedimentation in the Pattani Basin	18
2.5 Organic characteristic in the Pattani Basin	24
CHAPTER 3 STRATIGRAPHY	27
3.1 Introduction	27
3.2 Lithology of each well	27
CHAPTER 4 COAL PETROGRAPHY	55
4.1 Definition of macerals	55
4.2 Fluorescence microscopy	55

4.3 Coal macerals	56
4.3.1 The huminite/vitrinite group	58
4.3.2 The liptinite group	61
4.3.3 The inertinite group	66
4.4 Fluorescence of maceral	69
4.5 Procedure	70
4.5.1 Sample preparation	70
4.5.2 Identification procedure	71
CHAPTER 5 RESULTS AND INTERPRETATION	74
5.1 Maceral distribution	74
5.1.1 Maceral distribution in the east graben	74
5.1.2 Maceral distribution in the west graben	96
5.2 Interpretation	120
5.2.1 Geographic distribution	120
5.2.1.1 Vertical distribution	120
5.2.1.2 Lateral distribution	120
5.2.2 Lithology and maceral distribution	121
5.2.3 Organic maturity	133
5.3 Paleoenvironment	133
CHAPTER 6 CONCLUSIONS	156
REFERENCES	172
APPENDIX A INFORMATION AND LITHOLOGY OF EACH WELL	177
APPENDIX B MACERALS CONTAIN IN EACH WELL	251
CURRICULUM VITAE	299

LIST OF TABLES

TABLES	PAGE
1.1 Pattani Basin, Gulf of Thailand production (as of January to September, 1998)	4
2.1 List of well's name, position grid, total cutting depth and wireline log depth	17
2.2 Stratigraphy and depositional environment of Tertiary synrift strata, Pattani Basin	22
2.3 Stratigraphy and depositional environment of Tertiary postrift strata, Pattani Basin	23
4.1 Classification of coal macerals	57
4.2 Classification of low-range coal macerals	58
4.3 Fluorescence color and intensity for the maceral groups at difference coal rank using blue - light excitation	70
5.1 Relationship between lithology and maceral distribution in A well	125
5.2 Relationship between lithology and maceral distribution in B well	126
5.3 Relationship between lithology and maceral distribution in C well	127
5.4 Relationship between lithology and maceral distribution in D well	127
5.5 Relationship between lithology and maceral distribution in E well	128
5.6 Relationship between lithology and maceral distribution in F well	129

TABLES	PAGE
5.7 Relationship between lithology and maceral distribution in G well	129
5.8 Relationship between lithology and maceral distribution in H well	130
5.9 Relationship between lithology and maceral distribution in I well	130
5.10 Relationship between lithology and maceral distribution in J well	131
6.1 Conclusion the relationship between lithology and maceral distribution in the Pattani Basin	167

LIST OF ILLUSTRATIONS

FIGURE	PAGE
1.1 Map showing petroleum fields in thailand	3
1.2 Tectonic map of SE Asia and South China showing the main fault patterns and the relative movement of crustal blocks in response to the collision of India with Asia	6
1.3 Diagrammatic representation of the development of pull-apart basin in the Gulf of Thailand by the major strike-slip fault systems	9
1.4 Map showing petroleum operated blocks in the Gulf of Thailand	10
2.1 Isopach map of Tertiary basins in the Gulf of Thailand showing location of the Pattani and other major basins	14
2.2 Study area and location of wells used in this study	16
2.3 Gulf of Thailand stratigraphic sequence	19
2.4 Stratigraphic summary of the major depositional sequences within the Pattani Basin	25
3.1 Symbol description of lithology used in Figure 3.2 and 3.3	28
3.2 Lithological correlation between well in the east graben of the Pattani Basin	29
3.3 Lithological correlation between well in the west graben of the Pattani Basin	30
4.1 MPM200 microscope photometer for fluorometry with axioplan universal microscope as base unit	73
5.1 Symbol description about macerals contain used in Figure 5.2, 5.3, 5.4, 5.5, 5.6, 5.7 and 6.1	75

FIGURE	PAGE
5.2 Macerals distribution in each well in the east graben of the Pattani Basin	76
5.3 Macerals distribution in each well in the west graben of the Pattani Basin, north-south direction	77
5.4 Macerals distribution in each well in the west graben of the Pattani Basin, east-west direction	77
5.5 Relationship between macerals distribution and lithology in each sequence in east graben of the Pattani Basin	122
5.6 Relationship between macerals distribution and lithology in each sequence in west graben of the Pattani Basin, in north-south direction	123
5.7 Relationship between macerals distribution and lithology in each sequence in east graben of the Pattani Basin, in east-west direction	124
5.8 Depositional environments of lower sequence 1	134
5.9 Idealized reconstruction of alluvial-fan peats within an intermontane basin	134
5.10 Hypothetical intermontane fluvial-deltaic system analogous to the Pattani Basin during Miocene deposition	136
5.11 Pattern of peat accumulation in backswamps adjacent to meandering fluvial channels	136
5.12 Exsudatinite impregnated in clay grain displayed pale green color and zoning at -3281.1 ft to -3393.5 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	138
5.13 Exsudatinite impregnated in clay grain displayed pale green color and zoning at -3393.5 ft to -3504.2 ft under (left) under	

FIGURE	PAGE
oil-fluorescence light (right) under oil-white light (10x20).	138
5.14 Liptodetrinite as groundmass, resinite body (bright yellow), sclerotinite has many cells infill by resinite (exsudatinite) at -3723.2 ft to -3831.6 ft (left) under oil-fluorescence light (right) under oil-white light (10x50).	139
5.15 Resinite mass at -4783.2 ft to -4889.2 ft (left) under oil-fluorescence light (right) under oil-white light (10x50).	139
5.16 Alginite A, sporinite, cutinite and resinite at -3938.1 ft to -4044.3 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	140
5.17 Sapropelic coal has suberinite, liptodetrinite and sporinite at -4254.4 ft to -4359.9 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	140
5.18 Exsudatinite filled in ulminite at -4571.2 ft to -4677.2 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	141
5.19 Suberinite band, liptodetrinite as groundmass and sporinite at -4995.3 ft to -5101.6 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	141
5.20 liptodetrinite as groundmass, sporinite and cutinite at -2910.8 ft to -2992 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	142
5.21 Alginite A at -3572 ft to -3656.3 ft (left) under oil-fluorescence light (right) under oil-white light (10x50).	142
5.22 oil shale has lamaginite rich, exsudatinite impregnated in clay grain at -3820.5 ft to -3902.5 ft (left) under oil-fluorescence	

FIGURE	PAGE
light (right) under oil-white light (10x20).	143
5.23 Lamaginite or alginite B at -3902.5 ft to -3985.5 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	143
5.24 Exsudatinitite impregnated in clay grain displayed pale green color and zoning at -6809.7 ft to -6913.6 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	144
5.25 Sapropelic coal has high liptinite content; suberinite type show degrading state, attrinite and gelinite at -7965.5 ft to -8053.9 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	144
5.26 Sapropelic coal has high liptinite content; suberinite type show degrading state, attrinite and gelinite at -8053.9 ft to -8167.9 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	145
5.27 Cutinite, resinite and liptodetrinite as a groundmass at measure depth 3150 ft to 3720 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	145
5.28 Resinite mass fill in cell cavity of sclerotinite and suberinite at measure depth 3300 ft to 3420 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	146
5.29 Exsudatinitite filled in ulminite at measure depth 3600 ft to 3720 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	146
5.30 Liptodetrinite, bituminite (brown) at measure depth 3750 ft to 3870 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	147

FIGURE	PAGE
5.31 Thick layer of suberinite, diffuse in texture, significant degrading at measure depth 4200 ft to 4320 ft (left) under oil-fluorescence light (right) under oil-white light (10x50).	147
5.32 Liptodetrinite as a groundmass, bituminite (brown) indicated degrading at -4218.9 ft to -4292.8 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	148
5.33 Oil shale has lamaginite or alginite B rich, Alginite A (bright yellow) at -3234.7 ft to -3384.7 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	148
5.34 suberinite band, sporinite, and liptodetrinite at -3984.5 ft to -4134.3 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	149
5.35 Alginite A (?) as an unidentified type or clay flocculate impregnated by exsudatinite show radial ring layer and oxidation rim (brown) at -5933.3 ft to -6047.3 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	149
5.36 Sapropelic coal has high liptinite content show degrading state, gelinite has exsudatinite fill in crack, and quartz grains at -6047.3 ft to -6163.7 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	150
5.37 Sapropelic coal has huminite and exsudatinite impregnated in clay grain displayed yellow color at -6771.1 ft to -6897.6 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	150
5.38 Sapropelic coal of high liptinite content shows degrading, in gelinite coal, exsudatinite impregnate, quartz grains at -7024.2 ft	

FIGURE	PAGE
to -7154.6 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	151
5.39 Sapropelic coal of high liptinite content show degrading liptodetrinite and resinite body at -3190.0 ft to -3340.0 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	151
5.40 Thick band of lamaginite and liptodetrinite as groundmass, resinite (bright yellow), bituminite (brown) at -3340.0 ft to -3490.0 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	152
5.41 Exsudatinite impregnated in clay grains displayed pale green color and show zoning, surround by liptodetrinite at -3939.7 ft to -4088.2 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	152
5.42 Alginite A (?) as an unidentified type at -4376.8 ft to -4515.1 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	153
5.43 Resinite trap at -3670.0 ft to -3790.9 ft (left) under oil-fluorescence light (right) under oil-white light (10x50).	153
5.44 Suberinite band at -4031.2 ft to -4150.1 ft (left) under oil-fluorescence light (right) under oil-white light (10x50).	154
5.45 Alginite B degrading as groundmass, sporinite at -3670 ft to -3790.9 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	154
5.46 Alginite A mass at -4268.9 ft to -4389.7 ft (left) under oil-fluorescence light (right) under oil-white light (10x50).	155

FIGURE	PAGE
5.47 Mass of sporinite in cannel coal at -3183.7 ft to -3305.6 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	155
5.48 Alginite A (?) at -6040.1 ft to -6169.5 ft (left) under oil-fluorescence light (right) under oil-white light (10x50).	156
5.49 Lamaginite or alginite B, resinite body (bright yellow) at -2984.5 ft to -3080.1 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	156
5.50 Sapropelic coal of high liptinite content shows degrading, exsudatinite filling in cell cavity of sclerotinite at -3516.7 ft to -3634.6 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	157
5.51 Alginite A(?) and clay flocculate impregnated by exsudatinite show ring layer at -3753.2 ft to -3871.4 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	157
5.52 Sapropelic coal of high liptinite content shows degrading, exsudatinite impregnated and interbedded, gelinite, and quartz grain, at -5195.9 ft to -5306.6 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	158
5.53 Exsudatinite impregnated in quartz grains displayed yellow to deepest yellow at -8064.3 ft to -8199.1 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	158
5.54 Alginite A (?) at -9020.7 ft to -9159.0 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	159
5.55 Oil shale has lamaginite rich at -3437.4 ft to -3542.1 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	159

FIGURE	PAGE
5.56 Mass of sporinite at -3642.8 ft to -3738.0 ft (left) under oil-fluorescence light (right) under oil-white light (10x50).	160
5.57 Exsudatinite impregnated in carbonate crystals displayed yellow color at -5312.9 ft to -5418.7 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	160
5.58 Liptinite maceral; exsudatinite and liptodetrinite type at -3642.8 ft to -3738.0 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	161
5.59 Textinite with exsudatinite filling in cell cavity at -9243.9 ft to -9378.7 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	161
5.60 Resinite trapped between cutinite, exsudatinite filled in cracks of gelinite at -2572.8 ft to -2706.0 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	162
5.61 Exsudatinite filled in ulminite at -3512.7 ft to -3647.4 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	162
5.62 Suberinite show clear structure at -3512.7 ft to -3647.4 ft (left) under oil-fluorescence light (right) under oil-white light (10x20).	163
6.1 Relationship between macerals distribution and lithology in each sequence in the Pattani Basin, east-west direction.	166