

**REFERENCES**

- Altman, D.F. (1998). Drugs used in Gastrointestinal Disease. In B.G. Katzung (ed.), *Basic & Clinical Pharmacolpgy* (7<sup>th</sup> ed., pp. 1017-1029). USA, Prentice Hall International.
- Al-Yahya, M.A., Rafatullah, S., Mossa, J.S., Ageel, A.M., Parmar, N.S. and Tariq, M. (1989). Gastroprotective activity of ginger *Zingiber officinale* Rosc., in albino rats. *American Journal Clinical Medicine*, 17(1-2), 51-56.
- Al-Yahya, M.A., Rafatullah, s., Mossa, S.S., Ageel, A.M., Al-Said, M.S. and Mariq, M. (1990). Gastric antisecretory, antiulcer and cytoprotective properties of ethanol extract of *Alpinia galanga* Willd in rats. *Phytotherapy Research*, 4 (3), 112-114.
- Andrews, C.W. and Goldman, H. (1998). Clinical & Physical Disorders. In S.C. Ming and H. Goldman (eds.), *Pathology of the Gastrointestinal Tract* (2<sup>nd</sup> ed., pp. 195-224). USA, Williams&Wilkins.
- Barrowman, J.A. and Pfeiffer, C.J. (1992). Carbenoxolone: a critical analysis of its clinical value in peptic ulcer. In C.J. Pfeiffer (ed.), *Drugs and Peptic Ulcer* (pp. 123-132). CRC Press, Boca Raton.

- Best, R., Lewis, D.A. and Nasser, N. (1984). The anti ulcerogenic activity of unripe plantain banana (*Musa spp.*). *British Journal of Pharmacology*, 82, 107-116.
- Brito, A.R., Rodroiguez, J.A., Hiruma-Lima, C.A., Hann, M. and Nunes, D.S. (1998). Antiulcerogenic activity of trans-dehydrocroton from *Croton cajucara*. *Planta Medica*, 64(2), 126-129.
- Brodie, D.A. and Hanson, H.M. (1960). A study of the factors involved in the production of gastric ulcers by the restraint technique. *Gastroenterology*, 38, 353-360.
- Brown, H.M., Christie, A.B. and Colin Jone, E., et al. (1959). Glycyrrhetic acid hydrogen succinate (disodium) salt, a new anti-inflammatory compound. *Lancet*, 2, 492-496.
- Brunton, L.L. (1996). Agents for control of gastric acidity and treatment of peptic ulcer. In J.G. Hardman, A.G. Gillman and L.E. Limbird (eds.), *Goodman and Gilman's The Pharmacological Basis of Therapeutic* (9<sup>th</sup> ed., pp. 901-915). New York, Macmillan Publishing.
- Bullock, J. and Boyle, J. (1995). Stomach. In J. Bullock, J. Boyle and M. B. Wong (eds). *Physiology* (3th ed., pp. 422-431). Philadelphia, Williams&Wilkins

- Chang, I.S. (1985). Protective effect of garlic and ethanol-induced gastric mucosal injury in rats. *Journal of Catholic Medical College*, 38(4), 1211-1221.
- Chatterjee, T.K., Chakraborty, A., Plathak, M. and Sengupta, G.C. (1992). Effect of plant extract *Centella asiatica* Linn. on cold restraint stress ulcer in rats. *Indian Journal Experimental and Biological*, 30(10), 889-891.
- Cho, K.H., Chang, T.J., Lee, T.H. and Yoon, C.M. (1981). Clinical experiences of madecasol (*Centella asiatica*) in the treatment of peptic ulcer. *Korean Journal of Gastroenterology*, 13(1), 49-56.
- Corne, S.J., Morrisey, S.M. and Woods, R.J. (1974). A methanolic for the quantitative estimation of gastric barrier mucus. *Journal of Physiology*, 242, 116-117.
- Dai, J.Z., Zhang, J.W., Li, K., Hu, S.L. and Lu, Y. P. (1992). Action against peptic ulcer of gangtosuan. *Zhongguo yiyo Gongye Zazhi*, 23(1), 5.9-511.
- Dau, U.V., Ham, N.N. et al. (1998). The effects of traditional drug, turmeric (*Curcuma longa*), and placebo on the healing of duodenal ulder. *Phytomedicine*, 5, 29-34.
- Friedman, L.S. and Peterson, W.L. (1998). Peptic Ulcer and related disorder. In A.S Fanci, E. Braunwald, K.J. Isselbacher, J.D. Wilson. J. B. Martin, D.L. Kasper, S.L.

- Hanser and D.L. Longo (eds.), *Disorder of the Gastrointestinal System on Harrison's Principles of Internal Medicine* (14<sup>th</sup> ed., pp. 1597-1610). USA, Mcgraw Hill Companies.
- Ganong, W.F. (1997). The adrenal medulla & adrenal cortex. In *Review of Medical Physiology* (18<sup>th</sup> ed. pp. 334-388), Printice-Hall International Inc.
- Gole, R.K., Gupta, S., Shankar, R. and Sanyal, A.K. (1986). Anti-ulcerogenic effect of banana powder (*Musa spientum* var. *paradisiaca*) and its effect on mucosal resistance. *Journal of Ethnopharmacology*, 18(1), 33-34.
- Goel, R.K., Tavares, I.A. and Bennet, A. (1989). Stimulation of gastric and colonic mucosal eicosanoid synthesis by plantain banana. *Journal Pharmacy and Pharmacology*, 41 (11), 747-750.
- Guth, P.H. (1972). Gastric blood flow in restraint stress. *Digestive Disease and Science*, 17, 807-813.
- Guth, P.H., Paulsen, G. and Nagata, H. (1984). Histological and microulatory changes in alcohol-induced gastric lesions in the rat: Effect of prostaglandin cytoprotection. *Gastroenterology*, 87, 1183-1190.
- Hirschhon, H.H. (1983). Botanical remedies of the former Dutch (Indonesia). I: Eumycetes, Pteridophyta, Gtmnospermac,

Angiospernal Monocotylendenes only). *Journal of Ethnopharmacology.*, 7(2), 123-156.

Hong, N.D., Kim, J.W., Jeong, G.M. and Kim, N.J. (1982).

Studies on the efficacy of combined preparation of crude drugs (9). Effect of "Jakyakgamaho Tang" on anti-inflammatory and antiulcerative actions, and motility of isolated intestine. *Korean Journal Pharmacy*, 13, 87-91.

Howden, C.W., Jones, D.B., Peace, K.E., Burget, D.W. and Hunt, R.H. (1988). The treatment of gastric ulcer with antisecretory drugs: relation of pharmacological effects to healing rates. *Disease of the Digestive System*, 33, 619-624.

Itokawa, H., Mihashi, S., Watanabe, K., Natsumoto, H. and Hamanaka, T. (1983). Studies on the constituents of crude drugs having inhibitory activity against contraction of the ileum caused by histamine or bariumchloride (1). Screening test for the activity of commercially available crude drugs and the related plant materials. *Shoyakugaku Zasshi*, 37(3), 223-228.

Joshi, D.I., Dikshit, R.H. and Mansuri, S.M. (1987). Gastrointestinl action of garlic oil. *Phytotherapy Reserch*, 13, 140-141.

- Kanjanapothi, D., Panthong, A., Taesotikul, T., Rujjanawate, C., Kaewpinit, D., Lertprasertsuke, N., Choochate, W., Chaitong, U., Tookyang, B. and Pakdichareon, A. (1997). Plants for control of mosquito borne disease vectors. *A final report submitted to Chulabhorn Research Institute*, Bangkok, Thailand
- Key, T.D. (1996). *Chinese herbs (Their Botany, Chemistry and Pharmacodynamic)*. Tokyo, Charle E. Tuttle Company Inc.
- Kitagawa, H., Fujwara, M. and Osumi, V. (1979). Effect of water immersion stress on gastric secretion and mucosal blood flow in rats. *Gastroenterology*, 77, 298-302.
- Kitazawa, E., Sato, A., takahashi, S., Kuwano, H. and Ogiso, A. (1980). Novel diterpenelactones with anti-peptic ulcer activity from *Croton sublyratus*. *Chemical Pharmacological Bulletin*, 28, 227-234.
- Kiuchi, F., Nakamura, N. and Tsuda, Y. (1987). 3-Caren-5-one from *Kaempferia galanga*. *Phytochemistry*, 26(12), 3350-3351.
- Kiuchi, F., Hioki, M., Nakamura, N., Miyashita, N., Tsuda, Y. and Kondo, K. (1989). Screening of crude drugs used in Sri Langka for nematocidal activity on the larva of *Toxixoria canis*. *Shoyakugaku Zasshi*, 43(4), 288-293.

- Kobayashi, S. and Tabata, K. (1992). Pharmacology of plannitol, an anti-ulcer agent extract from a Thai medicinal plant (plau-noi). *JSPS-NRCT 1<sup>st</sup> Joint Seminar: Advance in Research on Pharmacologically Active Substances from Natural Sources*, Chiang Mai, Thailand, 3-5 December 1992: 23.
- Konturex, S.J., Obtalowicz, W., Kwiecion, N. and Oleksy, I. (1984). Generation of prostaglandins in gastric mucosa of patients with peptic ulcer disease: Effect of non steriodal anti-inflammatory compounds. *Scandinavian Gastroenterology*, 19, 75-77.
- Koo, M.W.L., Ogle, C.M. and Cho, C.H. (1986). Effect of verapamil, carbenoxolone and N-acetylcysteine on gastric wall mucus and ulceration in stressed rats. *Pharmacology*, 32, 236-334.
- Kosuge, T., Yokota, M., Sugiyama, K., Saito, M., Iwata, Y., Nakura, M. and Yamamoto, T. (1985). Studies on anticancer principles in Chinese medicines II. Cytotoxic principles in *Biota orientalis* L. Endl and *Kaempferia galanga* L. *Chemical Pharmacological Bulletin*, 33(12), 5565-5567.
- Kumar, D.S. and Prabhakar, V.S. (1987). On the ethnomedical significance of the Arjun Tree, *Terminalia Arjuna* (Roxb.)

Weight & Amot. *Journal of Ethnopharmacology*, 20 (2), 173-190.

Lange, K., Peskar, B.A. and Peskar, B.M. (1985). Stimulation of rat mucosal leukotriene formation by ethanol. *Naunyn Schmiedebergs Archive for Pharmacology*, 27, 3305.

Madulid, D.A., Gaealan, F.S.M., Romero, E.M. and Agoo, E.M.G. (1989). Ethnopharmacological study of the anti-trible in Nagpana, Barotac Viejo, Iloilo. *Acta Manilana.*, 38 (1), 25-40.

Maity, S., Vedasiromoni, J.R. and Ganguly, D.K. (1995). Anti-ulcer effect of the hot water of black tea (*Camellia sinensis*). *Journal of Ethnopharmacology* 46, 167-174.

Marhuenda, E., Martin, M. J., Alarcon dela Lastra, C. (1993). Antiuclerogenic activity of aescine in different experimental models. *Phytotherapy Research*, 7, 13-16.

Merh, P.S., Daniel, M. and Sabnis, S.D. (1986). Chemistry and taxonomy of some members of the Zingiberales. *Current Science.*, 55 (17), 835-839.

Minano, J.F., Serano, J.S., Pascual, J. and Sancibrian, M. (1987). Effects of GABA on gastric acid secretion and ulcer formation in rats. *Life Science*, 41, 1651-1658.

Mitsui, S., Kobayashi, S., Naghori, H. and Ogiso, A. (1976). Constituents from seeds of *Alpinia galanga* and their

- antiulcer activities. *Chemical Pharmacological Bulletin*, 24, 2377.
- Mokkhasmit, M., Swatdimongkol, K. and Satrawatia, P. (1971). Study on toxicity of Thai medicinal plants. *Bulletin of Dept Medical Scinces*, 12, 936-965.
- Mukhopadhyaya, K., Ehatattacharya, D., Chakraborty, A., Goel, R.K. and Sanyal, A.K. (1987). Effect of banana powder (*Musa sapientum* var. *paradisiaca*) on gastric mucosal shadding. *Journal of Ethnopharmacology*, 21(1), 11-19.
- Murakami, M., Lam, S.K., Inada, M. and Miyake, T. (1985). Pathophysiology and pathogenesis of acute gastric mucosal lesions after hypothermic restraint stress in rats. *Gastroenterology*, 88, 660-665.
- Ogiso, A., Kitazawa, E., Kuraboyashi, M. et al., (1978). Isolation and structure of antipeptic ulcer diterpene from Thai medicinal plant. *Chemical Pharmacological Bulletin*, 26(10), 3117-3123.
- Ohya, Y. and Guth, P.H. (1988). Ethanol-induced gastric mucosal blood flow and vascular permeability change in the rat. *Digestive Disease and Sciences*, 33, 883-888.
- Paijmans, K.P. (1976). *New Guinea Vegetation*. New York. Elsevier Scientific Publ Co.

- Paiva, L.A.F., Rao, V.S.N., Gramosa, N.V., Silveira, E.R. (1998). Gastroprotective effect of *Copaifera langsdorffii-eleorisin* on experimental gastric ulcer models in rats. *Journal of Ethnopharmacology* 62, 73-78.
- Prucksunand, C., Indrasukhisri, B., Leethochawalit, M., Nilvises, N., Prijavudhi, A. and Wimolwatthapun, S. (1986). Effect of the long turmeric (*Curcuma longa* Linn.) on healing of peptic ulcer: A preliminary report of 10 case study. *Thai Journal of Pharmacology*, 8(3), 139-151.
- Rafatullah, S., Tariq, M., Al-Yahya, M.A., Mossa, J.S. and Ageel, A.M. (1990). Evaluation of tuemeric (*Curcuma longa*) for gastric and duodenal antiulcer activity in rat. *Journal of Ethnopharmacology*, 29(1), 25-34.
- Rhee, J.C. and Chol, K.W. (1981). Clinical effect of the triturated extract of *Centulla asiatica* (madecassol) on the peptic ulcer. *Korean Journal Gastroenterology*, 13 (1), 35-40.
- Richardson, C.T. (1989). Gastric ulcer. In M.H. Sleisenger and J.S. Fordtran (eds.), *Gastrointestinal Disease Pathophysiology/ diagnosis/ management* (4<sup>th</sup> ed., pp. 879-925). Philadelphia, W.B. Saunders.
- Robert, A., Nezamine, J.E., Lancaster, C. and Honcher, A.J. (1979). Cytoprotective by prostaglandins in rats: Prevention of gastric necrosis produced by alcohol, HCl.

NaOH, hypertonic NaCl and Thermal injury.

*Gastroenterology*, 77, 433-443.

Robert, A., Nezemis, J.E., Lancaster, C., Davis, J.P. and Field, S.D. (1983). Mild irritants prevent gastric necrosis through "adaptive cytoprotection" mediated by prostaglandins. *American Journal of Physiology*, 245, 113-121.

Sanyal, A.K., Das, P.K., Sinha, S. and Sinha, Y.K. (1961). Banana and gastric secretion. *Journal of Pharmacy and Pharmacology*, 13, 318-319.

Sanyal, A.K., Gupta, K.K. and Chowdhury, N.K. (1964). Studies on peptic ulceration, Part I: Role of banana in phenylbutazone induced ulcers. *Archives Internationales de Pharmacodynamie et de Therapie*, 149, 393-400.

Sanyal, A.K., Banerjee, C.R. and Dos P.K. (1965). Studies on peptic ulceration, Part II: Role of banana in restraint and prednisolone induced ulcer in albino rats. *Archives Internationales de Pharmacodynamie et de Therapie*, 155, 244-248.

Shay, H., David, C.H., Sun, M.D. and Gruensteil, M. (1945). A sample method for the uniform production of gastric ulceration in the rat. *Gastroenterology*, 5, 43-61.

Shin, H.S., Choi, I.C., Lee, M.H. and Park, K.N. (1982). Clinical trials of madecassol (*Centulla asiatica*) on gastrointestinal

ulcer patient. *Korean Journal of Gastroenterology*, 14(1), 49-56.

Shinha, M., Mukherjee, B.P., Muknerjeee, B., Sikdar, S. and Dasgupta, S.P. (1975). Study of the mechanism of action of curcumin: on antiulcer agent. *Indian Journal of Pharmacology*, 7, 98-99.

Singh, T.J., Dos Gupta, P., Khan, S.Y. and Mishra, K.C. (1970). Preliminary pharmacological investigations of *Ocimum sanctum*. *Indian Journal Pharmacology*, 32, 92-94.

Singh, Y.N. (1986). Traditional medicine in Fiji: some herbal folk cures used by Fiji Indians. *Journal of Ethnopharmacology*, 15 (1), 57-58.

Staff of the Department of Pharmacology University of Edinbergh. (1970). *Pharmacological experimental on isolated preparations*. pp. 2, 96-97, 112-113. NewYork Churchill Livingstone Inc.

Takagi, T., Kasuya, Y. and Watanabe, K. (1963). Studies on the drugs of peptic ulcer: A reliable method for producing stress ulcer in rats. *Chemical Pharmacological Bulletin*, 12, 465-472.

Takeuchi, K., Okada, M., Niida, H. and Okabe, S. (1989). Role of sulphhydryls in mucosal injury caused by ethanol: relation to microvascular permeability, gastric motility and

- cytoprotection. *Journal of Pharmacology and Experimental Therapeutics*, 248, 836-841.
- Tookyang, B. (1997). *Plant extracts for control of mosquito-borne disease vector*. Ph.D. Thesis, Chiang Mai University, Thailand.
- Watanabe, K. (1966). Some pharmacological factors involved in formation and prevention of stress ulcers in rats. *Chemical Pharmaceutical Bulletin*, 14, 101-107.
- Yamahara, J., Kobayashi, M., Miki, K., Kozuka, M., Sawada, T. and Fujimura, H. (1985). Cholagogic and anti-ulcer effect of *Saussureae radix* and its active components. *Chemical Pharmacological Bulletin*, 33(3), 1285-1288.
- Yamahara, J., Mochizuki, M., Fujimura, F., Takaishi, Y., Yoshida, M., Tomimatsu, T. and Tamai, Y. (1988). The anti-ulcer effect in rats of ginger constituents. *Journal of Ethnopharmacology*, 23, 299-304.
- Yoshikawa, M., Hakeyama, S., Taniguchi, K., Matuda, H. and Yamahara, J. (1992). 6-Gingesulfonic acid, a new anti-ulcer principle, and gingerglycolipids A, B, and C, three new monoacyldigalactosylglyceral from *Zingiberis rhizoma* originating in Taiwan. *Chemical Pharmacological Bulletin*, 40(8), 2239-2241.