

TABLE OF CONTENTS

	page
ACKNOWLEDGEMENT	iii
ENGLISH ABSTRACT	iv
THAI ABSTRACT	v
TABLE OF CONTENTS	vii
LIST OF TABLES	ix
LIST OF FIGURES	x
ABBREVIATION	xii
CHAPTER 1 INTRODUCTION	1
1.1 Statement and significance of the problem	1
1.2 Objective	2
CHAPTER 2 LITERATURE REVIEWS	3
2.1 Structure and function of human hemoglobin	3
2.2 Molecular genetics and expression of the human α -globin gene cluster	7
2.3 Alpha-thalassemia	9
2.3.1 Non-deletional α -thalassemia	9
2.3.2 Gene deletion	11
2.3.3 Hemoglobin H disease	14
2.3.4 Hemoglobin Bart's hydrop fetalis syndrome	15
2.4 Polymerase chain reaction	17
2.5 Standard PCR amplification protocol	20

2.6 Optimization of PCR	21
2.7 Principle of preimplantation genetic diagnosis	24
2.8 Principle of primer-extension-preamplification	25
CHAPTER 3 METHODS	26
CHAPTER 4 RESULTS	37
CHAPTER 5 DISCUSSION AND CONCLUSION	50
REFERENCES	55
APPENDIX	62
Appendix A	62
Appendix B	64
VITA	67

LIST OF TABLES

Table	page
1 Normal human hemoglobins and their globin subunits	4
2 Nondeletional α -thalassemia mutations	10
3 Mutations of the α -globin genes found in Thailand	16
4 Optimization of PCR components	30
5 Detection of the α -thalassemia-1 (SEA) haplotype by PCR in prospective screening of expecting couples which had EOF values <60%	44
6 Detection of the α -thalassemia-1 of (SEA) haplotype by PCR from amniotic fluids	44

LIST OF FIGURES

Figure	page
1 The locations of the β -globin gene group on chromosome 11 and the α -globin genes on chromosome 16	4
2 The changes in hemoglobin composition during gestation and development	6
3 The organization of the α -globin gene complex	7
4 The duplicated xyz box arrangement containing the α -globin genes	12
5 The major deletions in α -thalassemia-1	13
6 The polymerase chain reaction	19
7 Diagram of step 1 procedure	26
8 The position of the primers in the α -globin gene cluster for detection of the α -thalassemia-1 (SEA) haplotype by PCR	29
9 Diagram of step 4 and 5 procedure	34
10 Comparison of the amplification using the conventional PCR without glycerol and PCR with glycerol at concentration 5, 10 and 15%, respectively	37
11 The PCRs were performed with annealing temperatures at 58, 59 and 60°C, respectively	38
12 The PCRs were performed at various $MgCl_2$ concentrations : 1.5, 1.75, 2.0, 2.25, 2.5 and 2.75 mM, respectively	39

13	Amplification for 40 cycles of PCR was performed with the concentration of each the primers A+B+C at 0.125, 0.25, 0.5, 0.75 and 1.0 μ M, respectively	40
14	The diluted <i>Taq</i> DNA polymerase from in house produced enzyme was varied at 0.06, 0.12 and 0.25 units in 10 μ l of reaction	41
15	The PCRs were performed for 30, 35 and 40 cycles, respectively	42
16	Detection of the α -thalassemia-1 (SEA) haplotype in 10 antenatal care women by PCR	43
17	Sensitivity test of DNA amplification for the α -thalassemia-1 (SEA) haplotype	46
18	Shows the activities of the BSA and the proteinase K by 12% SDS-PAGE	47
19	Shows the DNA bands obtained from the amniocytes amplification	48

ABBREVIATIONS

A	adenine
α	alpha
β	beta
bp	base pair
C	cytosine
δ	delta
γ	gamma
G	guanine
Hb	hemoglobin
kb	kilobase
ψ	psi
rpm	revolution per minute
T	thymine
θ	theta
ζ	zeta