

REFERENCES

1. Vazquez M.J., Perez-Marcos B., Gomez-Amoza J.L., Martinez-Pacheco R., Souto C. and Concheiro A., Influence of technological variables on release of drug from hydrophilic matrices. *Drug Dev. Ind. Pharm.* 1992; 18(11&12) : 1355-1375
2. Alan W. and Stephen P.C., Pentoxifylline – A review of its pharmacodynamic and pharmacokinetic properties and its therapeutic efficacy. *Drug* 1987; 34 : 50 – 97
3. Budavari S. et al., *Merck Index*, 11th ed. DC : Merck and Co.Inc., 1998
4. Gerald K.M., Editor, *AHFS Drug Information 96*. MD : American Society of Health System Pharmacist, Inc., 1996
5. Makoto O. and Yoshihida M., Programable drug release of highly water-soluble pentoxifylline from dry – coated wax matrix. *J.Pharm.Sci.* 1995; 84(4) : 443-446
6. Gleixner K., Muller R., Lehrach F., “Pharmaceutical compositions” *U.S. Patent 4189469* Feb. 19 , 1980
7. Salsa T., Veiga F. and Pina M.E., Oral controlled – release dosage forms. I. cellulose ether polymers in hydrophilic matrices. *Drug Dev. Ind. Pharm.* 1997; 23(9) : 929-938
8. Longer M.A. and Robinson J.R, Sustained-release drug delivery systems, in *Reminton's Pharmaceutical Sciences*, 18th ed. NY : Mark Easton, 1990, p. 1676

9. Robinson J.R. and Vincent H.L. *Controlled Drug Delivery Fundamentals and Applications*. 2nd ed. NY : Marcel Dekker, Inc., 1987, p. 1-16
10. Ping I.L. and William R.G., *Controlled-Release Technology Pharmaceutical Applications*. DC : American Chemical Society, 1987
11. Capan Y., Influence of technological factors on formulation of sustained release tablets. *Drug Dev. Ind. Pharm.* 1989; 15(6&7) : 927-956
12. Higuchi T., Mechanism of Sustained-Action Medication: Theoretical analysis of rate of release of solid drugs dispersed in solid matrices. *J.Pharm.Sci* 1963; 52(12) : 1145-1149
13. Khanna S.C. and Speiser, In vitro release of chloramphenicol from polymer beads of alpha-methacrylic acid and methylmethacrylate. *J.Pharm.Sci* 1970; 59(12) : 1398-1401
14. McGinity J.W., Cameron C.G. and Cuff G.W., Controlled release theophylline tablet formulation containing acrylic resin. part1: dissolution properties of tablets. *Drug Dev. Ind. Pharm.* 1983; 9(1-2) : 57-68
15. Carli F., Capone G., Colombo I., Magarotto L. and Motto A., Surface and transport properties of acrylic polymer influencing drug release from porous matrices. *Int. J. Pharm.* 1984; 21(Oct) : 317-329
16. Nakano M., Ohmori N., Ogata A., Sukimoto K., Tobino Y., Iwaoku R. and Juni K., Sustained release of theophylline from hydroxypropylcellulose tablets. *J.Pharm.Sci* 1983; 72(7) : 378-380

17. Desai S.J., Simonelli A.P. and Higuchi W.I., Investigation of factors influencing release of solid drug dispersed in inert matrices. *J. Pharm.Sci* 1965; 54(10) : 1459-1464
18. Daly P.B., Davis S.S. and Kennerley J.W., Effect of anionic surfactants on the release of chlorpheniramine from a polymer matrix tablet. *Int. J. Pharm.* 1984; 18(Jan) : 201-205
19. Lapidus H and Lordi N.G., Drug release from compressed hydrophilic matrices *J. Pharm.Sci* 1968; 57(8) : 1292-1301
20. Huber H.E. and Christenson G.L., Utilization of hydrophilic gums for the control of drug substance release from tablet formulations. II. Influence of tablet hardness and density on dissolution behavior. *J. Pharm.Sci* 1968; 57(1) : 164-167
21. Jambhekar S.S. and Cobby J., pH- and flow-rate-independent release of drug from uncoated slow-release tablets. *J. Pharm.Sci* 1985; 74(9) : 991-994
22. Skoug J.W., Mikelsons M.V., Vigneron C.N. and Stemm L., Qualitative evaluation of the mechanism of release of matrix sustained release dosage forms by measurement of polymer release. *J. Control. Rel.* 1993; 27(Dec) : 227-245
23. Abdou M.H , *Dissolution Bioavailability and Bioequivalence*, Pensilvania : Mack Printing Company, 1989
24. Foster T.P. and Parrott E.L., Release of highly water-soluble medicinal compounds from Inert, heterogeneous matrix. I: physical mixture, *J. Pharm.Sci* 1990; 79(9) : 806-810
25. The United State Pharmacopeial Committee. *USP DI 1997 Vol.1*: Drug Interaction for

the healthcare Professional, Massachusets : Ran MacNally, 1997

26. Kim N.S., Kawata M., Uchida T and Goto S., Preparation and evaluation of eudragit gels part3. rectal gel preparations for sustained release of pentoxifylline.
J.Pharm.Sci 1992; 81(9) : 537-540
27. Makoto O. and Yoshihida M., Controlled drug release of highly water soluble pentoxifylline from time limit disintegration type wax matrix tablet. *Pharm. Res.* 1994; 11 : 351-354
28. Baverova K., Rak J., Chalabala M., Heinrich J. and Tyrackova N., Study of in vitro kinetics and liberation mechanism of pentoxifylline from coated pellets and compacts base on 2-hydroxyethylmethacrylate-butyl acrylate copolymers. *Drug Dev. Ind. Pharm.* 1988; 14(15-17) : 2477-2497
29. Rynzo I., Hiroyuki Y., Yochiyuki H. and Kazno N., Dessign and preparation on pulsatile release tablet as a new oral drug delivery system. *Chem. Pharm.Bull.* 1992; 40(11) : 3036-3041
30. Makoto O. and Yoshihida M., Effect of cogrinding time on the release of pentoxifylline from wax matrix tablets. *J.Pharm.Sci* 1994; 83(7) : 956-960
31. Banakar U.V., *Materials Used in Controlled Release Technology*. MO : Technomic Publishing Co., Inc., 1994
32. Ainley W., Paul J.W. *Handbook of Pharmaceutical Excipients*. 2th Edition.
Washington : American Pharmaceutical Association., 1994

33. Nanette R.R. and Evone S.G., Mechanism of drug release from matrices prepared with aqueous dispersion of ethylcellulose. *Drug Dev. Ind. Pharm.* 1997; 23(1) : 113-117
34. Marini J.O., Mendes R.W. Rekhi J.S. and Jambhekar S.S., Some factors affecting the release of drug from membrane coated slow release tablets. *Drug Dev. Ind. Pharm.* 1991; 17(6) : 865-877
35. Klinger G.H., Ghaly E.S., Porter S.C. and Schwartz J.S., Formulation of controlled release matrices by granulation with a polymer dispersion. *Drug Dev. Ind. Pharm.* 1990; 16(9) : 1473-1490
36. Porter S.C., Controlled-release film coatings base on ethylcellulose. *Drug Dev. Ind. Pharm.* 1989; 15(9) : 1495
37. Document from Colorcon Methocel
38. Chowdary K.P. and Sankar G.G., Eudragit microcapsules of nifedipine and its dispersions in HPMC-MCC: physicochemical characterization and drug release studies. *Drug Dev. Ind. Pharm.* 1997; 23(3) : 325-330
39. Pabon C.V., Frutos P., Lastres J.L. and Frutos G., In vitro study of mixed controlled release matrix tablets containing HMPC and polyamide 12. *Drug Dev. Ind. Pharm.* 1992; 18(20) : 2163-2171
40. Sheskey P.J. and Cabelka T.D., Reworkability of sustained release tablet formulations containing HPMC polymers. *Pharm. Technol.* 1992; 16(Jul) : 60, 62, 64, 66, 68, 70, 72, 74

41. Harcum W.W., Skinner G.W. Altekarr M., Joneja S.K. Barnum P.E. and Guo J.H.,
Modelling the effects of hydroxypropylcellulose in acetaminophen tablet
formulation. *Drug Dev. Ind. Pharm.* **1998**; 24(10) : 911-918
42. Senel S., Capan Y., Dalkara T., Inanc N. and Hincal A.A., Formulation bioavailability
and pharmacokinetics of sustained release potassium chloride tablets. *Pharm. Res.*
1991; 8(Oct) : 1313-1317
43. Ebube N.K., Hikal A.H., Wyandt C.M., Beer D.C., Jones A.B., et al., Effect of drug,
formulation, and process variables on granulation and compaction characteristics of
heterogeneous matrices. part 2. HPMC and PVP systems. *Drug Dev. Ind. Pharm.*
1996; 22(7) : 561-567
44. Bansal P., Haribhakti K., Subramanian, V. and Plakogiannis F., Effect of formulation
and process variables on the dissolution profile of naproxen sodium from tablets.
Drug Dev. Ind. Pharm. **1994**; 20(13) : 2151-2156
45. Lieberman A.H., Lechman L. and Schwartz J.B. *Pharmaceutical Dosage Forms:
Tablets Volume 2*. New York : Marcel Dekker, Inc., 1990
46. Ford J.L. and Mitchell K., Molecular Interactions Between Water and HPMC Polymers.
Thermochim Acta. **1995**; 248 : 329-345
47. Doelker, E.L., Water-swollen cellulose derivatives in pharmacy. In Peppas, N.A.(Ed.).
Hydrogels in Medicine and Pharmacy, Vol 11: Polymers. CRC Press, Boca
Raton, FL. 1987

48. Ford. J.L., Rubinstien. M.H. and Hogan J.E., Formulation of sustained release promethazine hydrochloride tablets using HPMC matrices. *Int.J. Pharm.* 1985; 24 : 327-338
49. Gao. P., Nixon P.R. and Skoug J.W., Diffusion in HPMC gels. II. prediction of drug release rates from hydrophilic matrix extended-release dosage forms. *Pharm. Res.* 1995; 12: 965-971
50. Gao. P., Skoug J.W., Nixon P.R., Ju. T.R., Stemm. N.L. and Sung K.C., Swelling of HPMC matrix tablets. II. mechanic study of the influence of formulation variables on matrix performance and drug release. *J.Pharm.Sci.* 1996; 85 : 732-740
51. Sung K.C., Nixon P.R., Skoug J.W., Ju. T.R., Gao. P., Topp E.M. and Patel M.V., Effect of formulation variables on drug and polymer release from HPMC-based matrix tablets. *Int.J. Pharm.* 1996; 142 : 53-60
52. Lieberman A.H., Lechman L. and Schwartz J.B., *Pharmaceutical Dosage Forms: Tablets Volume I*. New York : Marcel Dekker, Inc., 1990
53. Fassihi A.R. and Kanfer I., "The effect of compressibility and powder flow properties on tablet weight variation" in *Pharmaceutical Technology: Tableting Technology Vol. I* . Westsussex, England : Ellis Horwood Ltd., 1987