

Table of contents

	Page
Acknowledgement	iii
Abstract (English)	iv
Abstract (Thai)	v
List of tables	viii
List of illustrations	ix
Chapter 1 Introduction	1
1.1 Historical Review	1
1.2 The Aim of the Present Work	7
1.3 Galactic Spiral Structure	8
Chapter 2 The Galactic Magnetic Field	12
2.1 Galactic Dynamo Theory	12
2.2 Regular and Irregular Field	13
2.3 Compression Profile	15
2.4 Synchrotron Radiation	19
2.5 Distribution of Electron Density	26
Chapter 3 Magnetic Field Model of Milky Way Galaxy	32
3.1 The Recent Model	32
3.2 The Adapted Model	41

	Page
Chapter 4 Methodology	44
4.1 408 MHz Allsky Survey	44
4.2 Methodology	47
Chapter 5 Discussion and Conclusion	55
5.1 Discussion and Conclusion	55
5.2 Future Work	59
References	60
Appendix	64
Curriculum vitae	67

List of tables

Table	Page
4.1 The four surveys at 408 MHz	45
4.2 Summary of all the equations and their parameters for Synchrotron Model.	53

List of illustrations

Figure	Page
2.1 The dependence of the regular magnetic field H_{reg} on the distance from the galactic plane at three distances from the galactic centre.	13
2.2 The regular magnetic field H_{reg} is shown in the plane perpendicular to the galactic plane, passing through the Sun and the galactic plane at several distances z from the plane.	14
2.3 The density of clouds is expressed relative to the average and marked underneath the curve is the full width of the arm amounting to the linear distance of 1 kpc.	17
2.4 The synchrotron spectrum from a single electron as a function of $x = \nu/\nu_c$ where ν_c is the synchrotron critical frequency.	21
2.5 The distribution of electron flux density with height above the plane at the solar galactic radius.	30
3.1 Radial variation of the magnitude of the regular component of magnetic field, $H_{reg}(R)/H_{reg}(R_{SUN})$ used by Kearsy (1983) but scaled the position of the Sun of 8.5 kpc.	33

Figure	Page
3.2 Radial variation of the magnitude of the regular component of magnetic field, $H_{reg}(R)/H_{reg}(R_{SUN})$ adapted from Han and qiao.	36
3.3 Bisymmetric spiral magnetic fields used for our model computations of RM.	38
3.4 Possible, two-armed, bisymmetric spirals fitted to the distribution of HII regions given by Georgelin and Georgelin (1976).	39
3.5 The adapted model	43
4.1 The coordinates of the galactic magnetic field show the component which perpendicular to the line of sight.	48
4.2 The calculated brightness temperature from the Adapted Model vary as the galactic longitudes in the range of -90° to 90° with the initial parameters.	49
4.3 The profile with adjusted $p = -6.8^\circ$ and $R_2 = 9.35$ kpc.	50
4.4 The profile with adjusted $p = -6.8^\circ$, $R_2 = 9.35$ kpc, $R_0 = 11.0$ kpc and $R_1 = 1.91$ kpc .	51
5.1 The calculated brightness temperature from Synchrotron Emission Model vary as the galactic longitude in the range of -90° to 90° .	56
5.2 The adapted model with the optimized parameters.	57
5.3 The calculated brightness temperature from the Adapted Model vary as the galactic longitude in the range of -90° to 90° .	58