

REFERENCES

- Abbas, A.K., Lichtman, A.H., Pober, J.S. (2000) Cellular and molecular immunology. W.B. Saunders. 39b-41b.
- Austin, AK., Hobbs, RN., Anderson, JC., Butler, RC., and Ashton BA. (1988) Humoral immunity to link protein in patients with inflammatory joint disease, osteoarthritis, and in non-arthritis controls. *Ann Rheum Dis.* 47, 886-892.
- Bajorath, J., Greenfield, B., Munro, S.B., Day, A.J., and Aruffo, A. (1998) Identification of CD44 residues important for hyaluronan binding and delineation of the binding site. *J Biol Chem.* 273, 338-343.
- Baker, J., Chritner, J. and Ekborg, S. (1991) An unsulphation region of the rat chondrosarcoma chondroitin sulphate chain and its binding to monoclonal antibodies 3B3. *Biochem. J.* 273, 237-239.
- Bank, R.A., Bayliss, M.T., Lafeber, F.P., Maroudas, A., and Tekoppele, J.M. (1998) Ageing and zonal variation in post-translational modification of collagen in normal human articular cartilage. The age-related increase in non-enzymatic glycation affects biomechanical properties of cartilage. *Biochem J.* 330, 345-351.
- Barta, E., Deak, F., and Kiss, I.(1993) Evolution of hyaluronan-binding module of link protein. *Biochem J.* 292,947-949.
- Bayliss, M.T., and Ali, S.Y. (1978) Age-related changes in the composition and structure of human articular-cartilage proteoglycans. *Biochem J.* 176,683-693.
- Benton, H.P., Jackson, T.R., and Hanley M.R. (1989) Identification of a novel inflammatory stimulant of chondrocytes, early events in cell activation by bradykinin receptors on pig articular chondrocytes. *Biochem J.* 258, 861-867.
- Bost, F., Diarra-mehrpor, M., and Martin J.P. (1998) Inter- α -trypsin inhibitor proteoglycan family, a group of proteins binding and stabilizing the extracellular matrix. *Eur. J. Biochem.* 252, 339-346.
- Bradford, M. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem.* 72, 248-254.

- Buschmann, M.D., Hunziker, E.B., Kim, Y.J., and Grodzinsky A.J. (1996) altered aggrecan synthesis correlates with cell and nucleus structure in statically compressed cartilage. *J Cell Sci.* 109, 499-508.
- Calabro, A., Hascall, V.C., and Caterson, B. (1992) Monoclonal antibodies directed against epitopes within the core protein structure of the large aggregating proteoglycan (aggrecan) from the swarn rat chondrosarcoma. *Arch Biochem Biophys.* 298, 349-360.
- Caterson, B. and Baker, J. (1979) The link proteins as specific components of cartilage proteoglycan aggregates in vivo (associative extraction of proteoglycan aggregate from swarn rat chondrosarcoma). *J. Biol. Chem.* 254, 2394-9.
- Caterson, B., Baker, J., Christner, J., Kearhey, J. and Stohrer, R. (1985) The characterization of clonal antibodies directed against bovine nasal cartilage proteoglycan and link protein. *Articular Cartilage Biochemistry.* Workshop Conference Hoechst- Werk Albertwiesbaden Sep 30-Oct 3. New York: Raven Press.
- Caterson, B., Baker, J.R., Christner, J.E., and Couchman, J.R. (1982) Immunological methods for the detection and determination of connective tissue proteoglycans. *J Invest Dermatol.* 79 Suppl, 45s-50s.
- Chandrosoma, P., and Taylor, C.R. (1998) a LANGE medical book, Concise pathology 3rd ed. Prentice-Hall International, Inc.
- Chuanbunyat, J. (2001) Monoclonal antibody hyaluronan binding proteins from shark cartilage. M.S. Thesis, Chiangmai University.
- Dahlberg, L., Ryd, L., Heinegard, D., and Lohmander, L.S. (1992) Proteoglycan fragments in joint fluid. Influence of arthrosis and inflammation. *Acta Orthop Scand.* 63,417-423.
- De Groot, J., Verzijl,N., and Blank, R.A. (1999) Age-related decrease in proteoglycan synthesis of human articular chondrocytes, the role of nonenzymatic glycation. *Arthritis Rheum.* 42,1003-1009.
- Dieppe, P. (1995) Osteoarthritis and molecular markers. A rheumatologist's perspective.*Acta Orthop Scand Suppl.* 266,1-5.
- Domowicz, MS., Pirok, E.W., Novak, T.E., and Schwartz, N.B. (2000) Role of the C-terminal G3 in Sorting and secretion of aggrecan core protein and ubiquitin-

- mediated degradation of accumulated mutant precursors. *J Biol Chem.* 275, 35098-35105.
- Dudhia, J., Davidson, C.M., Wells, T.M., Vynios, D.H., Hardingham, T.E., and Bayliss, M.T. (1996) Age-related changes in the content of the C-terminal region of aggrecan in human articular cartilage. *Biochem J.* 313 (Pt 3), 933-940.
- Eastell, R., Robins, S.P., Colwell, T., Assiri, A.M., Riggs, B.L., and Russell, R.G. (1993) Evaluation of bone turnover in type I osteoporosis using biochemical markers specific for both bone formation and bone resorption. *Osteoporos Int.* 3, 255-260.
- Engstrom-Laurent, A., and Hallgren, R. (1985) Circulating hyaluronate in rheumatoid arthritis: relationship to inflammatory activity and the effect of corticosteroid therapy. *Ann Rheum Dis.* 44, 83-88.
- Franzen, A., Inerot, S., Hejderup S.O., and Heinegard, D. (1981) Variations in the composition of bovine hip articular cartilage with distance from the articular surface. *Biochem J.* 195, 535-543.
- Garnero, P., Rousseau, J.-C., and Delmas, P.D. (2000) Molecular Basic and clinical use of biochemical markers of bone, cartilage, and synovium in joint diseases. *Arthritis Rheum.* 43, 953-968.
- Glant, T., Csongor, J., and Szucs, T. (1980) Immunopathologic role of proteoglycan antigens in rheumatoid joint disease. *Scand J Immunol.* 11, 247-252.
- Goldman, L., and Bennette, JC. (2000) Textbook of medicine, 21st ed: W.B. Saunders Company.
- Guerassimov, A., Zhang, Y., Cartman, A., Rosenberg, L.C., Esdaile, J., Fitzcharles, M.-A., and Poole, A.R. (1999) Immune responses to cartilage link protein and the G1 domain of proteoglycan aggrecan in patients with osteoarthritis. *Arthritis Rheum.* 42, 527-533.
- Guerne, P.A., Blanco, F., Kaelin, A., Desgeorges, A., and Lotz, M. (1995) Growth factor responsiveness of human articular chondrocytes in aging and development. *Arthritis Rheum.* 38, 960-968.
- Hardingham, T. (1981) Proteoglycan: Their structure, interaction and molecular organization in cartilage. *Biochem. Soc. Trans.* 9, 489-97.

- Hardingham, T. (1998) Cartilage: Aggrecan-link protein-hyaluronan aggregates[Online]. Available: <http://www.Glycoforum.gr.jp/science/hyaluronan/HA05/HA05E.html> [1999, December 25].
- Hardingham, T. and Fosang, A. (1992) Proteoglycans: many forms and many functions. *FASEB J.* 6, 861-70.
- Hedlund, H., Hedbom, E., Heinegard, D., Mengarelli-Widholm, S., Reinholt, FP., and Svensson , O. (1999) Association of the aggrecan keratan sulfate-rich region with collagen in bovine articular cartilage. *J Biol Chem.* 274, 5777-5781.
- Heinegard, D. (1977) Polydispersity of cartilage proteoglycans. Structural variations with size and buoyant density of the molecules. *J. Biol. Chem.* 252, 1980-9.
- Heinegard, D., Inerot, S., Wieslander, J., and Lindblad, G. (1985) A method for the quantification of cartilage proteoglycan structures liberated to the synovial fluid during developing degenerative joint disease. *Scand J Clin Lab Invest.* 45, 421-427.
- Hochberg, M., Altman, R., Brandt, K., Clark, B., Dieppe, P., Griffin, M., Moskowitz, R. and Schnitzer, T. (1995) Guidelines for the medical management of osteoarthritis. Part II Osteoarthritis of the knee-American College of Rheumatology. *Arth. Rheum.* 38, 1541-1546.
- Hogervorst, T., and Brand, R.A. (1998) Mechanoreceptors in joint function. *J Bone Joint Surg.* 80A, 1365-1378
- Kempson, G.E., Muir, H., Pollard, C., and Tuke, M. (1973) The tensile properties of the cartilage of human femoral condyles related to the content of collagen and glycosaminoglycans. *Biochim Biophys Acta.* 297, 465-472.
- Kiani, C., Lee, V., Cao, L., Chen, L., Wu, Y., Zhang, Y., Adams, M.E., and Yang, B.B. (2001) Roles of aggrecan domains in biosynthesis, modification by glycosaminoglycans and product secretion. *Biochem. J.* 354, 199-207.
- Kida, D., Yoneda, M., Miyaura, S., Ishimaru, T., Yoshida, Y., Ito, T., Ishiguro, N., Iwata, H., and Kimata, K. (1999) The SHAP-HA complex in sera from patients with rheumatoid arthritis and osteoarthritis. *J Rheumatol.* 26, 1230-1236.
- Knutson, J.R., Iida, J., Fields, G.B., and McCarthy, J.B. (1996) CD44/chondroitin sulfate proteoglycan and alpha 2 beta 1 integrin mediate human melanoma cell migration

- on type IV collagen and invasion of basement membranes. *Mol Biol Cell.* 7, 383-96.
- Kobayashi, H., Hirashima, Y., Sun, G.W., Fujie, M., Nishida, T., Takigawa, M., and Terao, T. (2000) Identity of urinary trypsin inhibitor-binding protein to link protein. *J Biol Chem.* 275, 21185-21191.
- Kohda, D., Morton, C.J., Parker A.A., Hatanaka, H., Inagaki, F.M., Campbell, I.D., and Day A.J. (1996) Solution structure of the link module: a hyaluronan-binding domain involved in extracellular matrix stability and cell migration. *Cell.* 86, 767-775.
- Koopman, W.J. (2001) Arthritis and allied conditions, a textbook of rheumatology 14th edition vol. 1., Lippincott Williams & Wilkins.
- Lohmander, L.S., Neame, P.J., and Sandy, J.D. (1993) The structure of aggrecan fragments in human synovial fluid. Evidence that aggrecanase mediates cartilage degradation in inflammatory joint disease, joint injury, and osteoarthritis. *Arthritis Rheum.* 36, 1214-1222.
- Mankin, H.J., Mow V.C., and Buckwalter J.A. (1994) Form and function of articular cartilage. Orthopaedic basic science. American of Orthopaedic Surgeons.
- Maroudas, A., Bayliss, M.T., Uchitel-Kaushansky, N., Schneiderman, R., and Gilav, E. (1998) Aggrecan turnover in human articular cartilage: use of aspartic acid racemization as a marker of molecular age. *Arch Biochem Biophys.* 350, 61-71.
- McKenna, L.A., Lui, H., Sansom, P.A. and Dean, M.F. (1998) An N-terminal peptide from link protein stimulates proteoglycan biosynthesis in human articular cartilage *in vitro*. *Arthritis Rheum.* 41, 157-161.
- McKibben, B., and Holdsworth, F.S. (1966) The nutrition of immature joint cartilage in the lamb. *J Bone Joint Surg.* 48B, 793-803.
- Microsoft corporation (2001) Encarta.msn.com. Joint damage caused by arthritis [Online]. Available://<http://encarta.msn.com/find/MediaMax.asp?pg=3&ti=761561517&idx=461547526>.
- Mikecz, K., Glant, T.T., and Poole, A.R. (1987) Immunity to cartilage proteoglycans in BALB/c mice with progressive polyarthritis and ankylosing spondylitis induced by injection of human cartilage proteoglycan. *Arthritis Rheum.* 30, 306-318.

- Moos, V., Rudwaleit, M., Herzog, V., Hohlig, K., Sieper, J., and Muller, B. (2000) Association of genotypes affecting the expression of interleukin-1 β or interleukin 1 receptor antagonist with osteoarthritis. *Arthritis Rheum.* 43, 2417- 2422.
- Moriuchi-Murakami, E., Yamada, H., Ishi, O., Kikukawa, T., and Igarashi, R. (2000) Treatment of established collagen induced arthritis with prostaglandin E₁ incorporated in lipid microspheres. *J Rheumatol.* 27, 2389-96.
- Muir, I.H.M. (1979) Biochemistry. Adult articular cartilage. 2nd ed. Pitman Medical. 145-214.
- Neame, P.J., and Barry, F.P. (1993) The link protein. *Experientia.* 49, 393-402.
- Parkar, AA, Kahmann, J.D., Howat, S.L., Bayliss, M.T., and Day, A.J. (1998) TSG-6 interacts with hyaluronan and aggrecan in a pH-dependent manner via a common functional element: implications for its regulation in inflamed cartilage. *FEBS Lett.* 29, 171-176.
- Petersson, IF., Boegard, T., Svensson, B., Heinegard, D., and Saxne, T. (1998) Changes in cartilage and bone metabolism identified by serum markers in early osteoarthritis of the knee joint. *Br J Rheumatol.* 37, 46-50.
- Plass, A.H., Wong-Plams, S., Roughley P.J., Midura, R.J., and Hascall, V.C. (1997) Chemical and immunological assay of the nonreducing terminal residues of chondroitin sulfate from human aggrecan. *J Biol Chem.* 272, 20603-20610.
- Poole, A.R. (1993) Cartilage in health and disease. In: *Arthritis and allied conditions.* 12th ed. Philadelphia: Lea & Fabiger.
- Poole, A.R. Witter, J., Roberts, N., Piccolo, F., Brandt, R., Paquin, J., and Baron, M. (1990) Inflammation and cartilage metabolism in rheumatoid arthritis. *Arthritis Rheum.* 790-799.
- Poole, A.R., Ionescu, M., Swan, A., and Dieppe, P.A. (1994) Changes in cartilage metabolism in arthritis are reflected by altered serum and synovial fluid levels of the cartilage proteoglycan aggrecan, Implications for pathogenesis. *J Clin Invest.* 94, 25-33.
- Poole, A.R., Pidoux, I., Reiner, A., and Rosenberg, L. (1982) An immunoelectron microscope study of the organization of proteoglycan monomer, link protein, and collagen in the matrix of articular cartilage. *J Cell Biol.* 93, 921-937.

- Poole, A.R., Webber, C., Pidoux, I. (1986) Localization of a dermatan sulfate proteoglycan (DS-PGII) in cartilage and the presence of an immunologically related species in other tissues. *J Histochem Cytochem.* 34, 619-625.
- Porath, J., Maisano, F., and Belew, M. (1985) Thiophic adsorption –a new method for protein fractionation. *FEBS Lett.* 185, 306-310.
- Pothacharoen, P. The quantitative analysis of chondroitin sulfate epitopes and hyaluronan as diagnostic markers for degenerative joint disease by ELISA technique. M.S. Thesis, Chiangmai University.
- Radhakrishnamurthy, B., Jeansson, N., Tracy, RE., and Berenson, GS. (1993) A monoclonal antibody that recognizes hyaluronic acid binding region of aorta proteoglycans. *Atherosclerosis.* 98, 179-92.
- Roughley, P.J., White, R.J., Poole, A.R. (1985) Identification of a hyaluronic acid-binding protein that interferes with the preparation of high-buoyant-density proteoglycan aggregates from adult human articular cartilage. *Biochem J.* 231, 129-138.
- Saxne, T., Heinegard, D., Wollheim, F.A., and Petersson, H. (1985) Difference in cartilage proteoglycan level in synovial fluid in early rheumatoid arthritis and reactive arthritis. *Lancet.* 2, 127-128.
- Seeley, R.R., Stephens, T.D., and Tate, P. (1999) Essentials of anatomy and physiology. 3rd ed. WCB /McGraw-Hill.
- Shikhman, AR., Brinson, DC., and Lotz, M. (2000) Profile of glycosaminoglycan-degrading glycosidases and glycoside sulfatases secreted by human articular chondrocytes in homeostasis and inflammation. *Arthritis Rheum.* 43, 1307-1314.
- Stockwell, R.S. Biology of cartilage cells. Cambridge, Cambridge University Press.
- Tengblad, A. (1979) Affinity chromatography on immobilized hyaluronate and its application to the isolation of hyaluronate binding proteins from cartilage. *Biochem. Biophys. Acta.* 578, 281-289.
- Tiengburanatam N. Production and characterization of monoclonal antibody against chondroitin sulfate. M.S. Thesis, Chiangmai University.
- Turley, E.A., Austen, L., Vandeligt, K., and Clary, C. (1991) Hyaluronan and a cell-associated hyaluronan binding protein regulate the locomotion of ras-transformed cells. *J Cell Biol.* 112, 1041-1047.

- Van den Berg, WB., Joosten, LAB., Kollias, G., and Van de Loo, AAJ. (1999) Role of tumor necrosis factor α in experimental arthritis: separate activity of interleukin 1 β in chrolicity and cartilage destruction. *Ann Rheum Dis.* 58 (S I), 140-148.
- Vilim, V., and Fosang, A.J. (1994) Proteoglycans isolated from dissociative extracts of differently aged human articular cartilage: characterization of naturally occurring hyaluronan-binding fragments of aggrecan. *Biochem J.* 304, 887-894.
- Watanabe, H., Cheung, S.C., Itano, N., Kimata, K., and Yamada, Y. (1997) Identification of hyaluronan-binding domains of aggrecan. *J Biol Chem.* 272, 28057-28065.
- Webber, C., Glant, T.T., Roughley, P.J., and Poole, A.R. (1987) The identification and characterization of two populations of aggregating proteoglycans of high buoyant density isolated from post-natal human cartilages of different ages. *Biochem J.* 248, 735-740.
- Wight, T.N., Heinegard, D.K., and Hascall, V.C. (1991) Proteoglycans: structure and function. *Cell biology of extracellular matrix.* Plenum Press, New York.
- Wisniewski, H.G., Hua, J.C., Poppers, D.M., Naime, D., Vilcek, J., and Cronstein BN. (1996) TNF/IL-1-inducible protein TSG-6 potentiates plasmin inhibition by inter-alpha-inhibitor and exerts a strong anti-inflammatory effect *in vivo.* *J Immunol.* 156, 1609-1615.
- Wisniewski, H.G., Maier, R., Lotz, M., Lee, S., Klampfer, L., Lee, T.H., and Vilcek, J. (1993) TSG-6: a TNF-, IL-1-, and LPS-inducible secreted glycoprotein associated with arthritis. *J Immunol.* 151, 6593-6601.
- Witter, J., Roughley, P.J., Webber, C., Roberts, N., Keystone, E., and Poole, A.R. (1987) The immunologic detection and characterization of cartilage proteoglycan degradation products in synovial fluids of patients with arthritis. *Arthritis Rheum.* 30, 519-529.
- Xu, T., Bianco, P., and Fisher, L.W. (1998) Targeted disruption of the biglycan gene leads to an osteoporosis-like phenotype in mice. *Nat Genet.* 20, 78-82.
- Yingsang, W. (1996) The development of a simple method for hyaluronan determination using biotinylation technique. M.S. Thesis, Chiangmai University.
- Zhang, Y., Guerassimov, A., Leroux, J.Y., Cartman, A., Webber, C., Lalic, R., de Miguel, E., Rosenberg, L.C., and Poole, A.R. (1998) Arthritis induced by proteoglycan aggrecan G1 domain in BALB/c mice. Evidence for t cell involvement and the

- immunosuppressive influence of keratan sulfate on recognition of t and b cell epitopes. *J Clin Invest.* 101, 1678-1686.
- Zheng, J., Luo, W., and Tanzer, M.L., (1998) Aggrecan synthesis and secretion. A paradigm for molecular and cellular coordination of multiglobular protein folding and intracellular trafficking. *J Biol Chem.* 273, 12999-13006.