

## REFERENCES

- Bandaru S. R., Chinthalapally V. R., Gary K. (1993). Chemoprevention of colon carcinogenesis by organosulfur compounds. *Cancer Res*, 53, 3493-3498
- Batist G., Tulpule A., Sinha B. K. (1986). Overexpression of a novel anionic glutathione transferase in multidrug resistant human breast cancer cells. *J Biol Chem*, 261, 15544-15549.
- Bell D. A., Taylor J. A., Paulson D. F. (1993). Genetic risk and carcinogen exposure: a common inherited defect of the carcinogen-metabolism gene glutathione S-transferase M1 (GSTM1) that increases susceptibility to bladder cancer. *J Natl Cancer Inst*, 85, 1159-64.
- Bird R. A. (1987). Observation and quantification of aberrant crypts in the murine colon treated with a colon carcinogenesis: preliminary findings. *Cancer Lett*, 37, 147-151.
- Boech E. M. A., Humboldt G. (1981). Cardio-circulatory effects of total water extract in normal persons and of stevioside in rats. *Ciencia e cultura*, 32, 208-210.
- Bonvie L., Bonvie B., Gates D. (1997). The stevia story, a tale of incredible sweetness and intrigue. B.E.D. Publications, Atlanta.
- Bradford M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem*, 72, 248-254.
- Bridel M., Lavielle R. (1931a). Le principe saveur sucre du Ka-h- (*Stevia rebaudiana*) Bertoni. *Bull Soc Chim Biol*, 13, 636-655.
- Bunnell E., Pacht E. R. (1993). Oxidized glutathione is increased in alveolar fluid of patients with adult respiratory distress syndrome. *Am Rev Resp Dis*, 148, 1174-1178.

- Carmmer B., Ikan R. (1986). Sweet glycosides from the stevia plant. *Chem Br*, 30, 915-917.
- Carr B. I. (1985). Chemical carcinogens and inhibitors of carcinogenesis in the human diet. *Cancer*, 55, 218-224.
- Chang S. S., Cook J. M. (1983). Stability studies of stevioside and rebaudioside A in carbonated beverages. *J Agric Food Chem*, 31, 409-412.
- Chang W. W. L. (1984). Histogenesis of colon cancer in experimental animals. *Scandinavian Journal of Gastroenterology*, 19 (suppl 104), 27-44.
- Chappell J. (1995). Biochemistry and molecular biology of the isoprenoid biosynthetic pathway in plants. *Annu Rev Plant Physiol Plant Mol Biol*, 46, 521-547.
- Chili E., Malvadi G., (1985). Role of Cyt P-450 dependent and FAA containing mono oxygenases in the bioactivation of thioacetamide, thiobezamide and their sulfoxides. *Biochemical Pharmacology*, 34, 395-396.
- Clairmont A., Ebert T., Weber H., Zoidl C., Eickelmann P., Schulz W. A., Sies H., Ryffel G. U. (1994). Lowered amounts of the tissue-specific transcription factor LFB1 (HNF1) correlate with decreased levels of glutathione S-transferase a mRNA in human renal cell carcinoma. *Cancer Research*, 54, 319-323.
- Coles B., Ketterer B. (1990). The role of glutathione and glutathione-S-transferase in chemical carcinogenesis. *Crit Rev Biochem Mol Biol*, 25, 47-70.
- Crammer B., Ikan R. (1986). Sweet glycosides from the stevia plant. *Chem Brit*, 22, 915-916.
- Cross C. E., Halliwell B., Borish E. T. (1987). Oxygen radicals and human disease (proceedings of a conference). *Ann Intern Med*, 107, 526-545.
- Daniel B. Mowry (1992). Life With Stevia: How Sweet It Is. Nutritional and Medicinal Uses, Reprinted of the publisher. Copyright 1992
- Das S., Das A. K., Murphy R. A., Punwani I. C., Nasution M. D., Kinghorn A. D. (1992). Evaluation of the cariogenic potential of the intense natural sweeteners stevioside and rebaudioside A. *Caries Res*, 26, 363-366.

- Dillio C., Del Boccio G., Aceto A., Cassacia R., Mucilli F., Fedirici G. (1985). Elevation of glutathione-S-transferase activity in human lung tumor. *Carcinogenesis*, 9, 335-340.
- Doll R., Peto R. (1981). The causes of cancer: quantitative assessment of avoidable risk of cancer in the United State today. *J Natl Cancer Inst*, 66, 1191-1308.
- Drotman R. B., Lawhorn G. T. (1978). Serum enzymes are indicators of chemical induced liver damage. *Drug and chemical Toxicology*, 1, 163-171.
- Duke R. C., Ojcius D. M., Young J. D. (1996). Cell suicide in health and disease. *Sci Am*, 275(6), 80-87.
- Eickelmann P., Morel F., Schulz W. A., Sies H. (1995). Turnover of Glutathione S-Transferase a mRNAs Accelerated by 12-0-Tetradecanoyl Phorbol-13-Acetate (TPA) in Human Hepatoma and Colon Carcinoma Cell Lines. *Eur J Biochem*, 229, 21-26.
- Esther M. M., Van Lieshout, Gary H. Posner, Benjamin T. Woodard, Wilbert H. M. Peters (1998). Effect of the sulforaphane analog compound 30, indole-3-carbinol, D-limonene or relafen on glutathione-S-transferase and glutathione peroxidase of the rat digestive tract. *Biochimica et Bioshysica Acta*, 1379, 325-336.
- Feinberg A., Zedeck M. S. (1980). Production of a highly reactive alkylating agent from the organospecific carcinogen methylazoxymethanol by alcohol dehydrogenase. *Cancer Res*, 40, 4446-4450.
- Fiala E. S. (1977). Investigations into metabolism and mechanism of action of the colon carcinogens, 1,2-dimethylhydrazine and azoxymethane. *Cancer*, 40, 2436-2445.
- Fitzhugh, O. G., Nelson, A. A. (1948). Liver tumours in rats fed thiourea or thioacetamide. *Science*, 108, 626-628.
- Friling R. S., Bensimon A., Tichauer Y., Daniel V. (1990). Xenobiotic-inducible expression of murine glutathione S-transferase Ya subunit gene is controlled by an electrophile-responsive element. *Proc Natl Acad Sci USA*, 87, 6258-6262.
-

- Kinghorn A. D., Soejarto D. D. (1991). Stevioside. In: L. O. Nabors and R. C. Gelardi (Eds.), *Alternative Sweeteners*. Marcel Dekker, Inc. New York, Basel, Hong Kong, 157.
- Fujita H., Edahiro T. (1979). Safety and utilization of stevia sweetener. *Shokuhin Kogyo*, 22 (20), 66-72.
- Geoffrey J. B., John D. H. (1993) Glutathione-S-transferase: biomedical applications. *Advance Clin Chem*, 30,281-279.
- Greenward P., Cullen J. W. (1985). The new emphasis in cancer control. *J Natl Cancer Inst*, 74, 543-551.
- Habig W. H. (1983). Glutathione-S-transferase: Versatile enzymes of detoxification. In "Radioprotectors and Anticarcinogens" (Nygaard O. F., Simic M. G., eds.). 169-190, New York, Academic Press.
- Haider Raza, Ljaz Ahmed, Mohammad S. Lakhani, Ashutosh K. Sharma, David Pallot, William Montague (1996). Effect of bitter melon (*Momordica Charantia*) Fruit Juice on the hepatic cytochrome P450-dependent monooxygenases and glutathione-S-transferase in Streptozotocin-induced diabetic rats. *Biochemical Pharmacology*, 52, 1639-1642.
- Hanson J. R., De Oliveira B. H. (1993). Stevioside and related sweet diterpenoid glycosides. *Nat Prod Rep*, 10, 301-309.
- Hatono S., Jimenez A., Wargovich M. J. (1996). Chemopreventive effect of S-allylcysteine and its relationship to the detoxification enzyme glutathione-S-transferase. *Carcinogenesis*, 17, 1041-1044.
- Hwang D. F., Hour J. L., Cheng H. M. (2000). Effect of taurine on toxicity of oxidized fish oil in rats. *Food Chem Toxicol*, 38, 585-591
- Jacobs, David S. (1996). *Laboratory Test Handbook*. 4th ed. Cleveland, OH: Lexi-Comp Inc.
- Katayama O., Sumida T., Hayashi H., Mitsuhashi H. (1976). The practical application of Stevia and research and development data (English translation). I.S.U. Company, Japan. 747 pp.

- Kenneth S. L. (2000). Hepatic Clinical Chemistry, College of Veterinary medicine, The university of Georgia.
- Kinghorn A. D., Soejarto D. D. (1985). Current status of stevioside as a sweetening agent for human use. Pages 1-52 in H. Wagner, H. Hikino and N. R. Farnsworth, eds. Economic and medicinal plant research. Academic Press, London.
- Kitahara A., Satoh K., Nishimura K., Ishikawa T., Ruike K., Sato K., Tsuda H., Ito N. (1984). Changes in molecular forms of rat hepatic glutathione S-transferase during chemical hepatocarcinogenesis. *Cancer Res*, 44(6), 2698-2703.
- Kodate C., Fukushi A., Narita T., Kudo H., Soma Y., Sato K. (1986). Human placental form of glutathione S-transferase (GST-pi) as a new immunohistochemical marker for human colonic carcinoma. *Jpn J Cancer Res*, 77(3), 226-229.
- Kosower N. S., Kosower E. M. (1978). The glutathione status of cells. *Int Rev Cytol*, 54, 109-160.
- Laemmli U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, 227, 680-685.
- Lafuente A., Pujol F., Carretero P., Villa J. P., Cuchi A. (1993). glutathione-S-transferase mu deficiency as a marker for the susceptibility to bladder and larynx cancer among smokers. *Cancer Lett*, 68, 49-54.
- Lomaestro B. M., Malone M. (1995). Glutathione in health and disease: pharmacotherapeutic issues. *Ann Pharmacother*, 29(12), 1263-1273.
- Mannervik B., Alin P., Guthenberg C., Jenson H., Tahir M. K., Warholm M., Jornvall H. (1985). Identification of three classes of cytosolic glutathione transferase common to several mammalian species: correlation between structural data and enzymatic properties. *Proc Natl Acad USA*, 82, 7202-7206.
- Massad L., de Waziers I., Ribrag V., Janot F., Beaune P. H., Morizet J., Gouyette A., Chabot G. G. (1992). phase I and phase II drug-metabolizing enzyme systems. *Cancer Res*, 52(23), 6567-6575.
-

- Matsui M., Matsui K., Kawasaki Y., Oda Y., Noguchi T., Kitagawa Y., Sawada M., Hayashi M. (1996). Evaluation of the genotoxicity of stevioside and steviol using six in vitro and one in vivo mutagenicity assays. *Mutagenesis*, 11(6), 573-579.
- McDonald P. J. (1989). The functions and the regulation of normal gut bacteria, and bacterial over growth syndromes. *Annu Rev Microbiol*, 43, 289-305.
- McGarvey D. J., Croteau R. (1995). Terpenoid metabolism. *Plant Cell*, 7, 1015-1026.
- McLellan E., Bird R. P. (1991). Effect of disulfiram on 1,2-dimethylhydrazine-and azoxymethane-induced aberrant crypt foci. *Carcinogenesis*, 12, 969-972.
- Moscow J. A., Fairchild C. R., Madden M. J., Ransom D. T., Wieand H. S., Poplack D. G., Cossman J. (1989). Expression of anionic glutathione-S-transferase and P-glycoprotein genes in human tissues and tumors. *Cancer Res*, 48, 1422-1428.
- Moscow J. A., Townsend A. J., Goldsmith M. E., Whang-Peng J., Vickers P. J., Poisson R., Cowan K. H. (1988). Isolation of anionic glutathione-S-transferase gene and the relation of its expression to estrogen receptor content in primary breast cancer. *Proc Natl Acad Sci USA*, 85, 6518-6522.
- Nakava K., Saijo N., Tsuchida S. (1990). Glutathione-S-transferase pi as a determinant of drug resistance in transfectant cell lines. *J Biol Chem*, 265, 4296-4301.
- Nalpas B., Vassault A., Charpin S., Lacour B., Berthelot P. (1988). Serum mitochondrial aspartate aminotransferase as a marker of chronic alcoholism: diagnostic value and interpretation in a liver unit. *Hepatology*, 6(4), 608-614.
- Nijhoff W. A., Peters W. H. (1992). Induction of rat hepatic and intestinal glutathione-S-transferase by dietary butylated hydroxyanisole. *Biochem Pharmacol*, 44(3), 596-600.
- Oveido C. A., Franciani G., Moreno R., Maas L. C. (1970). Accion hipoglyceminante de la *Stevia rebaudiana* Bertoni (Kaa-he-e). *Excerpta Medica*, 209, 92.
- Pacht E. R., Timerman A. P., Lykens M. G. (1991). Deficiency of alveolar fluid glutathione in patients with sepsis and the adult respiratory distress syndrome. *Chest*, 100, 1397-1403.

- Pagana, Kathleen Deska. (1998). *Mosby's Manual of Diagnostic and Laboratory Tests*. St. Louis, MO: Mosby, Inc.
- Pendergast W. R. (1991). GRAS Affirmation Petition Number 2G0390.
- Pereira M. A., Barnes L. H., Rassman V. L., Kelloff G. V., Steele V. E. (1994). Use of azoxymethane-induced foci of aberrant crypts in rat colon to identify potential cancer chemopreventive agents. *Carcinogenesis*, 15, 1049-1054.
- Pezzuto J. M., Compadre C. M., Swanson S. M., Nanayakkara D., Kinghorn A. D. (1985). Metabolically activated steviol, the aglycone of stevioside, is mutagenic. *Proc Natl Acad Sci USA*, 82 (8), 2478-2482.
- Phillips K. C. (1989). Stevia: steps in developing a new sweetener. Pages 1-43 in T.H. Grenby ed. *Developments in sweeteners*, Volume 3. Elsevier Applied Science, London.
- Pickett C. B., Lu A. Y. (1989). Glutathione-S-transferase: gene structure, regulation and biological function. *Annu Rev Biochem*, 58, 743-764.
- Planas G. M., Kuc J. (1968). Contraceptive properties of *Stevia rebaudiana*. *Science*, 162, 1007.
- Planas G. M., Kue J. (1968). Contraceptive properties of *Stevia rebaudiana*. *Science*, 162, 1007.
- Pretlow T. P., Barrow B. J., Ashton W. S., O'Riordan M. A., Pretlow T. G., Jurcisek J. A., Stellato T. A. (1991). Aberrant crypts: putative preneoplastic foci in human colonic mucosa. *Cancer Res*, 51, 1564-1567.
- Procinska E., Bridges B. A., Hanson J. R. (1991). Interpretation of results with the 8-azaguanine resistance system in *Salmonella typhimurium*: no evidence for direct acting mutagenesis by 15-oxosteviol, a possible metabolite of steviol. *Mutagenesis*, 6, 165-167.
- Ranjana P. Bird, Elizabeth A. Mclellan, W. Robert Bruce (1989). *Cancer Surveys*, 8 (1), 189-200.

- Raza H., Qureshi M. M., Montague W. (1995). Alteration of glutathione, glutathione-S-transferase and lipid peroxidation in mouse skin and extracutaneous tissues after topical application of gasoline. *Int J Biochem cell Biol*, 27 (3), 271-277.
- Rebbeck, T.R. (1997). Molecular epidemiology of the human glutathione S-transferase genotypes GSTM1 and GSTT1 in cancer susceptibility. *Cancer Epidemiology, Biomarkers & Prevention*, 6, 733-43.
- Reitman S., Franxel S. A. (1957). Colorimetric method for the Determination of serum Glutamic-oxaloacetic acid and Glutamic-pyruvic acid transaminase. *Am J Clin Pathol*, 28, 56-63.
- Robinson B.L. (1930). Contributions from the Gray Herbarium of Harvard University. The Gray Herbarium of Harvard University, Cambridge.
- Seidegard J., Pero R. W., Markowitz M. M., Roush G., Miller D. G. (1990). Isoenzyme of glutathione-S-transferase (class mu) as a marker for the susceptibility to lung cancer: a follow up study. *Carcinogenesis*, 11, 33-36.
- Sherwood P., Lyburn I., Brown S., Ryder S. (2001). How are abnormal results for liver function tests dealt with in primary care? *Audit of yield and impact British Medical Journal*, 322, 276-278.
- Sincholle D., Marcorelles P. (1989). Study of the anti-androgenic activity of extracts of *Stevia rebaudiana* Bertoni (English abstr.). *Med Plants and Phytotherapy*, 23, 282-287.
- Slater A. F., Stefan C., Nobel I., Van den Dobbelsteen D. J., Orrenius S. (1995). Signalling mechanisms and oxidative stress in apoptosis. *Toxicol Lett*, 82, 149-153.
- Smith M. T., Evans C. G., Doane-Setzer P. (1989). Denitrosation of 1,3-bis(chloroethyl)-1-nitrosourea by class mu glutathione transferase and its role in cellular resistance in rat brain tumor cells. *Cancer Res*, 49, 2621-2625.
- Smoliar V. I., Karpilovskaia E. D., Sali N. S., Tsapko E. V., Lavrushenko L. F., Gulich M. P., Kryshevich L. P., Grigorenko S. N. (1992). Effect of a new sweetening agent from *Stevia rebaudiana* on animals. *Vopr Pitan*, 1, 60-63.



- Soejarto D. D., Compadre C. M., Medon P. J., Kamath S. K., Kinghorn A. D. (1983). Potential sweetening agents of plant origin. II. Field search for sweet-tasting Stevia species. *Econ Bot*, 37, 71-79.
- Soejarto D.D., Compadre C. M., Medon P. J., Kamath S. K., Kinghorn A. D. (1983). Potential sweetening agents of plant origin, II. Field search for sweet-tasting Stevia species, *Econ Bot*, 37, 74-82.
- Sohn S. O., Ishizaki H., Yang S. C., Fiala E. (1991). Metabolism of azoxymethane, methylazoxymethanol and N-nitrosodimethylamine by cytochrome P450IIE1, *Carcinogenesis*, 12(1), 127-131.
- Suttajit M., Vinitketkaumnue U., Meevatee U., Buddhasukh D. (1993). Mutagenicity and human chromosomal effect of stevioside, a sweetener from Stevia rebaudiana Bertoni. *Environ health perspect*, 101(3), 53-56.
- Theresa P. P., Gregory A. S., Saeid B. A., Thomas G. P. (1992). Aberrant crypts correlate with tumor incidence in F344 rats treated with azoxymethane and phytate. *Carcinogenesis*, 3(9), 1509-1512.
- Theodorus P. M., Akerboom T. P., Sies H. (1981). Assay of glutathione, glutathione disulfide, and glutathione mixed disulfides in biological samples. *Methods Enzymol*, 77, 373-382.
- Toyada K., Matsui H., Shoda T., Uneyama C., Takada K. (1997). Assessment of the carcinogenicity of stevioside in F344 rats. *Food Chem Toxicol*, 35 (6), 597-603.
- Toyoda K., Matsui H., Shoda T., Uneyama C., Takada K., Takahashi M. (1997). Assessment of the carcinogenicity of stevioside in F344 rats. *Food Chem Toxicol*, 35(6), 597-603.
- Ward J. M., Yamamoto R. S., Brown C. A. (1973). Pathology of intestinal neoplasms and other lesions in rats exposed to azoxymethane. *J Natl Cancer Inst*, 51, 1029-1039.
- Wareing C. J., Balck S. M., Hayes J. D., Wolf C. R. (1993). Increased level of alpha class and pi class glutathione-S-transferases in cell lines resistant to 1-chloro-2,4-dinitrobenzene. *Eur J Biochem*, 217, 617-676.

- Wareing C. J., Balck S. M., Hayes J. D., Wolf C. R. (1993). Increased level of alpha class and pi class glutathione-S-transferases in cell lines resistant to 1-chloro-2,4-dinitrobenzene. *Eur J Biochem*, 217, 617-676.
- Wargovich M. J., Jimenez A., Mckee K., Steele V. E., Velasco M., Woods J., Price R., Gray K., Kelloff G. J. (2000). Efficacy of potential chemopreventive agents on rat colon aberrant crypt formation and progression. *Carcinogenesis*, 21, 1149-1155.
- Wasuntarawat C., Temcharoen P., Toskulkao C., Mungkormkarn P., Suttajit M., Glinsukon T. (1998). Developmental toxicity of steviol, the aglycone of stevioside, is mutagenic. *Drug Chem Toxicol*, 82(8), 2478-2482.
- Weisburger J. H. (1971). Colon carcinogen: Their metabolism and mode of action. *Cancer*, 28, 60-70.
- Weisburger J. H., Reddy B. S., Wynder E. L. (1997). Colon cancer: Its epidemiology and experimental product. *Cancer*, 40, 2414-2420.
- Wingard W. L., Brown J. P., Enderlin F. N. (1980). Intestinal degradation and absorption of the glycosidic sweetener stevioside and rebaudioside A. *Experientia*, 84, 510.
- Xili L., Chengjiayn B., Eryi X., Reiming S., Yuengming W., Haodong S., Zhiyian H. (1992). Chronic oral toxicity study of stevioside in rats. *Food Chem Toxicol*, 30 (11), 965-975.
- Yodyingyuad V., Bunyawong S. (1991). Effect of stevioside on growth and reproduction. *Hum Reprod*, 6(1), 158-165.
- Yodyingyuad V., Bunyawong S. (1991). Effects of stevioside on growth and reproduction. *Human Reprod*, 6, 158-165.
- Zhong S., Wyllie A. H., Barnes D. (1993). Relationship between the GSTM1 genetic polymorphism and susceptibility to bladder, breast and colon cancer. *Carcinogenesis*, 14, 1821-1814.