

TABLE OF CONTENTS

	page
ACKNOWLEDGEMENTS	iii
ABSTRACT	iv
TABLE OF CONTENTS	viii
LIST OF FIGURES	xii
LIST OF TABLES	xvi
ABBREVIATIONS	xvi
CHAPTER 1 INTRODUCTION	
1.1 Statement of problems	1
1.2 <i>Stevia rebaudiana</i> Bertoni	4
1.2.1 The chemistry of the diterpene glycoside sweeteners	4
1.2.2 Biosynthesis of the sweet glycosides	6
1.2.3 Safety of Stevia sweeteners	8
1.3 Aberrant Crypt Foci Formation	9
1.4 Serum Aminotransferase Enzyme	11
1.4.1 Aspartate aminotransferase (AST)	11
1.4.2 Alanine aminotransferase (ALT)	12
1.5 Glutathione	12
1.6 Glutathione-S-transferase	14
1.6.1 Glutathione-S-transferase Isoenzymes	15
1.6.2 GST as a tumor marker and susceptibility to toxic chemical	16
1.7 Objectives of the study	19
CHAPTER 2 MATERIALS AND METHODS	
2.1 Chemicals and instruments	20

2.2 Preparation of Stevia extract mixed in Matoom juice	20
2.3 Animals	20
2.4 Effects of Matoom juice containing Stevia extract as sweetener on aberrant crypts foci (ACF) formation	21
2.5 Aberrant crypt foci (ACF) assay	24
2.6 Determination of serum transaminase activity	25
2.7 Preparation of cytosolic fraction	28
2.8 Reduced glutathione assay	30
2.9 Glutathione-S-transferase activity assay	32
2.10 Characterization of glutathione-S-transferase subunits	34
2.10.1 SDS gel electrophoresis	34
2.10.2 Protein Western blot analysis	34
2.11 Statistic Evaluation	34
CHAPTER 3 RESULTS	
3.1 Effects of Matoom juice containing Stevia extract as sweetener on the number of aberrant crypt foci	36
3.1.1 Effects of Matoom juice containing Stevia extract as sweetener on the number of aberrant crypt foci at initiation stage	36
3.1.2 Effects of Matoom juice containing Stevia extract as sweetener on the number of aberrant crypt foci at promotion stage	38
3.2 Effects of 35 or 100 days administration of Matoom juice containing Stevia extract as sweetener on rat body weight	40
3.3 Effects of 35 or 100 days administration of Matoom juice containing Stevia extract as sweetener on AST and ALT activities in rats serum	42
3.3.1 Effects of 35 days administration of Matoom juice containing Stevia extract as sweetener on AST and ALT activities in rats serum	42
3.3.2 Effects of 100 days administration of Matoom juice containing Stevia extract as sweetener on AST and ALT activities in rats serum	42

3.4 Effects of 35 or 100 days administration of Matoom juice containing Stevia extract as sweetener on reduced glutathione (GSH) levels in rat liver and intestinal mucosa	45
3.4.1 Effect of 35 or 100 days administration of Matoom juice containing Stevia extract as sweetener on reduced glutathione levels in rat liver	45
3.4.2 Effects of 35 or 100 days administration of Matoom juice containing Stevia extract as sweetener on reduced glutathione levels in rat intestinal mucosa	45
3.5 Effects of 35 or 100 days administration of Matoom juice containing Stevia extract as sweetener on glutathione-S-transferase (GST) activity in rat liver and intestinal	48
3.5.1 Effects of 35 or 100 days administration of Matoom juice containing Stevia extract as sweetener on glutathione-S-transferase activity in rat liver	48
3.5.2 Effects of 35 or 100 days administration of Matoom juice containing Stevia extract as sweetener on glutathione-S-transferase activity in rat intestinal mucosa	48
3.6 Effects of 35 or 100 days administration of Matoom juice containing Stevia extract as sweetener on glutathione-S-transferase subunit level in rat liver and intestinal mucosa	51
3.6.1 Effects of 35 or 100 days administration of Matoom juice containing Stevia extract as sweetener on glutathione-S-transferase subunit level in rat liver	51
3.6.2 Effects of 35 or 100 days administration of Matoom juice containing Stevia extract as sweetener on glutathione-S-transferase subunit level in rat intestinal mucosa	60

CHAPTER 4 DISCUSSION AND CONCLUSION	63
REFERENCES	70
APPENDIXES	80
CURRICULUM VITAE	88

มหาวิทยาลัยเชียงใหม่
Chiang Mai University

LIST OF FIGURES

Figures	page
1. <i>Stevia rebaudiana</i> Bertoni	3
2. Chemical structure of the steviol-glycosides	5
3. Biosynthetic pathway for synthesis of steviol glycosides	7
4. Morphological of ACF	10
5. Structure of glutathione (GSH, γ -L-glutamyl-L-cysteinylglycine)	13
6. The role of glutathione-dependent pathways in cell survival	14
7. The three-dimensional structure of two subunits in the dimer of GST and GSH-binding site.	16
8. Experiment schedule to study of Matom juice containing Stevia extract as sweetener on ACF formation in initiation stage	22
9. Experiment schedule to study of Matom juice containing Stevia extract as sweetener on ACF formation in promotion stage	23
10. Protocol of aspartate aminotransferase (AST) activity determination	26
11. Protocol of alanine aminotransferase (ALT) activity determination	27
12. Scheme of cytosolic fraction preparation	29
13. Scheme of cytosolic protein determination	29
14. Protocol of sample preparation to determine reduced glutathione	31
15. Scheme of reduced glutathione assay	31
16. Scheme of glutathione-S-transferase activity assay	33
17. Protocol of glutathione-S-transferase subunits determination	35
18. Effects of 35 days administration of Matom juice containing Stevia extract as sweetener on rat body weight	40
19. Effects of 100 days administration of Matom juice containing Stevia extract as sweetener on rat body weight	41

20. Coomessie blue stain of rats cytosolic GSTs on running SDS-PAGE	53
21. Western blot analysis using antibodies raised against GST Yc subunit of rats hepatic cytosol	54
22. Western blot analysis using antibodies raised against GST Ya subunit of rats hepatic cytosol	55
23. Western blot analysis using antibodies raised against GST Yb subunit of rats hepatic cytosol	56
24. Western blot analysis using antibodies raised against GST YT subunit of rats hepatic cytosol	57
25. Matoom (<i>Aegle marmelos</i> Corr.), Family: Rutaceae	85

LIST OF TABLES

Tables	page
1. Number of ACF in rats after received Matoom juice containing Stevia extract as sweetener at initiation stage	37
2. Number of ACF in rats after received Matoom juice containing Stevia extract as sweetener at promotion stage	39
3. Effect of 35 days administration of Matoom juice containing Stevia extract as sweetener on AST and ALT activities in rat serum	43
4. Effect of 100 days administration of Matoom juice containing Stevia extract as sweetener on AST and ALT activities in rat serum	44
5. Effect of 35 or 100 days administration of Matoom juice containing Stevia extract as sweetener on reduced glutathione levels in rat liver	46
6. Effect of 35 or 100 days administration of Matoom juice containing Stevia extract as sweetener on reduced glutathione levels in rat intestinal mucosa	47
7. Effect of 35 or 100 days administration of Matoom juice containing Stevia extract as sweetener on glutathione-S-transferase activity in rat liver	49
8. Effect of 35 or 100 days administration of Matoom juice containing Stevia extract as sweetener on glutathione-S-transferase activity in rat intestinal mucosa	50
9. Effect of 35 days administration of Matoom juice containing Stevia extract as sweetener on glutathione-S-transferase subunit level in rat liver	58
10. Effect of 100 days administration of Matoom juice containing Stevia extract as sweetener on glutathione-S-transferase subunit level in rat liver	59

- | | |
|--|----|
| 11. Effect of 35 days administration of Matoom juice containing Stevia extract as sweetener on glutathione-S-transferase subunit level in rat intestinal mucosa | 61 |
| 12. Effect of 100 days administration of Matoom juice containing Stevia extract as sweetener on glutathione-S-transferase subunit level in rat intestinal mucosa | 62 |

มหาวิทยาลัยเชียงใหม่
Chiang Mai University

ABBREVIATIONS

Abs	absorbance
ACF	aberrant crypt foci
AOM	azoxymethane
APS	ammonium persulfate
BSA	bovine serum albumin
°C	degree celsius
CDNB	1-chloro-2,4-dinitrobenzene
DMSO	dimethyl sulfoxide
g	gram
GSH	reduced glutathione
GST	glutathione-S-transferase
HCl	hydrochloric acid
H ₂ O	water
H ₂ O ₂	hydrogen peroxide
hr	hour
KCl	potassium chloride
kg	kilogram
kg.bw.	kilogram body weight
M	molarity
mM	millimolar
mg	milligram
min	minute
ml	millilitre
MOPS	3-[N-morpholino] propanesulfuric acid
NaOH	sodium hydroxide
nm	nanometer

No.	number
OD	optical density
PAGE	polyacrylamide gel electrophoresis
PBS	phosphate buffer saline
rpm	round per minute
RT	room temperature
s.c	subcutaneous
sec	second
TEMED	N,N,N',N'-tertramethylethylene diamine
U	unit
w/v	weight by volume
μ g	microgram
μ l	microlitre
%	percent