

## References

- Abdel-Rahman SM, O'Reilly KL, Malone JB. Evaluation of a diagnostic monoclonal antibody-based capture enzyme-linked immunosorbent assay for detection of a 26- to 28-kd *Fasciola hepatica* coproantigen in cattle. Am J Vet Res 1998; 59:533-7.
- Agi PI. Comparative helminth infections of man in two rural communities of the Niger Delta, Nigeria. West Afr J Med 1997; 16:232-6.
- Agi PI. Pattern of infection of intestinal parasites in Sagbama community of the Niger Delta, Nigeria. West Afr J Med 1995; 14:39-42.
- Allan JC, Mencos F, Garcia-Noval J, Sarti E, Flisser A, Wang Y, et al. Dipstick dot ELISA for the detection of *Taenia* coproantigens in humans. Parasitology 1993; 107 (Pt 1):79-85.
- Allan JC, Craig PS, Garcia Noval J, Mencos F, Liu D, Wang Y, et al. Coproantigen detection for immunodiagnosis of echinococcosis and taeniasis in dogs and humans. Parasitology 1992; 104:347-56.
- Anantaphruti MT, Nuamtanong S, Muennoo C, Sanguankiat S, Pubampen S. *Strongyloides stercoralis* infection and chronological changes of other soil-transmitted helminthiases in an endemic area of southern Thailand. Southeast Asian J Trop Med Public Health 2000; 31:378-82.

- Arakaki T, Iwanaga M, Kinjo F, Saito A, Asato R, Ikeshiro T. Efficacy of agar-plate culture in detection of *Strongyloides stercoralis* infection. J Parasitol 1990; 76:425-8.
- Bailey JW. A serological test for the diagnosis of *Strongyloides* antibodies in ex Far East Prisoners of War. Ann Trop Med Parasitol 1989; 83:241-7.
- Bangs MJ, Purnomo, Andersen EM, Anthony RL. Intestinal parasites of humans in a highland community of Irian Jaya, Indonesia. Ann Trop Med Parasitol 1996; 90:49-53.
- Bannon JP, Fater M, Solit R. Intestinal ileus secondary to *Strongyloides stercoralis* infection: case report and review of the literature. Am Surg 1995; 61:377-80.
- Beaver PC, Jung RC, Cupp EW. Examination of specimens for parasites. In: Clinical Parasitology. 9th ed. Philadelphia: Lea & Febiger; 1984. p. 733-58.
- Bogish BJ, Cheng TC. Intestinal nematodes. In: Human parasitology. 2nd ed. San Diego: Academic Press; 1998. p. 349-56.
- Brown HW, Neva FA. Intestinal nematodes of human beings. In: Basic clinical parasitology. 5th ed. New Jersey: Prentice/Hall International Inc.; 1983. p 115-9.
- Carp NZ, Nejman JH, Kelly JJ. Strongyloidiasis. An unusual cause of colonic pseudopolyposis and gastrointestinal bleeding. Surg Endosc 1987; 1:175-7.
- Carroll SM, Karthigasu KT, Grove DI. Serodiagnosis of human strongyloidiasis by an enzyme-linked immunosorbent assay. Trans R Soc Trop Med Hyg 1981; 75:706-9.

- Chai JY, Hongvanthong B. A small-scale survey of intestinal helminthic infections among the residents near Pakse, Laos. Korean J Parasitol 1998; 36:55-8.
- Chaicumpa W, Ybanez L, Kitikoon V, Pungpak S, Ruangkunaporn Y, Chongsanguan M, et al. Detection of *Opisthorchis viverrini* antigens in stools using specific monoclonal antibody. Int J Parasitol 1992; 22:527-31.
- Conway DJ, Atkins NS, Lillywhite JE, Bailey JW, Robinson RD, Lindo JF, et al. Immunodiagnosis of *Strongyloides stercoralis* infection: a method for increasing the specificity of the indirect ELISA. Trans R Soc Trop Med Hyg 1993a; 87:173-6.
- Conway DJ, Bailey JW, Lindo JF, Robinson RD, Bundy DA, Bianco AE. Serum IgG reactivity with 41-, 31-, and 28-kDa larval proteins of *Strongyloides stercoralis* in individuals with strongyloidiasis. J Infect Dis 1993b; 168:784-7.
- Conway DJ, Lindo JF, Robinson RD, Bundy DA, Bianco AE. *Strongyloides stercoralis*: characterization of immunodiagnostic larval antigens. Exp Parasitol 1994; 79:99-105.
- Craig PS, Gasser RB, Parada L, Cabrera P, Parietti S, Borgues C, et al. Diagnosis of canine echinococcosis: comparison of coproantigen and serum antibody tests with arecoline purgation in Uruguay. Vet Parasitol 1995; 56:293-301.
- de Kaminsky RG. Evaluation of three methods for laboratory diagnosis of *Strongyloides stercoralis* infection. J Parasitol 1993; 79:277-80.

- de Rezende CH, Costa-Cruz JM, Gennari-Cardoso ML. Enteroparasitoses in food handlers of the public schools in Uberlandia (Minas Gerais), Brazil. Rev Panam Salud Publica 1997; 2:392-7.
- Deplazes P, Gottstein B, Eckert J, Jenkins DJ, Ewald D, Jimenez-Palacios S. Detection of *Echinococcus* coproantigens by enzyme-linked immunosorbent assay in dogs, dingoes and foxes. Parasitol Res 1992; 78:303-8.
- Douce RW, Brown AE, Khamboonruang C, Walzer PD, Genta RM. Seroepidemiology of strongyloidiasis in a Thai village. Int J Parasitol 1987; 17:1343-8.
- Dreyer G, Fernandes-Silva E, Alves S, Rocha A, Albuquerque R, Addiss D. Patterns of detection of *Strongyloides stercoralis* in feces specimens: implications for diagnosis and clinical trials. J Clin Microbiol 1996; 34:2569-71.
- Espino AM, Diaz A, Perez A, Finlay CM. Dynamics of antigenemia and coproantigens during a human *Fasciola hepatica* outbreak. J Clin Microbiol 1998; 36:2723-6.
- Espino AM, Finlay CM. Sandwich enzyme-linked immunosorbent assay for detection of excretory secretory antigens in humans with fascioliasis. J Clin Microbiol 1994; 32:190-3.
- Gam AA, Neva FA, Krotoski WA. Comparative sensitivity and specificity of ELISA and IHA for serodiagnosis of strongyloidiasis with larval antigens. Am J Trop Med Hyg 1987; 37:157-61.

Gann PH, Neva FA, Gam AA. A randomized trial of single- and two-dose ivermectin versus thiabendazole for treatment of strongyloidiasis. J Infect Dis 1994; 169:1076-9.

Grove DI, Blair AJ. Diagnosis of human strongyloidiasis by immunofluorescence, using *Strongyloides ratti* and *S. stercoralis* larvae. Am J Trop Med Hyg 1981; 30:344-9.

Grove DI, Warren KS, Mahmoud AA. Algorithms in the diagnosis and management of exotic diseases. III. Strongyloidiasis. J Infect Dis 1975; 131:755-8.

Grove DI. Strongyloidiasis: A major roundworm infection of man. London: Taylor and Francis; 1989.

Grove DI. Treatment of strongyloidiasis with thiabendazole: an analysis of toxicity and effectiveness. Trans R Soc Trop Med Hyg 1982; 76:114-8.

Hanvanich M, Viscidi R, Laughon BE, Bartlett JG, Yolken RH. Stool desorbing activity: a possible cause of false-positive reactions in competitive enzyme immunoassays. J Clin Microbiol 1985; 21:184-8.

Jongwutiwes S, Charoenkorn M, Sitthichareonchai P, Akaraborvorn P, Putaporntip C. Increased sensitivity of routine laboratory detection of *Strongyloides stercoralis* and hookworm by agar-plate culture. Trans R Soc Trop Med Hyg 1999; 93:398-400.

Kasuya S, Khamboonruang C, Amano K, Murase T, Araki H, Kato Y, et al. Intestinal parasitic infections among schoolchildren in Chiang Mai, northern Thailand: an analysis of the present situation. J Trop Med Hyg 1989; 92:360-4.

- Kobayashi J, Hasegawa H, Forli AA, Nishimura NF, Yamanaka A, Shimabukuro T, *et al.* Prevalence of intestinal parasitic infection in five farms in Holambra, Sao Paulo, Brazil. Rev Inst Med Trop Sao Paulo 1995; 37:13-8.
- Kobayashi J, Hasegawa H, Soares EC, Toma H, Dacal AR, Brito MC, *et al.* Studies on prevalence of *Strongyloides* infection in Holambra and Maceio, Brazil, by the agar plate faecal culture method. Rev Inst Med Trop Sao Paulo 1996; 38:279-84.
- Koga K, Kasuya S, Khamboonruang C, Sukavat K, Nakamura Y, Tani S, *et al.* An evaluation of the agar plate method for the detection of *Strongyloides stercoralis* in northern Thailand. J Trop Med Hyg 1990; 93:183-8.
- Koga K, Kasuya S, Khamboonruang C, Sukhavat K, Ieda M, Takatsuka N, *et al.* A modified agar plate method for detection of *Strongyloides stercoralis*. Am J Trop Med Hyg 1991; 45:518-21.
- Kongtawelert P, Ghosh P. A new sandwich-ELISA method for the determination of keratan sulphate peptides in biological fluids employing a monoclonal antibody and labelled avidin biotin technique. Clin Chim Acta 1990; 195:17-26.
- Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. Nature 1970; 227:680-5.
- Lindo JF, Conway DJ, Atkins NS, Bianco AE, Robinson RD, Bundy DA. Prospective evaluation of enzyme-linked immunosorbent assay and immunoblot methods for the diagnosis of endemic *Strongyloides stercoralis* infection. Am J Trop Med Hyg 1994; 51:175-9.

Little MD . Comparative morphology of six species of *Strongyloides* (Nematoda) and redefinition of the genus. J Parasitol 1966; 52:69-84.

Machado ER, Costa-Cruz JM. *Strongyloides stercoralis* and other enteroparasites in children at Uberlandia city, state of Minas Gerais, Brazil. Mem Inst Oswaldo Cruz 1998; 93:161-4.

Mahmoud AA. Strongyloidiasis. Clin Infect Dis 1996; 23:949-52.

Malgor R, Nonaka N, Basmadjian I, Sakai H, Carambula B, Oku Y, et al. Coproantigen detection in dogs experimentally and naturally infected with *Echinococcus granulosus* by a monoclonal antibody-based enzyme-linked immunosorbent assay. Int J Parasitol 1997; 27:1605-12.

Miranda RA, Xavier FB, Menezes RC. Intestinal parasitism in a Parakana indigenous community in southwestern Para State, Brazil. Cad Saude Publica 1998; 14:507-11.

Mishell BB, Shiigi SM. Selected methods in cellular immunology. San Francisco: W.H. Freeman and company; 1980.

Moonsom S, Khunkeawla P, Kasinrerk W. Production of polyclonal and monoclonal antibodies against CD54 molecules by intrasplenic immunization of plasmid DNA encoding CD54 protein. Immunol Lett 2001; 76:25-30.

Morakote N, Khamboonruang C, Siriprasert V, Suphawitayanukul S, Marcanantachoti S, Thamasonthi W. The value of enzyme-linked immunosorbent assay (ELISA) for diagnosis of human trichinosis. Trop Med Parasitol 1991; 42:172-4.

Moustafa MA. An evaluation of the modified agar plate method for diagnosis of *Strongyloides stercoralis*. J Egypt Soc Parasitol 1997; 27:571-9.

Nageswaran C, Craig PS, Devaney E. Coproantigen detection in rats experimentally infected with *Strongyloides ratti*. Parasitology 1994; 108:335-42.

Neva FA, Gam AA, Burke J. Comparison of larval antigens in an enzyme-linked immunosorbent assay for strongyloidiasis in humans. J Infect Dis 1981; 144:427-32.

Neva FA. Biology and immunology of human strongyloidiasis. J Infect Dis 1986; 153:397-406.

Nonaka N, Iida M, Yagi K, Ito T, Ooi HK, Oku Y, et al. Time course of coproantigen excretion in *Echinococcus multilocularis* infections in foxes and an alternative definitive host, golden hamsters. Int J Parasitol 1996; 26:1271-8.

Ritchie LS, Lin S, Moon AP, Frick LP, Williams JE, Asakura S, et al. The possible effects of pH and specific gravity on the ether sedimentation procedure in concentrating eggs and cysts. Am J Trop Med Hyg 1960; 9:444-9.

Sato Y, Inoue F, Kiyuna S, Shiroma Y. Immunoblot analysis of three antigen preparations from *Strongyloides stercoralis* Larvae in human strongyloidiasis. Jpn J Parasitol 1990; 39:258-66.

Sato Y, Kobayashi J, Toma H, Shiroma Y. Efficacy of feces examination for detection of *Strongyloides* infection. Am J Trop Med Hyg 1995; 53:248-50.

Sato Y, Takara M, Otsuru M. Detection of antibodies in strongyloidiasis by enzyme-linked immunosorbent assay (ELISA). Trans R Soc Trop Med Hyg 1985; 79:51-5.

Siddiqui AA, Koenig NM, Sinensky M, Berk SL. *Strongyloides stercoralis*: identification of antigens in natural human infections from endemic areas of the United States. Parasitol Res 1997; 83:655-8.

Singh K, Vinayak VK, Bhasin DK, Ganguly NK. A monoclonal antibody-based test system for detection of *Entamoeba histolytica*-specific coproantigen. Indian J Gastroenterol 1999; 18:104-8.

Sirisinha S, Chawengkirttikul R, Haswell-Elkins MR, Elkins DB, Kaewkes S, Sithithaworn P. Evaluation of a monoclonal antibody-based enzyme linked immunosorbent assay for the diagnosis of *Opisthorchis viverrini* infection in an endemic area. Am J Trop Med Hyg 1995; 52:521-4.

Sirisinha S, Chawengkirttikul R, Sermswan R, Amornpant S, Mongkolsuk S, Panyim S. Detection of *Opisthorchis viverrini* by monoclonal antibody-based ELISA and DNA hybridization. Am J Trop Med Hyg 1991; 44:140-5.

Sukhavat K, Morakote N, Chaiwong P, Piangjai S. Comparative efficacy of four methods for the detection of *Strongyloides stercoralis* in human feces specimens. Ann Trop Med Parasitol 1994; 88:95-6.

Towbin H, Staehelin T, Gordon J. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. Proc Natl Acad Sci USA 1979; 76:4350-4.

Udonsi JK, Behnke JM, Gilbert FS. Analysis of the prevalence of infection and associations between human gastrointestinal nematodes among different age classes living in the urban and suburban communities of Port Harcourt, Nigeria. J Helminthol 1996; 70:75-84.

Uparanukraw P, Phongsri S, Morakote N. Fluctuations of larval excretion in *Strongyloides stercoralis* infection. Am J Trop Med Hyg 1999; 60:967-73.

Viscidi R, Laughon BE, Hanvanich M, Bartlett JG, Yolken RH. Improved enzyme immunoassays for the detection of antigens in fecal specimens. Investigation and correction of interfering factors. J Immunol Methods 1984; 67:129-43.

Voller A, Bidwell DE, Bartlett A. Enzyme immunoassays in diagnostic medicine. Theory and practice. Bull World Health Organ 1976; 53:55-65.

Widjana DP, Sutisna P. Prevalence of soil-transmitted helminth infections in the rural population of Bali, Indonesia. Southeast Asian J Trop Med Public Health 2000; 31:454-9.

Youssef FG, Mansour NS, Aziz AG. Early diagnosis of human fascioliasis by the detection of coproantigens using counterimmunoelectrophoresis. Trans R Soc Trop Med Hyg 1991; 85:383-4.

Zaha O, Hirata T, Kinjo F, Saito A. Strongyloidiasis--progress in diagnosis and treatment. Intern Med 2000; 39:695-700.

Zola H. Monoclonal antibodies: A manual of techniques. Boca Raton: CRC Press; 1987. p 23-62.

Zumla A, Croft SL. Chemotherapy and immunity in opportunistic parasitic infections in AIDS. *Parasitology* 1992; 105 Suppl:S93-101.

Zygmunt DJ. *Strongyloides stercoralis*. *Infect Control Hosp Epidemiol* 1990; 11:495-7.