

Table of Contents

	Page
Acknowledgements	iii
Abstract	iv
List of Tables	ix
List of Figures	x
Abbreviations	xi
Principle	1
Literature Reviews	5
1. Etiology	5
2. Clinical syndromes	6
3. Prevalence	8
4. Diagnosis	9
5. Therapy	15
6. Objectives of this study	16
Materials and Methods	17
1. Studied population	17
2. Examinations of fecal specimens	17
2.1 Formalin-ether concentration method	17
2.2 Agar plate culture method	17
3. Standard direct smear	18
4. Collection of L3	18
4.1 Agar plate culture method	18
4.2 Filter paper slant culture method	19
5. Collection of L1	19
6. Preparation of antigens	20
6.1 L1	20
6.2 L3 and adult worms	20
7. Partial purification of the 41-kDa protein of L3	20

	Page
8. Removal of SDS from the protein sample	21
9. Protein concentrations	21
10. Induction of antibodies against crude somatic antigens or 41-kDa protein of <i>S. stercoralis</i>	22
10.1 Animals	22
10.2 Immunization	22
-Mice	22
-Rabbit	22
11. Hybridoma production	23
11.1 Preparation of myeloma cells	23
11.2 Preparation of immunized spleen cells	23
11.3 Preparation of feeder cells	24
11.4 Fusion procedure	24
11.5 Limiting dilution	25
12. Screening of immunized sera and hybridoma culture supernatants	26
12.1 Indirect enzyme-linked immunosorbent assay (indirect ELISA)	26
12.2 Western blotting	27
13. Isotyping of mAbs	28
14. Purification of IgG from rabbit antiserum to L3	29
15. Purification of mAbs from culture supernatants	29
16. Biotinylation of Igs	29
17. Labeling of Igs with peroxidase	30
18. Absorption of biotinylated rIgG α L3	30
19. Fecal supernatant preparation	31
20. Preparation of pooled positive and negative fecal samples	31
21. Coproantigen sandwich ELISA	31
22. Statistical analyses	32
Results	33
1. Examination of fecal samples by the formalin-ether concentration method	33

2. Density of <i>S. stercoralis</i> larvae in fecal samples	33
3. Collection of adult worms, L3 and L1 for antigen preparation	37
4. Determination of protein content of <i>S. stercoralis</i> antigen	37
5. Demonstration of the 41-kDa protein in L1, L3 and adult worms of <i>S. stercoralis</i>	39
6. Production of mAbs	39
7. Sandwich ELISA	47
8. Coproantigen detection by sandwich ELISA	47
Discussion	53
References	58
Appendix	69
Vita	75

List of Tables

Table	Page
1. Number of stool samples positive for intestinal parasites by formalin-ether concentration method	34
2. Stool examinations by agar plate culture as compared to formalin-ether concentration method	35
3. Densities of <i>S. stercoralis</i> larvae examined by Beaver method	36
4. The ratio between adult female worms and L1 of <i>S. stercoralis</i> after overnight culture of adult worms	38
5. Amounts of proteins extracted from L3 of <i>S. stercoralis</i> as determined by the BCA protein assay kit	38
6. Optical densities of mAbs reacting with antigens of various stages of <i>S. stercoralis</i> by ELISA.	46
7. Optical densities of sandwich ELISA using mAbs as capture antibodies	48
8. Optical densities of sandwich ELISA, using rIgG α L3 as capture antibodies	48
9. Optical densities of sandwich ELISA tested on pooled positive and pooled negative feces.	50
10. Coproantigen sandwich ELISA of stool samples spiked with <i>S. stercoralis</i> antigens	52

List of Figures

Figure	Page
1. Western blot analysis of proteins of <i>S. stercoralis</i>	40
2. Western blot analysis of the mAbs from the first fusion	41
3. Western blot analysis of hybridoma supernatants from the fourth fusion	43
4. Western blot analysis of the culture supernatants from the seventh fusion	45
5. Optical densities of sandwich ELISA of 42 <i>S. stercoralis</i> -positive fecal samples and 40 negative fecal samples	51

Abbreviations

°C	degree Celsius
µg	microgram
µl	microliter
2-ME	2-mercaptoethanol
BaCl ₂	barium chloride
BSA	Bovine serum albumin
DAB	3,3'-diaminobenzidine tetrahydrochloride
EDTA	ethylenediaminetetraacetic acid
ELISA	enzyme-linked immunosorbent assay
FCS	fetal calf serum
g	gram
g	gravity
H ₂ SO ₄	sulfuric acid
HAT	hypoxanthine aminopterin thymidine
HCl	hydrochloric acid
hr	hour
HT	hypoxanthine thymidine
IFA	indirectfluorescent assay
Ig	immunoglobulin
IgG	immunoglobulin G
IgM	immunoglobulin M
IHA	indirecthaemagglutination assay
IMDM	Iscoe's Modified Dulbecco's Medium
kDa	kilodalton
L	liter
L1,L2	rhabditiform larvae
L3	filariform larvae
M	Molar

mAb	monoclonal antibody
mg	milligram
min	minute
ml	milliliter
N	normal
Na ₂ CO ₃	sodium carbonate
Na ₂ SO ₄	sodium sulfate
nm	nanometer
NSS	normal saline solution
OD	optical density
OPD	<i>o</i> -phenylenediamine
PBS	phosphate buffer saline
PEG	polyethylene glycol
PMSF	phenylmethylsulfonyl fluoride
rIgG α L3	Rabbit IgG against L3 antigens
SDS	sodium dodecyl sulfate
SDS-PAGE	sodium dodecyl sulfate polyacrylamide gel electrophoresis
Tween 20	polyoxyethylenesorbitan monolaurate
V	volts