

REFERENCE

- Altaman, D.F. (1998). Drugs used in Gastrointestinal Disease. In B.G. Katzung (Ed.), *Basic & Clinical Pharmacology*, (7th ed., pp. 1017-1029). USA: Prentice Hall International.
- Al-Yahya, M.A., Rafatullah, S., Mossa, J.S., Ageel, A.M., Parmar, N.S. and Tariq, M. (1989). Gastroprotective activity of ginger *Zingiber officinale* rosc., in albino rats. *Am. J. Chin. Med.*, 17(1-2), 51-56.
- Best, R., Lewis, D.A. and Nasser, N. (1984). The anti ulcerogenic activity of unripe plantain banana (*Musa* spp.). *Br. J. Pharmacol.*, 82, 107-116.
- Bhakuni, D.S., Dhar, M.L., Dhar, M.M., Dhawan, B.N., Gupta, B. and Srimali, R.C. (1971). Screening of Indian plants for biological activity. *Indian J. Exp. Biol.*, 9, 91.
- Brodie, D.A. and Hanson, H.M. (1960). A study of the factors involved in the production of gastric ulcers by the restraint technique. *Gastroenterology*, 38, 353-360.
- Brunton, L.L. (1996). Agents for control of gastric acidity and treatment of peptic ulcer. In J.G. Hardman, A.G. Gillman and L.E. Limbird (Eds.), *Goodman and Gilman's The Pharmacological Basis of Therapeutics* (9th ed., pp. 901-915). New York: Macmillan Publishing Co., Inc.
- Bullock, J. and Boyle, J. (1995). Stomach. In J. Bullock, J. Boyle and M. B. Wong (Eds). *Physiology* (3th ed., pp. 422-431). Philadelphia: Williams&Wilkins.
- Crawford, J.M. (1999). The Gastrointestinal Tract. In R.S. Cotran, V. Kumar and T. Collins (Eds). *Robbins: Pathologic Basis of Disease* (6th ed., pp. 755-829). USA: W.B. Saunders Company.

- Chuakul, W., Saralamp, P., Paonil, W., Temsiririrkkul R. and Clayton, T.(1997). *Croton oblongifolius* in *Medicinal Plants in Thailand Volume II* (pp. 84-85). Bangkok: Amarin Printing and Publishing.
- Corne, S.J., Morrissey, S.M. and Woods, R.J. (1994). A method for the quantitative estimation of gastric barrier mucous. *J. Physiol.*, 242, 116-117.
- Dajani, E.Z. and Agrawal, N.M. (1995). Prevention and treatment of ulcers induced by nonsteroidal anti-inflammatory drugs: an update. *J. Physiol. Pharmacol.*, 46,3-16.
- Dau, U.V., Ham, N.N. (1998). The effects of traditional drug, turmeric (*Curcuma longa*), and placebo on the healing of duodenal ulcer. *Phytomedicine*, 5, 29-34.
- Farnsworth, N.R., Blomster, R.N., Messmer, W.M., King, J.C. ,Persinos, G.J. and Wilkes, J.D. (1969). A phytochemical and biological review of the genus *Croton*. *Lloydia*, 32(1), 1-28.
- Friedman, L.S. and Peterson, W.L. (1998). Peptic ulcer and Related Disorder. In A.S Fanci, E. Braunwald, K.J. Isselbacher, J.D. Wilson. J. B. Martin, D.L. Kasper, S.L. Hanser and D.L Longo (Eds.), *Disorder of the Gastrointestinal System in Harrison's Principles of Internal Medicine* (14th ed., pp. 1597-1610). USA: Mcgraw Hill Companies Inc.
- Ganong, W.F. (1997). The Adrenal Medulla & Adrenal Cortex. In *Review of Medical Physiology* (18th ed. pp. 334-388), USA: Printice-Hall International, Inc.
- Gole, R.K., Gupta, S., Shankar, R. and Sanyal, A.K. (1986). Anti-ulcerogenic effect of banana powder (*Musa spientum var. paradisiaca*) and its effect on mucosal resistance. *J. Ethnopharmacol.*, 18(1), 33-34.
- Guth, P.H. (1972). Gastric blood flow in restraint stress. *Dig. Dis. Sci.*, 17, 807-813.

- Guth, P.H., Paulsen, G. and Nagata, H. (1984). Histological and microculatory changes in alcohol-induced gastric lesions in the rat: Effect of prostaglandin cytoprotection. *Gastroenterology*, 87, 1183-1190.
- Hua, X.Y., Pahlen, S.E., Lundberg, J.M., Hammarstrom, S. and Hedqvist, P. (1985). Leukotrienes C₄, D₄ and E₄ cause widespread and extensive plasma extravasation in the guinea pig. *Naunyn-Schmiedeberg's Arch. Pharmacol.*, 330 (2), 136-141.
- Hudson, N., Everitt, S. and Edwards, T. (1991). Elevation of gastric mucosal leukotriene B₄ levels of patients on long-standing NSAIDs therapy. *Gastroenterology*, 100, A86.
- Ito, S. and Lacy, E.R. (1985). Morphology of rat gastric mucosal damage, defence and restitution in the presence of luminal ethanol. *Gastroenterology*, 88, 250-260.
- Jain, S.P. (1989). Tribal remedies from Saranda forest, Bihar, India-I. *Intern. J. Crude Drug Res.*, 27(1), 29-32.
- Kitagawa, H., Fujwara, M. and Osumi, V. (1979). Effect of water immersion stress on gastric secretion and mucosal blood flow in rats. *Gastroenterology*, 77, 298-302.
- Kobayashi, S. and Tabata, K. (1992). Pharmacology of plaunotol, an anti-ulcer agent extract from a Thai medicinal plant (plau-noi). *Thai J. Pharm. Sci.*, 16, 328.
- Koga, T., Watanabe, H., Kawada, H., Takahashi, K., Utsui, Y., Domon, H., Ishii, C., Narita, T. and Yasuda, H. (1998). Interactions of plaunotol with bacterial membranes. *J. Antimicrob. Chemother.*, 42, 133-140.
- Konturek, S.J., Brzozowski, T., Piastucki, I., Radecki, T. and Dembinska-Kiec, A. (1983). Role of prostaglandin and thromboxane biosynthesis in gastric necrosis produced by taurocholate and ethanol. *Dig. Dis. Sci.* 28, 154-160.

- Konturek, S.J., Brzozowski, T., Drozdowicz, D. and Beck, G. (1988). Role of leukotrienes in acute gastric lesions induced by ethanol, taurocholate, aspirin, platelet activating factor and stress in rats. *Dig. Dis. Sci.*, 33, 806-813.
- Lange, K., Peskar, B.A. and Peskar, B.M. (1985). Stimulation of rat mucosal leukotriene formation by ethanol. *Naunyn-Schmiedeberg's Arch. Pharmacol.*, 3305, R27.
- Lanza, L.L., Walker, A.M., Bortnichak, E.A. and Dreyer, N.A. (1995). Peptic ulcer and gastrointestinal haemorrhage associated with nonsteroidal anti-inflammatory drug use in patients younger than 65 years. *Arch. Intern. Med.*, 155, 1371-1377.
- Lau, A.T.S., Graham, G.G., Dry, R.O. and Perry, M.A. (1992). Effect of aspirin on ulcer site blood flow in cat stomachs. *Am. J. Physiol.*, 263(26), G155-160.
- Maity, S., Vedasiromoni, J.R. and Ganguly, D.K. (1995). Anti-ulcer effect of the hot water extract of black tea (*Camellia sinensis*). *J. Ethnopharmacol.*, 46, 167-174.
- Maity, S., Vedasiromoni, J.R. and Ganguly, D.K. (1998). Role of glutathione in the Anti-ulcer effect of the hot water extract of black tea (*Camellia sinensis*). *Jpn. J. Pharmacol.*, 78, 285-292.
- Malone, M.H. and Robichaud, R.C. (1962). A Hippocratic screening for pure or crude drugs materials. *Lloydia*, 25, 320-332.
- Manatsathit, S., Kachintorn, U., Boonyapisit, S., Loengrojanakul, P., Tanwadee, T. and Chinapak, O. (1993). Gastric acid secretion in Thai patients with particular reference to the characteristics differentiating types of gastric and duodenal ulcer. *J. Med. Assoc. Thai*, 76, 677-681.
- Marhuenda, E., Martin, M. J., Alarcon dela Lastra, C. (1993). Antiulcerogenic activity of aescine in different experimental models. *Phytotherapy Res.*, 7, 13-16.

- Miller, T.A. (1983). Protective effects of prostaglandins against gastric mucosal damage: Current knowledge and proposed mechanisms. *Am. J. Physiol.* 245, G610-G623.
- Miyoshi, A., Asaki, S., Yoshida, Y., Goto, Y., Asaki, S., and Ishikawa, M. (1985). A clinical trial of CS-684; plaunotol in gastric ulcer patients (phase III). *Prog. Med.* 5, Suppl. 1, 981-1002.
- Mizui, T. and Doteuchi, M. (1988). Effect of polyamines on acidified ethanol-induced gastric lesions in rats. *Jpn. J. Pharmacol.*, 33, 939-945.
- Mokkhasmit, M., Swatdimongkol, K. and Satrawaha P. (1971a). Study on toxicity of Thai medication plants. *Bull. Sci.*, 12(2/4), 36-65.
- Mokkhasmit, M., Ngarmwathana, W., Swatdimongkol, K. and Permiphat,U. (1971b). Pharmacological evaluation of Thai medicinal plants. *J. Med. Assoc. Thailand*, 54(7), 490-504.
- Morimoto, Y., Shimohara, K., Oshima, S. and Sukamoto, T. (1991). Effect of the new anti-ulcer agent KB-5492 on experimental gastric mucosal lesions and gastric mucosal defensive factor, as compared to those of Teprenone and Cimetidine. *Jpn. J. Pharmacol.*, 57, 495-505.
- Mudgal, V. and Pal, D.C. (1980). Medicinal plants used by tribals of Mayurbhanj. (Orissa.) *Bull. Bot. Surv. India.*, 22, 59-62.
- Murakami, M., Lam, S.K., Inada, M. and Miyake, T. (1985). Pathophysiology and pathogenesis of acute gastric mucosal lesions after hypothermic restraint stress in rats. *Gastroenterology*, 88, 660-665.
- Nayar, S.L. (1954). Poisonous seed of India: Part I. *J. Bombay. Nat. Hist. Soc.*, 52(1), 1-18.

- Nayar, S.L.(1955). Vetgetable insecticides. *Bull. Nat. Instit. Sci. India*, 4, 137-145.
- Ogiso, A., Kitazawa, E., Mikuriya, I. and Promdej, C. (1981). Original plants of Thai crude drug, Plau-noi. *Shoyakugaku Zasshi*, 35, 287-290.
- Pamaham, M.J. and Brune, K. (1987). Therapeutic control of inflammatory diseases. *Agents Actions*, 21, 232-234.
- Rainsford, K.D. (1987). The effect of 5-lipoxygenase inhibitors and leukotriene antagonists on the development of gastric lesions induced by nonsteroidal inflammatory drugs in mice. *Agents Actions*, 21, 316-319.
- Rafatullah, S., Tariq, M., Al-Yahya, M.A., Mossa, J.S. and Ageel, A.M. (1990). Evaluation of turmeric (*Curcuma longa*) for gastric and duodenal antiulcer activity in rat. *J. Ethnopharmacol.*, 29(1), 25-34.
- Richardson, C.T. (1989). Gastric ulcer. In M.H. Sleisenger and J.S. Fordtran (Eds.), *Gastrointestinal Disease Pathophysiology/ Diagnosis/ Management* (4th ed., pp. 879-925). Philadelphia: W.B. Saunders.
- Robert, A., Nezamine, J.E., Lancaster, C. and Honcher, A.J. (1979). Cytoprotective by prostaglandins in rats: Prevention of gastric necrosis produced by alcohol, HCl, NaOH, hypertonic NaCl and thermal injury. *Gastroenterology*, 77, 433-443.
- Robert, A., Nezemis, j.E., Lancaster, C., Davis, J.P. and Field, S.D. (1983). Mild irritants prevent gastric necrosis through "adaptive cytoprotection" mediated by prostaglandins. *Am. J. Physiol.*, 245, 113-121.
- Roengsumran, S., Achayindee, S., Petsom, A., Pudhom, K., Singtothong, P., Surachetapan, C. and Vilaivan, T. (1998). Two new cembranoids from *Croton oblongifolius*. *J. Nat. Prod.*, 61, 652-654.

- Roengsumran, S., Petsom, A., Sommit, D. and Vilaivan, T. (1999). Labdane diterpenoids from *Croton oblongifolius*. *Phytochemistry*, 50, 449-453.
- Sabnis, S.D. and Bedi, S.J. (1983). Ethnobotanical studies in Dadra-Nagar Haveli and Daman. *Indian. J. For.*, 6(1), 65-69.
- Selvanayahgma, Z.E., Gnanevendhan, S.G., Balakrishna, K. and Rao, R.B. (1994). Antisnake venom botanicals from ethnomedicine. *J. Herbs Spices Med. Plants*, 2(4), 45-100.
- Shay, H., David, C.H., Sun, M.D. and Gruensteil, M. (1945). A sample method for the uniform production of gastric ulceration in the rat. *Gastroenterology*, 5, 43-61.
- Shiratori, K., Watanabe, S., Takeuchi, T., Chang, J.H. and Moriyoshi, Y. (1989). Effect of plaunotol on release of plasma secretin and pancreatic exocrine secretion in humans. *Pancreas*, 4(3), 323-328.
- Souza, A.R.M., Rodriguez, J.A., Hiruma-Lima, C.A., Haun, M. and Nunes, D.S. (1998). Antiulcerogenic activity of *trans*-dehydrocrotonin from *Croton cajucara*. *Planta Med.*, 64, 126-129.
- Subramanian, S.S., Nagarajan, S. and Sulochana, N. (1971). Flavonoids of some Euphorbiaceae plants. *Phytochemistry*, 10, 2548-2549.
- Taha, A.S., Sturrock, R.D. and Russel, R.I. (1995). Mucosal erosions in longterm nonsteroidal anti-inflammatory drug users: predisposition to ulceration and relation to *Helicobacter pylori*. *Gut*, 36, 334-336.
- Takagi, T., Kasuya, Y. and Watanabe, K. (1963). Studies on the drugs of peptic ulcer: A reliable method for producing stress ulcer in rats. *Chem. Pharm. Bull.*, 12, 465-472.

Takeuchi, K., Okada, M., Niida, H. and Okabe, S. (1989). Role of sulfhydryls in mucosal injury caused by ethanol: relation to microvascular permeability, gastric motility and cytoprotection. *J. Pharmacol. Exp. Ther.*, 248, 836-841.

Yamahara, J., Mochizuki, M., Fujimura, F., Takaishi, Y., Yoshida, M., Tomimatsu, T. and Tamai, Y. (1988). The anti-ulcer effect in rats of ginger constituents. *J. Ethnopharmacol.*, 23, 299-304.

Yamamoto, K., Kakegawa, H., Ueda, H., Matsumoto, H., Sudo, T., Miki, T. and Satoh, T. (1992). Gastric cytoprotective anti-ulcerogenic actions of hydroxychalcones in rats. *Planta Med.*, 58(5), 389-393.