

## TABLE OF CONTENTS

	Page
Acknowledgements	iii
Abstract (in English)	v
Abstract (in Thai)	vii
List of Tables	x
List of Figures	xiii
List of Appendices	xviii
Abbreviations	xix
Chapter 1 Introduction	1
Chapter 2 Literature Review	7
Chapter 3 Study site	25
Chapter 4 Methodology	32
Chapter 5 Results	44
Chapter 6 Discussion	99
Chapter 7 Conclusions and Recommendations	110
References	112
Appendices	125
<i>Curriculum Vitae</i>	154

### LIST OF TABLES

Table	Page
1 Soil characteristics of the study site (n=16) compared with those in undisturbed primary evergreen forest (Reusee Cave, east side of Doi Suthep, elevation 1,100 m about 9 km from the study site)(n=20)	29
2 Number of seedlings and number of seedling species divided by dispersal means in all plots	44
3 Number of seedlings found in each species from all isolated tree species combined	46
4 Average density (no./m <sup>2</sup> ) and species richness (no.species/m <sup>2</sup> ) of total seedlings in each studied tree plots	50
5 Average density (no./m <sup>2</sup> ) and species richness (no.species/m <sup>2</sup> ) of wind-dispersed seedlings in each studied tree plots	51
6 Average density (no./m <sup>2</sup> ) and species richness (no.species/m <sup>2</sup> ) of animal-dispersed seedlings in each studied tree plots	52
7 Mean ratios of animal / wind seedlings using number of individuals and number of species	53
8 Ecological indices of natural tree seedlings in each plots	55
9 Similarity coefficient (Sorensen's index) and difference coefficient (Chord distance; CRD) of natural seedling communities between tree crowns and control plots	75

**LIST OF TABLES (CONTINUED)**

Table	Page
10 Mean Sorensen's index and mean Chord distance of natural seedling communities between tree crowns and control plots	77
11 Summary results of <i>Kruskal Wallis</i> -test on density (no./m <sup>2</sup> ) and species richness (no.species/m <sup>2</sup> ) of seedling establishment among the 7 species of isolated tree studied	78
12 Summary results of <i>Mann Whitney</i> -test on average density (no./m <sup>2</sup> ) of seedling establishment beneath isolated tree species (P<0.05)	79
13 Summary results of <i>Mann Whitney</i> -test on average species richness (no.species/m <sup>2</sup> ) of seedling establishment beneath isolated tree species (P<0.05)	80
14 Summary results of <i>Mann Whitney</i> -test on average density (no./m <sup>2</sup> ) of seedling establishment between control plots of remnant tree species (P<0.05)	80
15 Summary results of <i>Mann Whitney</i> -test on average species richness (no.species/m <sup>2</sup> ) of seedling establishment between control plots of remnant tree species (P<0.05)	81
16 Summary results of linear-regression test between remnant tree size with density (no./m <sup>2</sup> ) and species richness (no.species/m <sup>2</sup> ) of natural seedling establishment beneath similar remnant tree species	82

**LIST OF TABLES (CONTINUED)**

Table	Page
17 Summary results of linear-regression test between remnant tree size with density (no./m <sup>2</sup> ) and species richness (no.species/m <sup>2</sup> ) of wind-dispersed seedlings beneath similar remnant tree species	83
18 Summary results of linear-regression test between remnant tree size with density (no./m <sup>2</sup> ) and species richness (no.species/m <sup>2</sup> ) of animal-dispersed seedlings beneath similar remnant tree species	84
19 Average relative growth rate (% per year) of naturally established tree seedlings	86
20 <i>Mann Whitney</i> -test on average RGRs of 10 seedling species between beneath remnant tree crowns and in control plots	90
21 Ratios of minutes bird observed / total observation minutes for each tree species	92
22 Percent of individual visit minutes by each bird group (according to diet) in the trees studied	93
23 Ratios of minutes bird observed / total observation minutes for each fruiting tree species	95

## LIST OF FIGURES

Figure	Page
1 The loss of viability of tree seeds from the seed bank shows that most trees tested have no dormancy	8
2 Map of Doi Suthep-Pui National Park, Chiang Mai	26
3 Average monthly temperature and rainfall at Nong Hoi Highland Agricultural Research Station	27
4 Average monthly minimum and maximum temperature at Nong Hoi Highland Agricultural Research Station	27
5 Land use map showing the deforested sites in Doi Suthep-Pui National Park, and Mae Sa Mai village	31
6 Materials and equipment	40
7 <i>Albizia chinensis</i> (Obs.) Merr. (Leguminosae, Mimosoideae)	40
8 <i>Callicarpa arborea</i> Roxb. var. <i>arborea</i> (Verbenaceae)	41
9 <i>Castanopsis diversifolia</i> (Kurz) King ex Hk. f. (Fagaceae)	41
10 <i>Erythrina stricta</i> Roxb. (Leguminosae, Papilionoideae)	42
11 <i>Eucalyptus camaldulensis</i> Dehnh. (Myrtaceae)	42
12 <i>Pinus kesiya</i> Roy. ex Gord. (Pinaceae)	43
13 <i>Schima wallichii</i> (DC.) Korth. (Theaceae)	43
14 Species/area curves of wind-dispersed seedlings in <i>Albizia chinensis</i> plots	59

**LIST OF FIGURES (CONTINUED)**

Figure	Page
15 Species/area curves of animal-dispersed seedlings in <i>Albizia chinensis</i> plots	59
16 Species/area curves of natural tree seedlings in <i>Albizia chinensis</i> plots	59
17 Species/area curves of wind-dispersed seedlings in <i>Callicarpa arborea</i> var. <i>arborea</i> plots	60
18 Species/area curves of animal-dispersed seedlings in <i>Callicarpa arborea</i> var. <i>arborea</i> plots	60
19 Species/area curves of natural tree seedlings in <i>Callicarpa arborea</i> var. <i>arborea</i> plots	60
20 Species/area curves of wind-dispersed seedlings in <i>Castanopsis diversifolia</i> plots	61
21 Species/area curves of animal-dispersed seedlings in <i>Castanopsis</i> <i>diversifolia</i> plots	61
22 Species/area curves of natural tree seedlings in <i>Castanopsis diversifolia</i> plots	61
23 Species/area curves of wind-dispersed seedlings in <i>Erythrina stricta</i> plots	62
24 Species/area curves of animal-dispersed seedlings in <i>Erythrina stricta</i> plots	62
25 Species/area curves of natural tree seedlings in <i>Erythrina stricta</i> plots	62
26 Species/area curves of wind-dispersed seedlings in <i>Eucalyptus</i> <i>camaldulensis</i> plots	63

**LIST OF FIGURES (CONTINUED)**

Figure	Page
27 Species/area curves of animal-dispersed seedlings in <i>Eucalyptus camaldulensis</i> plots	63
28 Species/area curves of natural tree seedlings in <i>Eucalyptus camaldulensis</i> plots	63
29 Species/area curves of wind-dispersed seedlings in <i>Pinus kesiya</i> plots	64
30 Species/area curves of animal-dispersed seedlings in <i>Pinus kesiya</i> plots	64
31 Species/area curves of natural tree seedlings in <i>Pinus kesiya</i> plots	64
32 Species/area curves of wind-dispersed seedlings in <i>Schima wallichii</i> plots	65
33 Species/area curves of animal-dispersed seedlings in <i>Schima wallichii</i> plots	65
34 Species/area curves of natural tree seedlings in <i>Schima wallichii</i> plots	65
35 Rarefraction curves of wind-dispersed seedlings in <i>Albizia chinensis</i> plots	67
36 Rarefraction curves of animal-dispersed seedlings in <i>Albizia chinensis</i> plots	67
37 Rarefraction curves of natural tree seedlings in <i>Albizia chinensis</i> plots	67
38 Rarefraction curves of wind-dispersed seedlings in <i>Callicarpa arborea</i> var. <i>arborea</i> plots	68
39 Rarefraction curves of animal-dispersed seedlings in <i>Callicarpa arborea</i> var. <i>arborea</i> plots	68
40 Rarefraction curves of natural tree seedlings in <i>Callicarpa arborea</i> var. <i>arborea</i> plots	68

**LIST OF FIGURES (CONTINUED)**

Figure	Page
41 Rarefraction curves of wind-dispersed seedlings in <i>Castanopsis diversifolia</i> plots	69
42 Rarefraction curves of animal-dispersed seedlings in <i>Castanopsis diversifolia</i> plots	69
43 Rarefraction curves of natural tree seedlings in <i>Castanopsis diversifolia</i> plots	69
44 Rarefraction curves of wind-dispersed seedlings in <i>Erythrina stricta</i> plots	70
45 Rarefraction curves of animal-dispersed seedlings in <i>Erythrina stricta</i> plots	70
46 Rarefraction curves of natural tree seedlings in <i>Erythrina stricta</i> plots	70
47 Rarefraction curves of wind-dispersed seedlings in <i>Eucalyptus camaldulensis</i> plots	71
48 Rarefraction curves of animal-dispersed seedlings in <i>Eucalyptus camaldulensis</i> plots	71
49 Rarefraction curves of natural tree seedlings in <i>Eucalyptus camaldulensis</i> plots	71
50 Rarefraction curves of wind-dispersed seedlings in <i>Pinus kesiya</i> plots	72
51 Rarefraction curves of animal-dispersed seedlings in <i>Pinus kesiya</i> plots	72
52 Rarefraction curves of natural tree seedlings in <i>Pinus kesiya</i> plots	72
53 Rarefraction curves of wind-dispersed seedlings in <i>Schima wallichii</i> plots	73

**LIST OF FIGURES (CONTINUED)**

Figure	Page
54 Rarefraction curves of animal-dispersed seedlings in <i>Schima wallichii</i> plots	73
55 Rarefraction curves of natural tree seedlings in <i>Schima wallichii</i> plots	73
56 Frequency of seedling species beneath remnant tree crowns separated by periods of average RGR	89
57 Frequency of seedling species in control plots separated by periods of average RGR	89
58 Frequency of seedling species in both, beneath tree crowns and control plots separated by periods of average RGR	89

**LIST OF APPENDICES**

Appendix	Page
A Remnant trees studied	126
B List of naturally established tree seedling species in the study plots	132
C Quantity of naturally tree seedlings in the study plots	136
D R squares of linear regressions between three parameters of remnant tree size and the density of natural seedlings, established beneath their crowns	144
E Average relative growth rate (% per year) of naturally established tree seedlings ( $h > 10$ cm) in each remnant tree species plot	146
F List of birds observed in this study	152

**ABBREVIATIONS**

ANR : accelerated natural regeneration

cm : centimetre

FORRU : Forest Restoration Research Unit

GBH : girth at breast height

km<sup>2</sup> : square kilometres

m<sup>2</sup> : square metres

no. : number

RFD : Thai Royal Forest Department

RGR : relative growth rate

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่

Copyright© by Chiang Mai University

All rights reserved