

VII. REFERENCES

1. Silpapojakul K. Scrub typhus in Western Pacific region. Ann Acad Med Singapore 1997; 26:794-800.
2. Brown GW , Robinson DM, Huxsoll DL, Ng TS, Lim KJ and Sannasey G. Scrub typhus: a common cause of illness indigenous populations. Trans. R. Soc. Trop. Med. Hyg 1976; 70: 444-8.
3. Brown GW, Saunders JP, Singh S, Huxsoll DL and Shirai A. Single dose doxycycline therapy for scrub typhus. Trans. R. Soc. Trop. Med. Hyg 1978; 72: 412- 416.
4. Kelly DJ, Wong PW, Gan E and Lewis GE Jr. Compative evaluation of the indirect immunoperoxidase test for the serodiagnosis of rickettsial disease. Am. J. Med. Hyg 1988; 38: 400- 6.
5. Kawamura A. Jr, Tanaka H and Tamura A. Tsutsugamushi Disease. University of Tokyo Press. 3-1 Hongo 7 Bankyo- ku Tokyo, Japan. 1995; 275- 306.
6. Fredricks DN and Relman DH. Application of polymerase chain reaction to the diagnosis of infectious diseases. Clin. Infect. Dis 1999; 29: 475-486.
7. Eisemamm CS and Osterman JV. Identification of strain-specific and group-reactive antigenic determinants on the Karp, Gilliam and Kato strains of *Rickettsia tsutsugamushi*. Am. J. Trop. Med. Hyg 1985; 34 : 1173-1178.
8. Hanson B. Identification and partial characterization of *Rickettsia tsutsugamushi* major protein immunogens. Infect. Immun 1985; 50: 603- 609.
9. Ewing HE. Jr. Takeuchi A, Shirai A and Osterman JV. Experimental infection of mouse peritoneal mesothelium with scrub typhus rickettsiae: an ultrastructural study. Infect. Immun 1978; 19: 1068-1075.
10. Tamura, A., Urakami, H. and Ohashi, N. 1990b. A comparative view of *Rickettsia tsutsugamushi* and the other groups of rickettsiae. Eur.J.Epidemiol 1978; 7: 259- 269.
11. Krieg NR and Holt JG. Bergey's manual of systematic bacteriology. Baltimore and London, Williams and wilkins, 1984; 1:687- 698
12. Takahashi K, Urakami H and Tamura A. Purification of cell envelopes of *Rickettsia tsutsugamushi*. Microbiol. Immunol 1985; 29: 475-8.

13. Tamura A, Ohashi N, Urakami H and Miyamura S. Classification of *Rickettsia tsutsugamushi* in a new genus, *Orientia* gen. Nov., as *Orientia tsutsugamushi* comb. Nov. Int. J. Syst. Bacteriol. 1995; 45:589-591.
14. Amono K, Tamura A, Ohashi N, Urakami H, Kaya S and Fukushi K. Deficiency of peptidoglycan and lipopolysaccharide components in *Rickettsia tsutsugamushi*. Infect. Immun 1987; 55: 2290- 2292.
15. Miyamura S, Ohta T and Tamura A. Comparison of *in vitro* susceptibilities of *Rickettsia tsutsugamushi* to chemotherapeutic agents J. Japan, Assoc. Infect. Dis 1989; 59: 486- 488. (in Japanese).
16. Urakami H, Tamura A, Miyamura S, Yamamoto S and Kawabata N. Susceptibilities of recent colinical isolates of *Rickettsia tsutsugamushi* to chemotherapeutic agents *in vitro*, J. Japan.Assoc. Infect. Dis 1988a;. 62: 931- 937. (in Japanese).
17. Myers WF, Baca OG and Wisseman CL. Jr. Genomic size of the rickettsia *Coziellaburnetii*. J. Bacteriol 1980; 144: 460- 461.
18. Myers WF and Wisseman CL. Jr. The taxonomic relationship of *Rickettsia Canada* to the typhus and spotted fever groups of the genes *Rickettsia*. In: Burgdorfer W and Anacker RL eds. *Rickettsiae and rickettsial diseases*. New York, London, Tront, Sydney, San Francisco. Academic Press 1981; 313-325
19. Shimada M, Ohashi N and Tamura A. Analysis of genomic DNA structure in *Rickettsia tsutsugamushi* by pulse field gel electrophoresis (AB). Japan. J.Bacteriol 1994; 49: 293. (in Japanese).
20. Tamura A, Ohashi N, Urakami H, Takahashi K and Oyanagi M. Analysis of polypeptide composition and antigenic components of *Rickettsia tsutsugamushi* by polyacrylamide gel electrophoresis and immunoblotting. Infect. Immun., 1985; 48: 671- 675.
21. Ohashi N, Tamura A and Suto T. Immunoblotting analysis of anti-rickettsial antibodies produced in patients of Tsutsugamushi disease. Microbiol. Immunol 1988; 32: 1085- 1092.
22. Smirnova NS, Popov VL and Kokorin IN. Peculiarities of *Rickettsia prowazekii* in the cell culture as revealed by cryoultramicrotomy. Acta Virol 1986; 30: 436- 439.
23. Silverman DJ, Wisseman CL. Jr and Waddell A. In vitro studies of rickettsia host cell interaction: ultrastructural study of *Rickettsia prowazekii* infected chicken embryo fibroblasts. Infect. Immun 1980; 29: 778- 790.

24. Wisseman CL. Jr, Waddell AD and Silverman DJ. *In vitro* studies on rickettsia-host cell interactions: lag phase in intracellular growth cycle as a function of stage of growth of infecting *Rickettsia prowazeki*, with preliminary observations on inhibition of rickettsial uptake by host cell fragments. *Infect. Immun* 1976; 13: 1749-1760.
25. Ito S, Vinson JW and Whitescarver J. Ultrastructural observations of *Rickettsia typhi* in the human body louse. In: Kazar J, Ormsbee RA. and Tarasevich In eds. *Rickettsiae and rickettsial diseases*. Bratislava VEDA Publ. House, Slovak Acad. Sci. 1978; 53- 64.
26. Lyle PB and Burgdorfer W. Fine structure of *Rickettsia Canada* in tissues of *Dermacentor andersoni* stiles. *J. Bacteriol* 1971; 105: 1149- 1159.
27. Silverman DJ, Wisseman CL. Jr, Waddell AD and Jones M. External layers of *Rickettsia prowazekii* and *Rickettsia rickettsii*: occurrence of a slime layer. *Infect. Immun* 1978; 22: 233-246.
28. Hayes SF and Burgdorfer W. Ultrastructure of *Rickettsia rhipicephali*, a new member of the spotted fever group rickettsiae tissues of the host vector *Rhipicephalus sanguineus*. *J. Bacteriol* 1979; 137: 605-613.
29. Wright JD, Hastriter MW and Robinson DM. Observations on the ultrastructure and distribution of *Rickettsia tsutsugamushi* in naturally infected *Leptotrombidium* (*Leptotrombidium*) *arenicola* (Acari: Trombiculidae). *J. Med. Entomol* 1984; 21: 17- 27.
30. Rikihisa Y and Ito S. Effect of antibody on entry of *Rickettsia tsutsugamushi* into polymorphonuclear leukocyte cytoplasm. *Infect. Immun* 1983; 39: 928- 938.
31. Urakami H, Tsuruhara T and Tamura A. Electron microscopic studies on intracellular multiplication of *Rickettsia tsutsugamushi* in L cells. *Microbiol Immunol* 1984; 28: 1191- 1201.
32. Urakami H, Takanashi M, Tamura A and Hori E. Electron microscopic observations of the embryo *Leptotrombidium* (*Leptotrombidium*) *pallidum* naturally infected with *Rickettsia tsutsugamushi*. *Microbiol. Immunol* 1988b; 32: 967- 972.
33. Ogata N, Nakajima G and Kajima S. Animals employed for experimental infection by pathogenic rickettsiae in laboratory- especially recommendation to use mouse for the detection of *Rickettsia tsutsugamushi*. *Tokyo Med. J* 1932; 2760: 155-160. (in Japanese).
34. Kundin WD, Liu C, Harmon P and Robison DM. Pathogenesis of scrub typhus infection (*Rickettsia tsutsugamushi*) as studied by immunofluorescence. *J. Immunol* 1964; 93: 772- 781.

35. Urakami H, Tsuruhara T and Tamura A. Observations of the same whole cells infected with *Rickettsia tsutsugamushi* by means of transmission and scanning electron microscopy. J. Electron Microsc 1982a; 31: 212- 215.
36. Urakami H, Tsurugara T and Tamura A. . Intranuclear *Rickettsia tsutsugamushi* in culture mouse fibroblasts (L cells). Microbiol. Immunol 1982b; 26: 445- 447.
37. Urakami H, Tsuruhara T and Tamura A. Penetration of *Rickettsia tsutsugamushi* into cultured mouse fibroblasts (L cells): and electron microscopic observation. Microbiol. Immunol 1983; 27: 251-263.
38. Kawamori F, Akiyama M, Sugida M, Kanda T, Akahane S, Uchikawa, Uamada Y, Kumada N, Furuya Y, Yoshida Y, Yamamoto S, Ohashi N and Tamura A. Epidemiology of Tsutsugamushi disease in relation to the serotypes of *Rickettsia tsutsugamushi* isolated from patients, field mice, and unfed chiggers on the eastern slope of Mount Fuji, Shizuoka Prefecture, Japan. J. Clin. Microbiol 1992; 30: 2842- 2846.
39. Kawamura A. Jr, Murata M, Osono M, Nogami S, Shirasaka A, Tanaka H, Sudo K, Suzuki K, Miyari T and Kijima H. Studies on inapparent infection of Tsutsugamushi disease in Izu Shichito Islands; seroepidemiology and demonstration of an avirulent rickettsia strain for mice. Jpn. J. Exp. Med 1980; 50: 91- 105.
40. Saito H, Sekikawa H and Kenmotsu M. Clinical review on new type tsutsugamushi disease from clinical cases in Niigata prefecture. Nihon Iji Shinpo 1981; 3007: 23- 30.
41. Bell EJ, Bennett BL and Whitman L. Antigenic differences between strains of scrub typhus as demonstrated by cross- neutralization test. Proc. Soc Exp. Biol. Med 1946; 2:134- 7.
42. Bebgton IA. Apparent serological heterogeneity among strains of Tsutsugamushi disease (scrub typhus). Publ. Hlth. Rep 1945; 60: 1483-88.
43. Bennett BL, Smadel JE and Gauld RL. Studies on scrub typhus (Tsutsugamushi disease), IV. Heterogeneity of strains of *R. tsutsugamushi* as demonstrated by cross-neutralization tests. J. Immunol 1949; 62: 453- 61.
44. Fox JP. The neutralization technique in tsutsugamushi disease (scrub typhus) and the antigenic differentiation of rickettsial strains. J. Immunol 1949; 62: 341- 352.
45. Shishido A, Ohtawara M, Hikita M and Kitaoka M. The nature of immunity against scrub typhus in mice. II. The cross-protection test with mice for identification and differentiation of

- several strains of *Rickettsia orientalis* newly isolated in Japan. Japan. J. Med. Sci. Biol 1959; 12: 391-404.
46. Kawamura A. Jr and Aoyama Y. Immunofluorescence in medical science. Tokyo: University of Tokyo Press, 1983; 1-262
47. Yamamoto S, Kawabata N, Ooura K, Murata M and Minamishima Y. Antigenic types of *Rickettsia tsutsugamushi* isolated from patients with tsutsugamushi fever and their distribution on Miyazaki Prefecture. J. Japan. Assoc. Infect. Dis 1989; 63: 109- 117.
48. Kobayashi Y, Kanemitsu N, Tange Y, Tachibana N and Tamara A. Immunological analysis of newly isolated strains of *Rickettsia tsutsugamushi* using monoclonal antibodies. J. Japan. Assoc. Infect. Dis 1987; 61: 1264- 1269.
49. Kim I-S, Seong S-Y, Woo. S-G, Choi M-S and Chang W-H . High-level expression of a 56-kilodalton protein gene (*bor 56*) of *Rickettsia tsutsugamushi* boryong and its application to enzyme- linked immunosorbent assay. J.Clin. Microbiol 1993a ; 31: 598- 605.
50. Chang WH. Current status of tsutsugamushi diseases in Korea. J. Korean Med. Sci 1995; 10:227-238.
51. Ohashi N, Koyama Y, Urakami H, Higo M and Tamura A. Characterization of *Rickettsia tsutsugamushi* isolated from rodents and chiggers in Taiwan (Ab). J. Japan. Bacteriol 1994; 49: 270. (in Japanese).
52. Guo HB, Yu MM, Xu MH, Tang J-G and Wu G-G. Studies on serological typing of anti-*Rickettsia* antibody in Jiangsu. Chines J. Microbiol. Immunol 1993; 13: 316- 317.
53. Furuya Y, Yoshida Y, Katayama T, Yamamoto and Kawamura A Jr. Serotype-specific Amplification of *Rickettsia tsutsugamushi* DNA by Nested Polymerase Chain Reaction. J.Clin. Microb. 1993; 31:1634- 40.
54. Kaiho I, Tokieda M, Yoshida Y, Furuya Y, Murata M, Tanaka H and Kawamura A. Jr. Epidemiology of Tsutsugamushi disease and typing of isolated *Rickettsia* in Chiba Prefecture. J. Japan. Assoc. Infect. Dis 1993; 67: 196-201. (in Japanese).
55. Shirai A, Tanskul PL, Andre RG, Dohane AI and Huxsoll DL. *Rickettsia tsutsugamushi* strains found in chiggers collected in Thailand. Southeast Asian J Trop Med Public Heath 1981; 12: 1-6.

56. Shirai A, Dohany AL, Ram S, Chiang GL and Huxsoll DL. Serological classification of *Rickettsia tsutsugamushi* organisms found in chiggers (Acarina: Trombiculidae) collected in Peninsular Malaysia. Trans. Roy. Soc. Trop. Med. Hyg 1981; 75: 580-582.
57. Shirai A, Robinson DM, Brown GW, Gan E and Huxsoll DL. Antigenic analysis by immunofluorescence of 114 isolates of *Rickettsia tsutsugamushi* recovered from febrile patients in rural Malaysia. Jpn J Med Sci Biol 1976; 32: 337- 344.
58. Eismann CS and Osterman JV. Antigens of scrub typhus rickettsiae separation by polyacrylamide gel electrophoresis and identification by enzyme-linked immunosorbent assay. Infect. Immun 1981; 32: 525-533.
59. Ohashi N, Tamura A, Ohta M and Hayashi K. Purification and partial characterization of type-specific antigen of *Rickettsia tsutsugamushi*. Infect. Immun., 1989; 57: 1427- 31.
60. Ohashi N, Nashimoto H, Ikeda H and Tamura A. Cloning and sequencing of the gene (*tsg 56*) encoding a type-specific antigen from *Rickettsia tsutsugamushi*. Gene 1990a; 91: 119- 122.
61. Stover CK, Marana DP, Carter JM, Roe BA, Mardis E and Oaks EV. The 56- kilodalton major protein antigen of *Rickettsia tsutsugamushi*: molecular cloning and sequence analysis of the *tsg 56* gene and precise identification of strain-specific epitope. Infect. Immun 1990; 58: 2076- 84.
62. Tamura A, Takahashi K, Tsuruhara T, Uekami H, Miyamura S, Sekikawa H, Kenmotsu M, Abe S and Nezu H. Isolation of *Rickettsia tsutsugamushi* antigenically different from Kato, Karp, and Gilliam strains from patients. Microbiol. Immunol 1984; 28: 873-882.
63. Oiver D. Protein secretion in *Escherichia coli*. Ann.Rev. Microbiol 1985; 39: 615-648.
64. Oaks EV, Stover CK and Rice RM. Molecular cloning and expression of *Rickettsia tsutsugamushi* genes for two major protein antigens in *Escherichia coli*. Infect. Immun 1987; 55: 1156-1162.
65. Hickman CJ, Stover CK, Joseph SW and Oaks EV. Molecular cloning and sequence analysis of a *Rickettsia tsutsugamushi* 22 kDa antigen containing B-and T-cell epitopes. Microb. Pathog 1991;11: 19-31.
66. Hanson B. Identification and partial characterization of *Rickettsia tsutsugamushi* major protein immunogens. Infect. Immun 1985; 50: 603-609.

67. Murata M, Yoshida Y, Osono M, Ohashi N, Oyanagi M and Kawamura A. Jr. Production and characterization of monoclonal strain-specific antibodies against prototype strains of *Rickettsia tsutsugamushi*. Microbiol. Immunol 1986; 30: 599-610.
68. Ohashi N, Nashimoto H, Ikeda H and Tamura A. Diversity of immunodominant 56-kDa type-specific antigen (TSA) of *Rickettsia tsutsugamushi*. Sequence and comparative analyses of the genes encoding TSA homologous from four antigenic variants. J.Biol. Chem 1992; 267: 12728- 35.
69. Oaks EV, Rice RM, Kelly DJ and Stover CK. Antigenic and genetic relatedness of eight *Rickettsia tsutsugamushi* antigens. Infect. Immun 1989; 57: 3116- 22.
70. Groves MG, Rosenstreich DL, Taylor BA and Osterman JV. Host defenses in experimental scrub typhus: mapping the gene that controls natural resistance in mice. J. Immunol 1980; 125: 1395- 1399.
71. Yamamoto S, Kawabata N, Murata M and Minamishima Y. Antigenic types of *Rickettsia tsutsugamushi* isolated from patients with Tsutsugamushi fever and their distribution in Miyazaki Prefecture. J. Japan. Assoc. Infect. Dis 1989; 63: 109- 117.
72. Weil E and Felix A. Zur serologischen Diagnose des Fleckfiebers. *Wien. Klin Wochenschr.*, 1916; 29: 33- 35.
73. Shishido A, Kohno S, Hikita M, Iida T, Kawashima H and Kawamura A Jr. Complement fixation and direct immunofluorescence for typing of Tsutsugamushi disease rickettsia. *Acta medica et Biologica* 1967; 15 (Suppl.): 87-95.
74. Iida T, Kawashima H and Kawanyra A. Jr. Direct immunofluorescence for typing of Tsutsugamushi disease rickettsia. J. Immunol 1966a; 95: 1129- 33.
75. Elisberg B.L, Sangkasuvana V, Campbell JM, Bozeman FM, Boahidatta P and Rapmud G.. Physiographic distribution of scrub typhus in Thailand. *Acta Medica et Viologica* 1967; 15 (Suppl.): 61- 67.
76. Elisberg BL, Needy CF and Bozeman FM. Antigenic inter-relationships among strains of *Rickettsia tsutsugamushi* In: Kazar J, Ormsbee RA. and Tarasevich IN editors. *Rickettsiae and Oriental diseases*. Bratislava: VEDA Publ. House Slovak. Acad. Sci. 1978; 253- 262.
77. Suto T. A ten years experience on diagnosis of Oriental diseases using the indirect immunoperoxidase methods. *Acta virol* 1991; 35: 580- 586.

78. Tamura A, Ohashi N, Sakurai H, Urakami H, Yamamoto S and Kawamori F. Serodiagnosis of scrub typhus by dot-blot method. J. Japan. Assoc. Inf. Dis 1990a; 64: 704- 714. (In Japanese).
79. Urakami H, Yamamoto S, Tsuruhara T, Ohashi and Tamura A. Serodiagnosis of scrub typhus with antigens immobilized on nitrocellulose sheet. J. Clin.Microbiol 1989; 27: 1841- 6.
80. Furuya Y, Yoshida Y, Katayama T, Kawamori F Yamamoto S, Ohashi N, Kamura A and Kawamura A. Jr. Specific amplification of *Rickettsia tsutsugamushi* DNA from clinical specimen by polymerase chain reaction. J. Clin. Microbiol. 1991; 29: 2628-30.
81. Sugita Y, Nagatani T, Okuda K, Yoshida Y and Nakajima H. Diagnosis of typhus infection with *Rickettsia tsutsugamushi* by polymerase chain reaction. J. Med. Microbiol 1992; 37: 357-360.
82. Sugita Y, Yamakawa Y, Takahashi K, Nagatani T, Okuda K and Nakajima H. A polymerase chain reaction system for rapid diagnosis of scrub typhus within six hours. Am. J. Trop. Med. Hyg 1993; 49: 636- 640.
83. Brown GW, Shirai A, Rogers C and Campbell JM. Diagnostic criteria for scrub typhus: probability values for immunofluorescent antibody and *Proteus* OX-K agglutinin titers. Am. J. Trop. Med. Hyg 1983; 32: 1101-1107.
84. Disease under surveillance. Division of Epidemiology, Ministry of Public Health, Thailand, Monthly Epidemiological Surveillance Report : 2000; 31: 404.
85. Monthly Epidemiological Surveillance Report 2000. Disease under surveillance. 31:(12) 404.
86. James C. Scrub typhus. Control of Communicable Diseases Manual, An official report of the American Public Health Association, Washington D.C 17thed. 2000; 545- 8.
87. Rollwagen FM, Dasch GA and Jerrells TR. Mechanisms of immunity to rickettsial infection: characterization of a cytotoxic effector cell. J. Immunol 1986; 136: 1418-21.
88. Nei M. Molecular evolutionary genetics. New York: Columbia University Press; 1987; 293-8.