

REFERENCES

- Adamsen, A.K., Lindhagen, L. and Ahring, B.K. (1995). Optimization of extracellular xylanase production by *Dityoglomus* sp B1 in continuous culture. *Appl. Microbiol. Biotechnol.* 44:327-332.
- Antanopoulos, V. T., Hernandez, M., Arias, M. E., Mavrakos, E. and Ball, A. S.(2001). The use of extracellular enzymes from *Streptomyces albus* ATCC 3005 for the bleaching of eucalyptus kraft pulp. *Appl. Microbiol. Biotechnol.* 57: 92-97
- Bajpai, P., Bhardwaj, N.K., Bajpai, P.K. and Jauhari, M.B.(1994). The impact of xylanases on bleaching of eucalyptus kraft pulp. *J Biotechnol*; 38:1– 6.
- Balakrishnan, H., Choudhury, M. D., Srinivasan, M. C. and Rele, M. V.(1992). Cellulase-free xylanase production from an alkalophilic *Bacillus* species. *World. J. Microbiol. Biotechnol.*, 8: 627-631.
- Bansod, S.M., Dutta-Choudhury, M., Srinivasan, M.C. and Rele, M.V. (1992). Xylanase active at high pH from an alkalotolerant *Cephalosporium* species. *Biotechnol. Lett.* 15:965-970.
- Bataillon, M., Cardinali, A.P.N., Castillon, N. and Duchiron, F.(2000). Purification and characterization of a moderately thermostable xylanase from *Bacillus* sp. strain SPS-0. *Enz. Microbiol. Technol.* 26:187-192.
- Beg, Q. K., Bhushan, B., Kapoor, M. and Hoondal, G. S. (2000). Enhanced production of a thermostable xylanase from *Streptomyces* sp. QG-11-3 and its application in biobleaching of eucalyptus kraft pulp. *Enz. Microbiol. Technol.*, 27: 459-466
- Beg, Q. K., Kapoor, M., Mahajan, L. and Hoondal, G. S. (2001). Microbial xylanases and their industrial applications: a review. *Appl. Environ. Microbiol.* 56:326–338.

- Bhardwaj, N.K., Bajpai, P. and Bajpai, P.K.(1996). Uses of enzymes in modification of fibers for improved beatability. *J. Biotechnol.*51:21-26.
- Biela, P. (1985). Microbial xylanolytic system. *Trends Biotech.* 3:286-290.
- Biely, M.J., Buchert, J. and Viikari, L.(1993). Effect of pH on production of xylanase by *Trichoderma reesei* on xylan and cellulose base media. *Appl. Microbiol. Biotechnol.* 40:224-229.
- Blanco, A., Vidal, T., Colom, J.F. and Paster, F.I.J. (1995). Purification and properties of xylanase A from alkali-tolerant *Bacillus* sp. Strain BP-23. *Appl. Environ. Microbiol.* 61(12):4468-4470.
- Box, G.E.P., Hunter, W.G. and Hunter, S.J.(1978). *Statistic of experiments.* John Wiley & Son, Inc., Newyork. Pp.35-353.
- Box, G.E.P. and Wilson, K.B. (1951). On the experimental attainment of optimum conditions. *J.Royal Statistic. Society.* 13;1-45
- Bradford M.M.. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding, *Anal. Biochem.* 72:248-254.
- Breccia, J.D., Siferiz, F., Baigori, M.D., Castro, G.R. and Hatti-Kaul, R.(1998). Purification and characterization of a thermostable xylanase from *Bacillus amyloliquefaciens*. *Enz. Microbiol. Technol.*22:42-49.
- Caroline, S.M. Furniss., Nigel, J. Belshaw., Marcos, J.C. Alcocer., Gary, Williamson., Giles, O. Elliott., Kurt, Gebruers., Nigel, P. Haigh., Neville, M. Fish.and Paul, A. Kroon. (2002). A family 11 xylanase from *Penicillium funiculosum* is strongly inhibited by three wheat xylanase inhibitors. *Biochimica et Biophysica Acta.* 1598; 24-29.

- Chen, C., Adolphson, R., Dean, J.F.D., Ericksson, K.L., Adam, M.W.W. and Westpheling, J. (1997). Release of lignin from kraft pulp by a hyperthermophilic xylanase from *Thermotoga maritima*. *Enz. Microbiol. Technol.* 20:39-45.
- Christakopoulos, P., Nerinckx, W., Kekos, D., Macris, B. and Claeysens, M.(1996). Purification and characterization of two low molecular mass alkaline xylanases from *Fusarium oxysporum* F3. *J. Biotechnol.* 51:181-189.
- Christov, L. P. and Prior, B. A. (1997). Bleaching Response of Sulfite Pulps to Pretreatment with Xylanases. *Biotechnol. Prog.* 13:695-698.
- Christov, L.P., Szakacs, G. and Balakrishnan, H.(1999). Production partial characterization and use of fungal cellulase-free xylanase in pulp bleaching. *Process Biochem.* 64:511-517.
- Christoph, W. and Wolfgang, L.(1995). Two extremely thermostable xylanases of the hyperthermophilic bacterium *Thermotoga maritima* MSB8. *Process Biochem.* 64:511-517.
- Clark, E.M., Tankanen, M., Setala, T.N. and Penttila, M. (1996). Cloning of gene encoding α -L-arabinofuranosidase and β -xylosidase from *Trichoderma reesei* by expression in *Saccharomyces cerevisiae*. *Appl. Environ. Microbiol.*62:3840-3846.
- Costa-Ferreira, M., Dias, A. Maximo, C., Morgado, M.J., Sena-Martins, G. and Duarts, J.C.(1994). Xylanolytic enzyme production by an *Aspergillus niger* isolate. *Appl. Biochem. Biotechnol.* 44:231-242.
- Dahlberg, L., Holst, O. and Kristjansson, J.K..(1993). Thermostable xylanolytic enzymes from *Rhodothermus marinus* grown on xylan. *Appl. Microbiol. Biotechnol.*40:63-68.

- De Souza, D.F., De Souza, C.G.M. and Peralta, R.M.(2001). Effect of easily metabolizable sugar in the production of xylanase by *Aspergillus tamaris* in solid-state fermentation. *Process Biochem.* 36:835-838.
- Dhillon, A., Gupta, J.K. and Khanna, S.(2000a). Enhanced production, purification and characterization of a novel cellulase-poor xylanase thermostable, alkalotolerant xylanase from *Bacillus circulans* AB16. *Process Biochem.* 35:849-856.
- Dhillon, A., Gupta, J.K. Jauhari, B.M. and Khanna, S.(2000b). A cellulase-poor, thermostable, alkalotolerant xylanase produced by *Bacillus circulans* AB16 grown on rice straw and its application in biobleaching of eucalyptus pulp. *Bioresource Technol.* 73:273-277.
- Dhillon, A and Khanna, S.(2000c). Production of thermostable alkalitolerant xylanase from *Bacillus circulans* AB16 grown on wheat straw, *J. Microbiol. Biotechnol.* 16 (4):325-327.
- Elegir, G., Sykes, G. and Jeffries, T.W.(1995). Differential and synergistic action of *Streptomyces* endoxylanase in prebleaching of kraft pulp. *Enz. Microbiol. Technol.* 17(10): 954-959.
- Elegir, G., Szakaacs, G. and Jeffries, T.W.(1994). Purification, characterization and substrate specificities of multiple xylanase from *Streptomyces* sp. strain B-12-2. *Appl. Environ. Microbiol.* 60(7): 2609-2615.
- Eriksson, K.E.L., Blanchette, R.A. and ander, P. (1990). Microbial and Enzymatic Degradation of Wood and Wood Components. Springer-Verlag Berlin Heidelberg Berlin: Germany.pp. 102-112,147,175.
- Fernandez-Espinar, M.T., Ramon, D., Pinaga, F. and Valles, S. (1992). Xylanase production by *Aspergillus nidulans*. *FEMS Microbiol.* 91:91-96.

- Garg, A.P., Roberts, J.C. and Jeffries, T.W.(1998). Purification, characterization and substrate specificities of multiple xylanases from *Streptomyces* sp. strain B-12-2. *Appl. Environ. Microbiol.* 60(7):2609-2615.
- George, S.P., Ahmad, A. and Rao, M.B. (2001a). Studies on carboxymethyl cellulase produced by an alkalothermophilic actinomycetes. *Bioresource Technol.* 77:171-175.
- George, S.P., Ahmad, A. and Rao, M.B. (2001b). A novel thermostable xylanase from *Thermomonospora* sp. :influence of additive on thermostability. *Bioresource Technol.* 7:221-224.
- Gessesse, A. and Mama, G.(1999). High-level xylanase production by an alkalophilic *Bacillus* sp. by using solid-state fermentation. *Enz. Microbiol. Technol.* 25:68-72.
- Gilbert, M., Breuil, C. and Saddler, J.N.(1992). Characterization of the enzymes present in the cellulase systems of *Thielavia terrestris* 225B. *Bioresource Technol.* 39:147-154.
- Gilead, S. and Shoham, Y. (1995). Purification and characterization of α -L-arabinofuranosidase from *Bacillus stearotherophilus* T-6. *Appl. Environ. Microbiol.* 61:170-174.
- Gomes, J., Gomes, I. And Steiner, W.(2000a). Thermostable xylanase of the Antarctic yeast *Cryptococcus adeliae*: production and properties. *Extremophiles.* 4:227-235.
- Gomes, J., Gomes, I., Terler, K., Gubala, N., Ditzelmuller, G. and Steiner, W.(2000b). Optimization of culture medium and condition for α -L-Arabinofuranosidase production by extreme thermophilic eubacterium *Rhodothermus marinus*. *Enz. Microbiol. Technol.* 27:414-422.
- Grabski,A.C. and Jeffries, T.W.(1991). Production, purification and characterization of β -(1,4)-endoxylanase of *Streptomyces roseiscleroticus*. *Appl. Environ. Microbiol.* 57:987-992.

- Gupta, S., Bhushan, B. and Hoondal, G.S.(2000). Isolation, purification and characterization of xylanase from *Staphylococcus* sp.SG-13 and its application in biobleaching of kraft pulp. *J. Appl. Microbiol.*88(2):325-334.
- Gupta, S., Kuhad, R.C., Bhushan, B. and Hoondal, G.S.(2001). Improved xylanase production from a haloalkalophilic *Staphylococcus* sp.SG-13 using inexpensive agricultural residues. *World J. Microbiol. Biotechnol.*17(1):5-8.
- Haltrich, D., Nidetzky, B., Kulbe, K.D., Steiner, W. and Zupancic, S.(1996). Production of fungal xylanases. *Bioresource Technol.* 58:137-163.
- Hinnman, R.L.(1994).The changing face of fermentation industry. *Chem. Technol.* 45-48.
- Inakaki, K., Nakahira, K., Mukai, K., Tamura, T. and Tanaka, H.(1998). Gene Cloning and characterization of an acidic xylanase from *Acidobacterium capsultum*. *Biosci. Biochem.* 62(4):1061-1067.
- Jain, A., Garg, S.K. and Johri, B.N.(1998). Properties of a thermostable xylanase production by *Melanocarpus albomyces* IIS-68 in solid-state fermentation. *Bioresource Technol.*64:255-228.
- Jeffries, T.W. (1992). Enzymatic treatment of pulp: emerging technologies for materials and chemicals for biomass. *ACS Symp.Ser.* 476: 313-329.
- Jimenze, L., Martinez, C., Perez, I., Lopez, F. (1997). Biobleaching procedures for pulp from agricultural residues using *Phanerochaete chrysosporium* and enzyme. *Process Biochem.* 32(4):297-304.
- Kang, S.W., Kim, S.W. and Lee, J.S. (1995). Production of cellulase and xylanase in a bubble column using immobilized *Aspregillus* sp. *Appl. Biochem Biotechnol.* 53:101-106.

- Karthikeyan, R.S., Rakshit, S.K. and Baradarajan, A.(1997). Optimization of batch fermentation conditions for dextran production. *Bioproc. Engg.* 15(5):247-252.
- Keneth W. B. (1970). Hand book of pulp and paper technology second edition revised and enlarged. Van Nostrand Reinhold Company. New York. Pp. 269-425.
- Khanna, P., Sundari, S.S. and Kumar, N.J. (1995). Production, Isolation and partial purification of xylanase from and *Aspergillus* sp. *World J. Microbiol. Biotechnol.* 11:242-243.
- Khanongnuch, C., Asada, K., Tsurua, H., Ooi, T., Kinoshita, S. and Llummyoung, S. (1998). Beta-mannanase and xylanase of *Bacillus subtilis* 5H active for biobleaching of curd pulp. *J. Ferm. Bioeng.* 86:461-466.
- Kitamoto, N., Yoshino, S., Ohmiya, K. and Tsukagoshi, N. (1999). Sequence analysis, overexpression, and antisense inhibit of a β -xylosidase gene, *xyl A*, from *Aspergillus oryzae* KBN616. *Appl. Environ. Microbiol.* 65:20-24.
- Kluefel, D., Vats-Mehta, S., Aumont, F., Shareck, F. and Morosoli R.(1990). Purification and characterization of a new xylanase (xylanase B) produced by *Streptomyces lividans* 66. *J.Biochem.* 267:45-50.
- Kohli, U., Nigam P., Singh D. and K. Chaudhary (2001). Thermostable, alkalophilic and cellulase-free xylanase production by *Thermoactinomyces thalophilus* subgroup C. *Enzyme. Microb. Technol.*, 28:606-610.
- Kulkarni, N. and Rao, M.(1996). Application of xylanase from alkalophilic thermophilic *Bacilus* sp.NCIM59 in biobleaching of bagasse pulp. *J. Biotechnol.* 51:167-173.
- Ledoux, P., Detroz, R., deBuyl, E., Shetty, J., Troughton, N. and Presly, J.R. (1993). Use of bacterial xylanase in chlorine free bleaching sequences. Tappi Proceedings of the Pulping Conference. Boston. Vol.3 pp. 1057-065.

- Li, K., Azadi, P., Collins, R., Tolan, J., Kim, J.S. and Eriksson, K.E.L. (2000). Relationship between activity of xylanases and xylan structure. *Enz. Microb. Technol.* 27:89-94.
- Lopez-Fernandez, C.L., Rodriguez, J., Ball, A.S., Copa-Patino, J.L., Perez-Leblic, M.I. and Arias, M.E.(1998). Application of the affinity binding of xylanases to oat-splet xylan in purification of endoxylanase CM-2 from *Streptomyces chattanoogensis* CECT3336. *Appl. Microbiol. Biotechnol.* 50(2):284-287.
- Lundgren, K.R., Bregkvist, L., Hogman, S., Joves, H., Erickson, G., Bartfai, T., Laan, J., Rosenberg, E. and Shonam, Y.(1994). TCF mill trial on soft wood pulp with Korsnas thermostable and alkaline stable xylanase T6. *FEMS Microbiol. Rev.*13:365-.68.
- Magnuson, T.S. and Crawford, D.L.(1997). Purification and characterization of an alkaline xylanase from *Streptomyces viridosporus* T7A. *Enz. Microbiol. Technol.* 21(3):160-164.
- Maheswari, M.U. and Chandra, T.S..(2000). Production and potential applications of xylanase from a new strain of *Streptomyces cuspidosporus*. *World J. Microbiol. Biotechnol.*16:257-263.
- Manmori, T., Watanabe, T., Shinke, R., Kohno, A. and Kawamera, Y.(1990). Purification and properties of thermostable xylanase and β -xylosidase production by newly isolated *Bacillus stearothermophilus* strain. *J. Bacteriol.*12:6669-6672.
- Miller, L.H. (1959). Use of dinitrosalilic acid reagent for determination of reducing sugars. *Anal. Chem.* 31:426-428.
- Montgomery, D.C.(1991) Design and analysis of experiments, 3rd ed., Wiley, New York
- Morosoli, R.M., Bertrand, J., Mondou, F., Shareck, F and Kueppel, D. (1986). Purification and properties of a xylanase from *Streptomyces lividans*. *J. Biochem.* 239:587-592.

- Murthy, M.S.R.C., Swaminathan, T., Rakshit, S.K. and Kosugi, Y.(2000). Statistical optimization of lipase catalyzed hydrolysis of methyloleate by response surface methodology. *Bioprocess Engineering*. 22:35-39.
- Nammori, T., Watanabe, T., Shinke, R., Kohno, A. and Kayamura, Y.(1990). Purification and properties of thermostable xylanase and β -xylosidase produced by a newly isolated *Bacillus stearothermophilus* strain. *J. Bacteriol.*172(12):6669-6672.
- Nath, D and Rao, M. (2000). pH dependent conformational and structural change of xylanase from an alkalophilic thermophilic *Bacillus* sp. NCIM59. *Enz. Microbiol. Technol.* 28:397-403.
- Okeke, B.C. and Obi, S.K.C..(1993). Production of cellulolytic and xylanolytic enzymes by an *Antrographis* species. *World J. Microbiol. Biotechnol.* 9:345-349.
- Panda, T., Babu, A., Kumari, J., Rao, D.S., Theodore, K., Rao, K.J., Kesava, K.S., Nair, S.R., Sinha, J., Srinivas, R., Prasanna, G.L., Das, V.V., Pazouki, M., Felse, P.A., Naidu, G.S., Gokul, B., Uma, S., Srividya, K., Muralidhar, R., Balamurugan, K. and Chandra Sekar, R.(1997). Bioprocess optimization a challenge. *J. Microbiol. Biotechnol.* 7:367-372.
- Poutanen, K. and Puls, J..(1998). Characteristics of *Trichoderma reesei* β -xylosidase and its use in the hydrolysis of solubilized xylans. *Appl. Microbiol. Biotechnol.* 28:425-432.
- Poutanen, K., Ratto, M., Puls, J. and Viikari, L. (1987). Evaluation of different microbial xylanolytic systems. *J. Biotechnol.* 6:49-60.
- Prasertsan, P., H-Kittikuk, A., Kunghae, A., Maneesri, J. and Oi, S. (1997). Optimization for xylanase and cellulase production from *Aspergillus niger* ATTC6275. In palm oil mill waste and its application. *World J. Microbiol. Biotechnol.* 13 (5): 555-559.

- Ragauskas, A.J., Poll, K.N. and Cesternino, A. (1994). Effects of xylanase pretreatment procedures on non chlorine bleaching. *Enz. Microbiol. Technol.* 16:492-5.
- Rama Mohan Reddy, P., Gopal Reddy and Seenayya, G. (1999). Production of thermostable pullulanase by *Clostridium thermosulfurogenes* SV2 in solid-state fermentation: optimization of nutrients levels using response surface methodology. *Bioprocess Engineering.* 21:497-503.
- Rani, D.S. and Nand, K. (2000). Production of thermostable cellulase-free xylanase by *Clostridium absonum* CFR-702. *Process Biochem.* 36:355-362.
- Reddy, V., Reddy, P., Pillay, B. and Singh, S. (2002). Effect of aeration on the production of hemicellulase by *T. lanuginosus* SSPB in a 30 l bioreactor. *Process Biochem.* 37:1221-1228.
- Sachslehner, A., Nidetzky, B., Kulbe, K.D. and Hartrich, D. (1998). Induction of mannanase, xylanase and endoglucanase activity in *Sclerotium rolfsii*. *Appl. Environ. Microbiol.* 64(3): 594-600.
- Sa-Pereira, P., Costa-Ferreira, M. and Atres-Barros, M.R. (2002a). Enzymatic properties of a neutral endo-1,3(4)- β -xylanase Xyl II from *Bacillus subtilis*. *J. Biotechnol.* 58:71-78.
- Sa-Pereira, P., Mesquita, A., Duarte, J.C., Barros, M.R.C. and Costa-Ferreira, M. (2002b). Rapid production of thermostable cellulase-free xylanase by strain of *Bacillus subtilis* and its properties. *Enz. Microbiol. Technol.* 30: 924-933.
- Saxena, S., Bahadur, J. and Varna, A. (1991). Production and localization of carboxymethylcellulase xylanase and β -glucosidase from *Cellulomonas* and *Micrococcus* sp. *Appl. Microbiol. Biotechnol.* 34:668-670.

Saxena, S., Fierobe, H.P., Gaudin, C., Guerlesquin, F. and Belaich, J.P. (1995).

Biochemical properties of a β -xylosidase from *Clostridium cellulolyticum*. *Appl. Environ. Microbiol.* 61:3509-3512.

Sen, R.K. and Swaminathan, T.(1997). Application of response surface methodology to evaluate the optimum environmental conditions for enhanced production of surfactin. *Appl. Microbiol. Biotechnol.* 47:358-363.

Shah, A.K., Sidid, S.S., Ahmad, A. and Rele, M.V. (1999). Treatment of bagasse pulp with cellulase-free xylanase from an alkalophilic *Bacillus* sp.Sam-3. *Bioresource Technol.*68:133-140.

Singh, S., du Peez, J.C., Pillay, B. and Prior, B.A. (2000a). The production of hemicellulases by *Thermomyces lanuginosus* strain SSBP: influent of agitation and dissolved oxygen tension. *Appl. Microbiol. Biotechnol.* 54:698-704.

Singh, S., Pillay, B. and Prior, B.A. (2000b). Thermal stability of β -xylanase produced by different *Thermomyces lanuginosus* strain. *Enz. Microbiol. Technol.*26:502-508.

Singh, S., Reddy, P., Haarhoff, J., Biely, P., Janse, B., Pillay, B., Pillay, D and Prior, B.A. (2000c). Relatedness of *Thermomyces lanuginosus* strain producing thermostable xylanase. *J.Biotechnol.*81:11-128.

Smits, J.P., Rinzema, A., Tramper, J., VanSonsbeek, H.M. and Knol, W. (1996). Solid-state fermentation of wheat bran by *Trichoderma reesei* QM9414: substrate composition changes, C balance, enzyme production, growth and kinetics. *Appl. Microbiol. Biotechnol.* 46:486-496.


Souza, M. C. de O., Roberto, I. C. and Milagres, A. M. F.(1999). Solid-state fermentation for xylanase production by *Thermoascus aurantiacus* using response surface methodology. *Appl. Microbiol. Biotechnol.* 52:768-772.

- Srinivasan, M., Natarajan, K. and Nagarajan, G.. (2000). Growth optimization of an ectomycorrhizal fungus with respect to pH and temperature in vitro, using Design of Experiments. *Bioprocess Engineering*, 22: 267-273.
- Steiner, W., Lafferty, R.M., Gomes, I. and Esterbauer, H.(1987). Studies on wild strain of *Schizophyllum commune*: cellulase and xylanase production and formation of extracellular *Schizophyllum*. *Biotechnol Bioeng*. 30:169-178.
- Sunna, A. and Antranikian, G. (1997). Xylanolytic enzymes from fungi and bacteria. *Crit. Rev. biotech*.17:39-67.
- Techapun C., Poosaran, N., Watanabe, M., and Sasaki, K. (2003). Thermostable and alkaline-tolerant microbial cellulase-free xylanases produced from agricultural wastes and the properties required for use in pulp bleaching bioprocesses: a review. *Process Biochemistry* 1 –14 (In press).
- Techapun C., Sinsuwongwat, S., Watanabe, M., Poosaran, N. and Sasaki, K. (2001). Feasibility of cellulase-free xylanase production in agricultural waste materials by thermotolerant microorganisms, *Biotechnol. Lett*. 23:1685-1698
- Tenkanen, M. and Siika-aho, M. (2000). An α -glucuronidase of *Schizophyllum commune* acting of polymeric xylan. *J. Biotechnol*. 78:149-161.
- Teruel, A.M.L., Gontier, E., Bienaime, C., Nava, J.E., Saucedo, and Barbotin, J.N. . (1997). Response surface analysis of chlortetracycline and tetracycline production with K. carrageenan immobilized *Streptomyces aureofaciens*. *Enz. Microbiol. Technol*. 21:314-320.
- Tseng, M., Yap, M., Ratanakanokchai, K., Kyu, K.L. and Chen, S.(2002). Purification and characterization of two cellulase-free xylanase from an alkalophilic *Bacillus firmus*. *Enz. Microbiol. Technol*. 30:590-595.

- Vicuna, R., Oyarzum, E. and Osses, M.(1995). Assessment of various commercial enzymes in bleaching of radiata pine kraft pulp. *J. Biotechnol.* 40:163-168.
- Viikari, L., Kantelinen, A., Sundquist, J. and Ranua, M. (1990). Characterization of pulps treated with hemicellulolytic enzymes period to bleaching. *Biotechnology in Pulp and Paper Manufacture*. Boston : Butterworth-Heinemann.pp.145-151.
- Viikari, L., Kantelinen, A., Sundquist, J. and Linko, M. (1994). Xylanases in bleaching: from an idea to industry. *FEMS Microbiol.Rev.*13:335-350.
- Vijayalakshmi, G., Shobha, B., Vanajakshi, V., Divakar, S. and Manohar, B.(2001). Response surface methodology for optimization of growth parameters for the production of carotenoids by a mutant strain of *Rhodotorula gracilis* . *Eur. Food Res. Technol.* 213:234–239
- Winkler, M.(1998). Optimization and time-profing in fermentation processes. In: M.E. Bushell, (Ed.) *Progress in industrial microbiology*. vol. 25, Elsevier, Amsterdam. pp. 91-150.
- Wong, K.K.Y., Tan, L.U.L. and Saddler, J.N. (1998). Multiplicity of β -1,4-Xylanases in microorganisms: function and applications. *Microbiological. Rev.* 52:305-317.
- Wong, K.Y., Nleson, S.L. and Saddler, J.N.(1996). Xylanase treatment for the peroxide bleaching of oxygen delignified kraft pulp derived form three soft wood species. *J. Biotechnol.* 48.137-145.
- Yang, V.W., Zhang, Z., Elegir, G. and Jiffries, T.W.(1995). Alkaline-active xylanase produced by alkalophilic *Bacillus* sp. isolated from kraft pulp.J. *Ind. Microbiol.* 15:434-443.
- Yu, X., Hallett, S.G., Sheppard, J. and Watson, A.K..(1997). Application of the Plackett-Burman experimental design to evaluate nutritional requirements for the production of *Colletotrichum coccodes* spores. *Appl. Microbiol. Biotechnol.* 47:301-305.

Zheng, L., Do, J.H. and Zhang, J. (2000). Biobleaching effect of xylanase preparation from an alkalophilic *Bacillus* sp. On ramie fiber. *Biotechnol. Lett.*22(17) : 1363-1367.

Zhengqiang, J., Kobayashi, A., Ahson, M.M., Lite, L., Kitaoka, M. and Hayashi, K. (2001). Characterization of a thermostable family-10 endo-xylanase (XynB) from *Thermotoga maritima* that cleaves *p*-nitrophenyl- β -D-xylosidase. *J. Biosci. Bioeng.* 92(5):423-428.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่

Copyright© by Chiang Mai University

All rights reserved