

## REFERENCES

- Augustinius, P.C., and Macphail, M.K. 1997. Early Pleistocene stratigraphy and timing of the Bulgobac Glaciation, western Tasmania, Australia. *Palaeogeography, palaeoclimatology, palaeoecology*, 128: 253-267.
- Awasthi, N., and Ahuja, M. 1982. Investigation of some carbonised woods from the Neogene of Varkala in Kerala Coast. *Geophytology*, 12(2): 245-259.
- Awasthi, N., Mehrotra, R.C., and Srivastava, R. 1996. Fossil woods from the Deccan Intertrappean beds of Madhya Pradesh. *Geophytology*, 25: 113-118.
- Awasthi, N., and Srivastava, R. 1992. Additions to the Neogene flora of Kerala Coast, India. *Geophytology*, 20(2): 148-154.
- Bande, M.B. 1992. Paleogene vegetation of peninsular India (megafossil evidences). *Palaeobotanist*, 40: 275-284.
- Bande, M.B., and Prakash, U. 1980. Fossil woods from the Tertiary of West Bengal, India. *Geophytology*, 10(1-2): 146-157.
- Baksi, S.K. 1962. Palynological investigation of Simsang River Tertiaries, South Shillong Front, Assam. *Bulletin of the Geological, Mining and Metallurgical Society of India*, 26: 1-22.
- Barr, S.M., Macdonald, A.S., Haile, N.S., and Reynolds, P.H. 1976. Paleomagnetism and age of the Lampang basalt (northern Thailand), and age of underlying pebble tools. *Journal of the Geological Society of Thailand*, 3(1-2): 1-10.
- Barron, E.J. 1987. Cretaceous plate tectonic reconstructions. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 59: 3-29.

- Baum, F., Braun, E.v., Hahn, L., Hess, A., Koch, K.-E., Kruse, G., Quarch, H., and Siebenhüner, M. 1970. On the geology of northern Thailand. *Beihefte zum Geologischen Jahrbuch*, 102: 1-24.
- Baum, F., Braun, E.v., Hess, A., and Koch, K.E. 1982. *Geological map of northern Thailand, Sheet Chiang Mai (5), scale 1: 250,000*. German Geological Mission in Thailand in cooperation with Geological Survey Division Department of Mineral Resources Thailand, Federal Institute for Geosciences and Natural Resources, Stuttgart.
- Benammi, M., Urrutia-Fucugauchi, J., Alva-Valdivia, L.M., Chaimanee, Y., Triamwichanon, S., and Jaeger, J.-J. 2002. Magnetostratigraphy of the Middle Miocene continental sedimentary sequences of the Mae Moh basin in northern Thailand: evidence for counterclockwise block rotation. *Earth and Planetary Science Letters*, 204: 373-383.
- Benammi, M., Urrutia-Fucugauchi, J., Chaimanee, Y., and Jaeger, J.-J. (2003). Preliminary paleomagnetic study of the Chiang Muan basin in northern Thailand. in B. Ratanasthien, S.L. Rieb, and S. Chantraprasert (Eds.), *Proceeding of the 8<sup>th</sup> International Congress on Pacific Neogene Stratigraphy: Pacific Neogene Paleoenvironments and Their Evolution* (pp. 352), Chiang Mai, Thailand.
- Biswas, S. 1962. Stratigraphy of the Mahodeo, Langpur, Cherra, and Tura Formations, Assam, India. *Bulletin of Geological, Mineralogical and Metallurgical Society of India*, 25: 1-48.
- Bos, J.A.A., and Punt, W. 1991. The Northwest European pollen flora, 47: Juglandaceae. *Review of Palaeobotany and Palynology*, 69(1-3): 79-95.
- Bradley, W.H. 1967. Two aquatic (Chytridiales) of Eocene age from the Green River Formation of Wyoming. *American Journal of Botany*, 54(5): 577-582.

- Brown, G.F., Buravas, S., Charaljavanaphet, J., Jalichandra, N., Johnson, W.D., Sresthaputra, V., and Taylor, G.C. 1951. Geologic reconnaissance of the mineral deposits of Thailand. *Geologic Investigation of Asia. U.S. Geological Survey Bulletin*, 984: 1-183.
- Buffetaut, E., Helmcke-Ingavat, R., Jaeger, J.J., Jongkanjanasootorn, Y., Suteethorn, S., and Tong, H. (1989). Fossil vertebrates and the age of the intermontane basins of Thailand. in: T. Thanasuthipitak and P. Ounchanum (Eds.), *Proceedings of the International Symposium on Intermontane Basins: Geology and Resources* (pp. 187-195), Chiang Mai, Thailand.
- Chaimanee, Y., Jolly, D., Benammi, M., Tafforean, P., Duzer, D., Moussa, I., and Jaeger, J.J., 2003. A Middle Miocene hominoid from Thailand and orangutan origin. *Nature*, 422: 61-65.
- Chaodumrong, P. (1985). *Sedimentological studies of some Tertiary deposits of Mae Moh basin, Changwat Lampang*. Master's thesis. Chulalongkorn University, Thailand.
- Chaodumrong, P., and Burrett, C.F. 1997. Stratigraphy of the Lampang Group in central North Thailand: new version. *CCOP Technical Bulletin* 26: 65-80.
- Chaodumrong, P., Snansieng, S., Bhoripatkosol, S., and Khamdhurian, W. (1982). "Geology of Li basin", Geological Survey Division, Department of Mineral Resources, Bangkok, Report No. 4.
- Charoenprawat, A., Chuaviroj, S., Hinthong, C., and Chonglakmani, C. (1995). "Geological map of Thailand, Sheet Lampang (NE47-7), scale 1: 250,000", Geological Survey Division, Department of Mineral Resources, Bangkok, Thailand.

- Cheneval, J., Ginsburg, L., Mourer-Chauvire, C., and Ratanasthien, B. 1991. The Miocene avifauna of the Li Mae Long locality, Thailand: systematics and paleoecology. *Journal of Southeast Asian Earth Sciences*, 6(2): 117-126.
- Chiji, M., and IGCP-246 National Working Group of Japan. 1990. Paleogeography and environments during the Opening of the Sea of Japan. in R. Tsuchi (Ed.) *Pacific Neogene Events, Their Timing, Nature and Interrelationship* (pp. 161-169), University of Tokyo Press.
- Chowdhury, K.A., and Tandon, K.N. 1964. A fossil wood of *Terminalia tomentosa* W. & A. from the Tertiary of Burma. *Annals of Botany*, 28(111): 445-450.
- Cookson, I.C. 1947. On fossil leaves (Oleaceae) and a new type of fossil pollen grain from Australian brown coal deposits. *Proceedings of the Linnean Society of New South Wales*, 72: 183-197.
- Cookson, I.C. (1957). *Plant microfossils from the lignites of Kerguelen Archipelago British and New Zealand Antarctic Research Expedition 1929-1931*. Reports-Series A, 2(8): 129-142.
- Couper, R.A. 1953. Upper Mesozoic and Cainozoic spores and pollen grains from New Zealand. *New Zealand Geological Survey, Palaeontological Bulletin*, 22: 1-77.
- Couper, R.A. 1960. New Zealand Mesozoic and Cainozoic plant microfossils. *New Zealand Geological Survey, Palaeontological Bulletin*, 32: 1-87.
- Cranwell, L.M. (1964). Antarctica: cradle or grave for its *Nothofagus*? in: L.M. Cranwell (Ed.), *Ancient Pacific Floras* (pp. 87-93), University of Hawaii Press, Honolulu.

- David, C.A. 1916. On the fossil algae of the petroleum-yielding shales of the Green River Formation of Colorado and Utah. *U.S. National Academic Science Proceedings*, 11: 114-119.
- Dettmann, M.E. 1963. Upper Mesozoic microfloras from southeastern Australia. *Proceedings of the Royal Society of Victoria*, 77: 1-148.
- Dettmann, M.E. 1973. Angiospermous pollen from Albian to Turonian sediments of eastern Australia. *Special Publication, Geological Survey of Australia*, 4: 3-34.
- Dilcher, D.L. 1965. Epiphyllous fungi from Eocene deposits in western Tennessee, U.S.A. *Palaeontographica Abt.B.*, 116: 10-13.
- Ducrocq, S., Buffetaut, E., Chaimanee, Y., Suteethorn, V., and Jaeger, J.J. 1995. Mammalian faunas and the ages of the continental Tertiary fossiliferous localities from Thailand. *Journal of Southeast Asian Earth Science*, 12(1-2): 65-78.
- Ducrocq, S., Chaimanee, Y., Suteethorn, V., and Jaeger, J.J. 1994. Ages and paleoenvironment of Miocene mammalian faunas from Thailand. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 108(1-2): 149-163.
- Dueñas, H. 1980. Some remarks about the genus *Magnastriatites* Germeraad, Hopping & Muller, 1968. *Review of Palaeobotany and Palynology*, 30: 329-331.
- Ediger, V.S., and Bati, Z. 1988. Morphological examination of *Pediastrum* (Chlorophyta) from the Tertiary strata of the Thrace basin (NW Turkey). *Pollen et Spores*, 30(2): 203-222.
- Edwards, W.N. 1922. An Eocene microthyriaceous fungus from Mull, Scotland. *Transactions British Mycology Society*, 8: 66-72.
- Elsik, W.C. 1968. Palynology of a Palaeocene Rockdale lignite, Milam County, Texas, I: morphology and taxonomy. *Pollen et Spores*, 10(2): 263-314, II: morphology and taxonomy. *Pollen et Spores*, 10(3): 599-664.

- Elsik, W.C., and Dilcher, D.L. 1974. Palynology and age of clays exposed in Lawrence Clay Pit, Henry County, Tennessee. *Palaeontographica Abt.B.*, 146: 65-87.
- Endo, S. 1963. Some older Tertiary plants from northern Thailand. *Japanese Journal of Geology and Geography*, XXXIV (2-4): 177-180.
- Endo, S. (1964). Some older Tertiary plants from northern Thailand. in: T. Kobayashi (Ed.), *Geology and Palaeontology of Southeast Asia*, 1 (pp. 113-117), The University of Tokyo Press, Tokyo.
- Endo, S. (1966). A supplementary note on the Paleogene Li flora in northern Thailand. in: T. Kobayashi and R. Toriyama (Eds.), *Geology and Palaeontology of Southeast Asia*, 3 (pp. 165-169), The University of Tokyo Press, Tokyo.
- Engelhardt, D.W. 1964. Plant microfossils from the Eocene Cockfield Formation, Hinds County, Mississippi, *Mississippi Geological Research Paper*, 104: 65-95.
- Erdtman, G. (1943). *An introduction to pollen analysis*. Waltham, Mass., U.S.A.
- Ethridge Glass, D.L., Brown, D.D., and Elsik, W.C. 1986. Fungal spores from the Upper Eocene Manning Formation, Jackson Group, East and South-Central Texas, U.S.A. *Pollen et Spores*, 28(3-4): 403-420.
- Evitt, W.R. 1963. Occurrence of freshwater alga *Pediastrum* in Cretaceous marine sediments. *American Journal of Science*, 261: 890-893.
- Faegri, K., and Iversen, J. 1989. *Textbook of pollen analysis*. IV edition, John Wiley & Sons.
- Fairchild, W.W., and Elsik, W.C. 1969. Characteristic palynomorphs of the Lower Tertiary in the Gulf Coast. *Palaeontographica*, Abt.B, 128: 81-89.



- Farabee, M.J., and Canright, J.E. 1986. Stratigraphic palynology of the lower part of the Lance Formation (Maastrichtian) of Wyoming. *Palaeontographica, Abt.B.*, 199: 1-89.
- Filatoff, J. 1975. Jurassic palynology of the Perth Basin, western Australia. *Palaeontographica Abt.B.*, 154: 1-113.
- Frederiksen, N.O. 1980a. Sporomorphs from the Jackson Group (Upper Eocene) and adjacent strata of Mississippi and western Alabama. *USGS Professional Paper*, 1084: 1-75.
- Frederiksen, N.O. 1980b. Paleogene sporomorphs from South Carolina and quantitative correlations with the Gulf Coast. *Palynology*, 4: 125-179.
- Frederiksen, N.O., and Ames, H.T. 1979. Status of the pollen genus *Pollenites*. *Taxon*, 28(5/6): 227-561.
- Frederiksen, N.O., and Christopher, R.A. 1978. Taxonomy and biostratigraphy of Late Cretaceous and Paleogene triatriate pollen from South Carolina. *Palynology*, 2: 113-145.
- Gardner, S., Sidisunthorn, P., and Anusarnsunthorn, V. 2000. A field guide to forest trees of northern Thailand. *CMU Herbarium, Biology Department, Science Faculty, Chiang Mai University*: 545 p.
- Germeraad, J.H., Hopping, C.A., and Muller, J. 1968. Palynology of Tertiary sediments from tropical areas. *Review of Palaeobotany and Palynology*, 6(3/4): 189-348.
- Ghosh, A.K., Jacob, K., and Lukose, N.G. 1964. On the spores of Parkeriaceae and Schizaeaceae from India. *Bulletin of Botany Society of Bengal*, 17(1&2): 23-28.

- Gibling, M., and Ratanasthien, B. 1980. Cenozoic basins of Thailand and their coal deposits: a preliminary report. *Bulletin of Geological Society of Malaysia*, 13: 27-42.
- Ginsburg, L., and Tassy, P. 1985. The fossil mammals and the age of the lignite beds in the intermontane basins of northern Thailand. *Journal of Geological Society of Thailand*, 8(1-2): 13-27.
- Ginsburg, L., and Ukkakimapan, Y. 1983. Uncervidé nouveau du Miocène d'Asie et l'âge des lignites des bassins intramontagneux du Nord de la Thaïlande. *C.R. Acad. Sci., Paris*: 297-300.
- Gonzalez-Guzman, E. 1967. A palynological study on the Upper Los Cuervos and Mirador Formations (Lower and Middle Miocene; Tibú Area, Colombia). *Brill, Leiden*: 129 pp.
- Groot, J.J., and Groot, C.R. 1962. Some plant microfossils from the Brightseat Formation (Paleocene) of Maryland. *Palaeontographica*, B111: 161-171
- Gortemaker, R.E. 1986. A method to identify pollen of some recent and fossil species of *Fagus* L. (Fagaceae). *Review of Palaeobotany and Palynology*, 47: 263-292.
- Guan, X., Fan, H., Song, Z., and Zheng, Y. 1989. *Researches on Late Cenozoic palynology of the Bohai Sea*. Nanjing University Press.
- Guleria, J.S. 1994. Fossil leaf of *Terminalia* from the Rajpardi lignite mine, District Bharuch, Gujarat, India. *Geophytology*, 24(1): 55-58.
- Gupta, K.M. 1936. *Leguminoxylon burmense* gen. and sp. nov., a dicotyledonous wood from the Tertiary of Burma. *Indian Science Congress Proceedings*, 305 p.
- Guy-Ohlson, D. 1992. *Botryococcus* as an aid in the interpretation of palaeoenvironment and depositional processes. *Review of Palaeobotany and Palynology*, 71: 1-15.



- Hart, G.F. (1969). Palynology of Permian period. in R.H. Tschudy and R.A. Scott (Eds), *Aspect of Palynology* (pp. 271-289), John Wiley and Sons: New York.
- Hekel, H. 1972. Pollen and spore assemblages from Queensland Tertiary sediments. *Geological Survey of Queensland, publication 355*: 1-34.
- Hughes, N.F. (1969). Jurassic and Early Cretaceous pollen and spores. In: R.H. Tschudy and R.A. Scott (eds.), *Aspects of Palynology* (pp. 311-329), John Wiley and Sons, New York.
- Itoigawa, J., and Yamanoi, T. (1990). Climatic optimum in the Mid-Neogene of the Japanese Islands. in: R. Tsuchi (Ed.) *Pacific Neogene Events, Their Timing, Nature and Interrelationship* (pp. 1-14), University of Tokyo Press.
- Jaeger, J.J., Tong, H., Buffetaut, E., and Ingavat, R. 1985. The first fossil rodents from the Miocene of northern Thailand and their bearing on the problem of the origin of the Muridae. *Revue de Paléobiologie*, 4(1): 1-7.
- Jitapankul, S. (1992). Geology of Ban Pa Kha coal deposits, Li basin, Lamphun. in: C. Piancharoen (Ed.), *Proceedings of the National Conference on Geologic Resources of Thailand: Potential for Future Development* (pp. 259-272), Bangkok, Thailand.
- Jolivet, M., Brunel, M., Seward, D., Xu, Z., Yang, J., Roger, F., Tapponnier, P., Malavieille, J., Arnaud, N., and Wu, C. 2001. Mesozoic and Cenozoic tectonics of the northern edge of the Tibetan plateau: fission track constraints. *Tectonophysics*, 343: 111-134.
- Kar, R.K. 1979. Palynological fossils from the Oligocene sediments and their biostratigraphy in the District of Kutch, western India. *The Palaeobotanist*, 26(1): 16-49.

- Kar, R.K., and Ambwani, K. 1991. Light-microscopy and SEM studies of *Striatriletes* and *Malayaeaspora* from India and Malaysia. *Geophytology*, 21: 45-47.
- Kedves, M., and Solé de Porta, N. 1963. Comparacion de las esporas del género *Cicatricosisporites* R. Potonié y Gell, 1933 de Hungría y Colombia. Algunos problemas referentes a su significado estratigráfico. *Bot. Geol.*, 12: 51-76.
- Kemp, E.M., and Harris, W.K. 1977. The palynology of Early Tertiary sediments Ninetyeast Ridge, Indian Ocean. *Special Paper in Palaeontology*, 19: 1-70.
- Khan, A.M., and Martin, A.R.H. 1971. A note on genus *Polypodiisporites* R. Potonié. *Pollen et Spores*, 13: 475-480.
- Klaus, W. 1954. Bau und form von *Sporotrapoidites illingensis* n. gen et sp. sporomorpharum. *Grana palynologica*, 1: 114-131.
- Krutzsch, W. 1959. Mikropaläontologische (sporen-palaontologische) untersuchungen in der Braunkohle des Geiseltales. *Geologie*, 8: 1-425.
- Krutzsch, W. 1960. Über *Thomsonipollis magnificus* (Th & Pf, 1953) n. fgen. N. comb. und Bemerkungen zur regionalen Verbreitung einiger Pollengruppen im älteren Paläogen. *Freiberger Forsch.*, C86: 53-65.
- Krutzsch, W. 1962. *Atlas der mittel-und jungtertiären dispersen sporen-und pollen-sowie der mikroplanktonformen des nördlichen Mitteleuropas*. Leif. I.-VEB Deutscher Verlag der Wissenschaften, Berlin: 108 p.
- Krutzsch, W. 1963. Atlas der mittel-und jungtertiären dispersen sporen-und pollen-sowie der mikroplankton- formen des nördlichen mitteleuropas. Lieferung 2: Die sporen der Anthocerotaceae und der Lycopodiaceae. *Veb Gustav Fischer Verlag Jena, Berlin*.
- Krutzsch, W. 1969. Über eigene stratigraphisch wichtige neue longaxones. Pollen aus dem mitteleuropäischen Alttertiär. *Geologie*, 11: 472-487.

- Kuprianova, L.A. 1960. Palynological data contributing to the history of *Liquidambar*. *Pollen et Spores*, 2(1): 71-88.
- Lange, R.T. 1976. Fossil ephiphyllous « germlings », their living equivalents and their palaeohabitat indicator value. *Neues Jahrbuch für Geologie un Palaeontologie, Abhandlungen*, 151(2): 142-165.
- Lee, T.-Y., and Lawver, L.A. 1995. Cenozoic plate reconstruction of Southeast Asia. *Tectonophysics*, 251: 85-138.
- Leffingwell, H.A. 1971. Palynology of the Lance (Late Cretaceous) and Fort Union (Paleocene) Formations of the type Lance area, Wyoming. *Geological Society of America Special Paper*, 127: 1-64.
- Leidelmeyer, P. 1966. The Paleocene to Lower Eocene pollen flora of British Guinea. *Ledise Geol. Meded.*, 38: 49-57.
- Leloup, P.H., Lacassin, R., Tapponnier, P., Schärer, U., Dalai, Z., Xiaohan, L., Liangshang, Z., Shaocheng, J., and Trinh, P.T. 1995. The Ailao Shan – Red River shear zone (Yunnan, China), Tertiary transform boundary of Indochina. *Tectonophysics*, 251: 3-84.
- Le Pichon, X., Fournier, M., and Jolivet, L. 1992. Kinematics, topography, shortening, and extrusion in the India-Eurasia collision. *Tectonics*, 11(6): 1085-1098.
- Mandaokar, B. 1993. A palynological investigation of the Tikak Parbat Formation (Oligocene) of Dangri Kumari colliery, Dibrugarh District, Assam, India. *Tertiary Research*, 14(4): 127-139.
- Mandaokar, B. 2000. Palynology of coal-bearing sediments of the Tikak Parbat Formation (Oligocene) from Namchik River section, Changlang District, Arunachal Pradesh, India. *Tertiary Research*, 20(1-4): 37-46.

- Manum, S. 1962. Studies in the Tertiary flora of Spitsbergen, with notes on Tertiary floras of Ellesmere Island, Greenland and Iceland. *Norsk Polarinst. Skrifter*, 125: 1-127.
- Martin, H.A., Macphail, M.K., and Partridge, A.D. 1996. Tertiary *Alangium* (Alangiaceae) in eastern Australia: evidence from pollen. *Review of Palaeobotany and Palynology*, 94(1-2): 111-122.
- Martin, H.A., and Rouse, G.E. 1966. Palynology of Late Tertiary sediments from Queen Charlotte Islands, British Columbia. *Canadian Journal of Botany*, 44: 171-208.
- Matsuoka, K. 1990. The Early Middle Miocene inland paleoclimate around the Central Kinki, Southwest Japan. In R. Tsuchi (ed.) *Pacific Neogene Events, Their Timing, Nature and Interrelationship* (pp. 15-22), University of Tokyo Press.
- Mauzy, G., Muller, J., and Lugardon, B. 1975. Notes on the morphology and fine structure of the exine of some pollen types in Dipterocarpaceae. *Review of Palaeobotany and Palynology*, 19(4): 241-289.
- Meesuk, J. (1986). *Geology of the Tertiary coal basins of Thailand*. Master's thesis, University of Aston in Birmingham, England.
- Mehrotra, R.C. 1988. Fossil wood of *Sonneratia* from the Deccan Intertrappean beds of Mandla District, Madhya Pradesh. *Geophytology*, 18(2): 129-134.
- Mein, P., Ginsburg, L., and Ratanasthien, B. 1990. Nouveaux rongeurs du Miocène de Li (Thaïlande). *Compte Rendu Acad. Sci. Paris*, 310(II): 861-865.
- Mohr, B.A.R. 1983. Nachweis von Pollen der Gattung *Hemitrapa* Miki (Trapaceae) aus mittelmiozänen Schichten von Gallenbach bei Dasing (Lkrs. Aichach-Friedberg). *Berichte Naturwissenschaften Vereins für Schwaben*, 87: 69-72.

- Molnar, P., and Burke, P. 1977. Penrose conference report: Erik Norin Penrose conference on Tibet. *Geology*, 5: 161-163.
- Molnar, P., and Tapponnier, P. 1975. Cenozoic tectonics of Asia: Effect of the continental collision. *Science*, 189: 419-426.
- Morley, C.K., Sangkumarn, N., Hoon, T.B., Chonglakmani, C., and Lambiase, J. 2000. Structural evolution of the Li Basin, northern Thailand. *Journal of the Geological Society, London*, 157: 483-492.
- Morley, C.K., Woganan, N., Sankumarn, N., Hoon, T.B., Aliet, A., and Simmons, M. 2001. Late Oligocene – Recent stress evolution in rift basins of northern and central Thailand: implications of escape tectonics. *Tectonophysics*, 334: 115-150.
- Morley, R.J. 1982. Fossil pollen attributable to *Alangium* Lamarck (Alangiaceae) from the Tertiary of Malesia. *Review of Palaeobotany and Palynology*, 36: 65-94.
- Morley, R.J. (2000). *Origin and evolution of tropical rainforests*. John Wiley & Sons.
- Mukherjee, A. 1942. A gymnospermous fossil wood from Burma. *Science and Culture*, 8(6): 273-274.
- Muller, J. 1968. Palynology of the Pedawan and Plateau Sandstone Formations (Cretaceous-Eocene) in Sarawak, Malaysia. *Micropaleontology*, 14(1): 1-37.
- Nagy, E. 1985. Sporomorphs of the Neogene in Hungary. *Geologica Hungarica, Series Palaeontologica*, 47, Inst. Geol. Hungar., Budapestini.
- Nakaya, H., Saegusa, H., Kunimatsu, Y., and Ratanasthien, B. 2002. Miocene mammalian faunas of northern Thailand and their geological age. *Primate Research*, 18: 131-141.
- Navale, G.K.B., and Misra, B.K. 1979. Some pollen grains from Neyveli lignite, Tamil Nadu, India. *Geophytology*, 8: 226-239.

- Nayar, B.K., Lata, P., and Tiwari, L.P. 1964. Spore morphology of the ferns of west tropical Africa. *Pollen et Spores*, 6: 545-582.
- Nichols, D.J. 1973. North American and European species of *Momipites* ("Engelhardtia") and related genera. *Geoscience and Man*, 7: 103-117.
- Nichols, D.J., and Ott, H.L. 1978. Biostratigraphy and evolution of the *Mimopites-Caryapollenites* lineage in the Early Tertiary in the Wind River Basin, Wyoming. *Palynology*, 2: 93-112.
- Norton, N.J., and Hall, J.W. 1969. Palynology of the Upper Cretaceous and Lower Tertiary in the type locality of the Hell Creek Formation, Montana, U.S.A. *Palaeontographica*, 125B(1-3): 1-64.
- Ogasawara, K., Ugai, H., and Kurihara, Y. 2003. Short-term Early Miocene climatic fluctuations in the Japanese Islands. In B. Ratanasthien and S.L. Rieb, *Proceedings of the 8<sup>th</sup> International Congress on Pacific Neogene Stratigraphy, Pacific Neogene Paleoenvironments and their evolution* (pp. 181-190), Chiang Mai, Thailand.
- Penny, J.S. 1969. Late Cretaceous and Early Tertiary palynology. In: R.H. Tschudy and R.A. Scott (eds.), *Aspects of Palynology* (pp. 271-289), John Wiley and Sons, New York.
- Playford, G. 1982. Neogene palynomorphs from the Huon Peninsula, Papua New Guinea. *Palynology*, 6: 29-54.
- Pocknall, D.T. 1989. Late Eocene to Early Miocene vegetations and climate history of New Zealand. *Journal of the Royal Society of New Zealand*, 19(1): 1-18.
- Pocknall, D.T. and Crosbie, Y.M., 1982. Taxonomic revision of some Tertiary tricolporate and tricolpate pollen grains from New Zealand. *New Zealand Journal of Botany*, 20: 7-15.



- Pocknall, D.T. and Mildenhall, D.C. 1984. Late Oligocene to Early Miocene spores and pollen from Southland, New Zealand. *New Zealand Geological Survey Palaeontological Bulletin*, 32: 1-66.
- Pocock, S.A.J. 1968. *Zonalapollenites* Pflug 1953 and related genera. *Imperial Oil Limited, Calgary, Canada*: 639-641.
- Potonié, R. 1931a. Zur mikroskopie der Braunkohlen. Tertiäre Blütenstaubformen. *Braunkohle*, 30(H16): 325-333.
- Potonié, R. 1931b. Pollenformen aus tertiären Braunkohlen. *Jahrb. Preuß. Geol. Landesanstalt Jahrb.*, 52: 1-7.
- Potonié, R. 1931c. Zur mikroskopie der Braunkohlen. Tertiäre Sporen-und Blütenstaubformen. *Braunkohle*, 30(H27): 554-556.
- Potonié, R. 1931d. Pollenformen der miocänen Braunkohlen. *Gesell. Naturforsch. Freunde Berlin Sitzungsber., Jahrg*, 1-3: 24-28.
- Potonié, R. 1951a. Pollen-und Sporenformen als Leitfossilien des Tertiärs. *Microscopie*, 6: 272-283.
- Potonié, R. 1951b. Revision stratigraphisch wichtiger Sporomorphen des mitteleuropäischen Tertiärs. *Palaeontographica*, 91B(1-3): 131-151.
- Potonié, R. 1956. Synopsis der Gattungen der Sporae dispersae, I Teil: Sporites. *Beihefte zum Geologischen Jahrbuch*, 23: 1-103.
- Potonié, R. 1958. Synopsis der Gattungen der Sporae dispersae, II Teil: Sporites (Nachtrage), Saccites, Aletes, Praecolpates, Polyplicates, Monocolpates. *Beihefte zum Geologischen Jahrbuch*, 31.
- Potonié, R. 1960. Synopsis der Gattungen der Sporae dispersae, III Teil: Nachtrage sporites, Fortsetzung Pollenites Mit Generalregister zu Teil I-III. *Beihefte zum Geologischen Jahrbuch*, 39: 1-189.

- Potonié, R., Thompson, P.W., and Thiergart, F. 1950. Zur nomenklatur der neogenen Sporomorphae (Pollen und Sporen). *Geologischen Jahrbuch.*, 65: 35-70.
- Potonié, R., and Venitz, A. 1934. Zur mikrobotanik des miozänen Humodils der niederrheinischen Bucht. *Arbeiten Institut für Paläobotanik und Petrographie der Brennsteine*, 5: 1-54.
- Prakash, U. 1971. Fossil woods from the Tertiary of Burma. *Palaeobotanist*, 20(1): 48-70.
- Prakash, U., and Bande, M.B. 1980. Some more fossil woods from the Tertiary of Burma. *Palaeobotanist*, 26(3): 261-278.
- Raatz, G. 1937. Mikrobotanisch-stratigraphische untersuchung der braunkohle des muskauer bogens. *Abhandlungen Preussischen Geologischen Landesanstalt*, 183: 1-35.
- Ramanujam, C.G.K. 1966. Palynology of the Miocene lignite from South Arcot District, Madras, India. *Pollen et Spores*, 8(1): 149-204.
- Rao, M.R. 1996. An Early Miocene palynological assemblage from Turavur bore-hole, Alleppey District, Kerala – its palaeoecological and stratigraphical significance. *Geophytology*, 25: 155-163.
- Ratanasthien, B. 1984. Spore and pollen dating of some Tertiary coal and oil deposits in northern Thailand. *Conference on Application of Geology and the National Development, Chulalongkorn University, Bangkok, 19-22 November 1984*: 273-280.
- Ratanastien, B. (1989). Depositional environment of Mae Lamao basin as indicated by palynology and coal petrography. in: T. Thanasuthipitak and P. Ounchanum (Eds.), *International Symposium on Intermontane Basins: Geology and Resources* (pp. 205-215), Chiang Mai, Thailand.

- Ratanasthien, B. (1990). Mae Long Formation of Li Basin, Thailand. in R. Tsuchi (Ed.), *Pacific Neogene Events, Their Timing, Nature and Interrelationship* (pp. 123-128), University of Tokyo Press.
- Ratanasthien, B. (1997). Algae types of oil source rocks in northern Thailand. in: P. Dheeradilok and others (Eds.), *Conference on Stratigraphy and Tectonic Evolution of Southeast Asia and the South Pacific* (pp. 606-612), Bangkok, Thailand.
- Ratanasthien, B., Kojima, T., Tokumitsu, T., Katoh, A., and Uyemura, N. 1992. Relationship between elementary analysis, origin and diagenesis of Tertiary Thai coals. *Proceedings of the National Conference on Geologic Resources of Thailand: Potential for Future Development, Bangkok: 273-282.*
- Ratanasthien, B., and Promkotra, S. (1994). Coal seams correlation of Li coal deposits. in: P. Angsuwathana and others (Eds.), *International Symposium on Stratigraphic Correlation of Southeast Asia* (pp. 282-290), Bangkok, Thailand.
- Reitsma, T. 1970. Pollen morphology of the Alangiaceae. *Review of Palaeobotany and Palynology*, 10(4): 249-332.
- Ross, N. 1949. On a Cretaceous pollen and spore bearing clay of Scania, a preliminary report. *Geological Institution of the University of Uppsala, Bulletin*, 34: 25-43.
- Sah, S.C.D., and Kar, R.K. 1969. Palynology of the Laki sediments in Kutch- 3. Pollen from the bore-holes around Jhulral, Baranda and Panandhro. *Palaeobotanist*, 18(2): 127-142.
- Salujha, S.K., Kindra, G.S., and Rehman, K. 1972. Palynology of the South Shillong Front part 1. The Paleogene of Garo Hills. *Proc. Sem. Palaeopalynology Indian Stratigraphy: 265-291.*

- Salvador, A. (1994). *International Stratigraphic Guide: A Guide to Stratigraphic Classification, Terminology, and Procedure*. International Subcommission on Stratigraphic Classification of IUGS International Commission on Stratigraphy, 2<sup>nd</sup> edition.
- Santisuk, T. 1997. Geographical and ecological distributions of the two tropical pines, *Pinus kesiya* and *Pinus merkusii*, in Southeast Asia. *Thai Forest Bulletin (Botany)*, Bangkok, 25: 102-123.
- Sasada, M., Ratanasthien, B., and Saponpongpiat, R. 1987. New K/Ar ages from the Lampang basalt, northern Thailand. *Geological Survey of Japan, Bulletin*, 38(1): 13-20.
- Savin, S.M., Douglas, R.G., and Stehli, F.G. 1975. Tertiary marine paleotemperatures. *Geological Society of America, Bulletin*, 86: 1499-1510.
- Sharma, C., and Gupta, A. 1996. LM and SEM pollen studies in Indian Alangiaceae. *Palaeobotanist*, 45: 134-142.
- Silaratana, T., Ratanasthien, B., Fyfe, W.S., Takayasu, K., Asanachinda, P., and Kandharosa, W. (2002). Mineralogical variation of clastic rocks and coal from the Chiang Muan coal field in northern Thailand. in: N. Mantajit (Ed.), *Proceedings of the Symposium on Geology of Thailand* (pp. 233-241), Bangkok, Thailand.
- Silaratana, T., Ratanasthien, B., Takayasu, K., Asanachinda, P., Kandharosa, W., and Kusakabe, M. (2003). Sulfur isotope implication of Middle Miocene marine incursion in northern Thailand. in B. Ratanasthien, S.L. Rieb, and S. Chantraprasert (Eds.), *Proceedings of the 8<sup>th</sup> International Congress on Pacific Neogene Stratigraphy: Pacific Neogene Paleoenvironments and their Evolution* (pp. 356-374), Chiang Mai, Thailand.

- Singh, R.S. 1999. Diversity of *Nypa* in the Indian subcontinent: Late Cretaceous to Recent. *Palaeobotanist*, 48: 147-154.
- Singh, H.P., and Khanna, A.K. 1978. Some fossil species of *Pediastrum* and their palaeoecological significance in the Subathu Formation of Himachal Pradesh. *Palaeobotanist*, 25: 466-473.
- Sithiprasasna, D. 1959. Occurrence of Mae Moh fossils. *Thailand Royal Department of Mines, Report of Investigation*, 2: 29-34.
- Smith, A.H.V., and Butterworth, M.A. 1967. Miospores in the coal seams of the Carboniferous of Great Britain. *Special Paper in Palaeontology, London*, 1: 1-321.
- Snansieng, S., and Manee kut, N. 1985. Li basin, an analysis of the oldest Cenozoic basin of Thailand. *Journal of the Geological Society of Thailand*, 8(1-2): 29-35.
- Sompong, W., Springbelt, G.M., and Evans, P.R. (1996). *Lignite mine development project (phase III) Mae Moh coal deposit*. Electricity Generating Authority of Thailand, Australia AID Geological Report.
- Song, Z., Guan, X., Zheng, Y., Li, Z., Wang, W., and Hu, Z. 1985. *A research on Cenozoic palynology of the Longjing structural area in the shelf basin of the East China Sea (Donghai) region*. Anhui Science and Technology Publishing House.
- Songtham, W. (1996) *Palynology, age and depositional environment Tertiary Laem Pho Section, Krabi Province, southern Thailand*. Investigation Report of Geological Survey Division, Department of Mineral Resources, Bangkok.
- Songtham, W., Ratanasthien, B., Mildenhall, D.C., Singharajwarapan, S, and Khandharosa, W. 2001. Palynological zonations and their paleovegetations of Ban Pa Kha coal mine, Li basin, Changwat Lamphun. *Paper presented at*



*Geological Survey Division Annual Academic Meeting, Department of Mineral Resources, 3-4 September 2001, Bangkok, Thailand: 1-11.*

- Songtham, W., Ratanasthien, B., and Mildenhall, D.C., Singharajwarapan, S., and Khandharosa, W. 2003. Oligocene – Miocene climatic changes in northern Thailand resulting from extrusion tectonics of Southeast Asian landmass. in B. Ratanasthien, S.L. Rieb, and S. Chantraprasert (Eds.), *Proceedings of the 8<sup>th</sup> International Congress on Pacific Neogene Stratigraphy: Pacific Neogene Paleoenvironments and their evolution* (pp. 248-264), Chiang Mai, Thailand.
- Songtham, W., Ratanasthien, B., Watanasak, M., and Mildenhall, D.C. 2000. Temperate palynological elements from Na Hong basin, northern Thailand. *Mahidol Journal*, 7(2): 121-126.
- Srivastava, S.K. 1966. Upper Cretaceous microflora (Maestrichtian) from Scollard, Alberta, Canada. *Pollen et Spores*, 8(3): 497-552.
- Stover, L.E., and Partridge, A.D. 1973. Tertiary and Late Cretaceous spores and pollen from the Gippsland basin, southeastern Australia. *Proceedings of the Royal Society of Victoria*, 85: 237-286.
- Suganuma, Y., Okada, M., and Nagaoka, S. 2002. Preliminary Magnetostratigraphy of the Tertiary sedimentary rocks, Mae Moh Group from Chiang Muan in northern Thailand. *Primate Research*, 18(2): 165-173 (in Japanese with English abstract).
- Sukto, P., Sutheethorn, V., and Boripatkosol, S., Meesook, A., and Sareerat, S. (1976). *Geological map of Thailand, Sheet Moulmein (NE47-14), scale 1: 250,000*. Geological Survey Division, Department of Mineral Resources, Bangkok, Thailand.



- Takahashi, K., and Jux, U. 1991. Miocene palynomorphs from lignites of the Soma Basin (West Anatolia, Turkey). *Bulletin of the Faculty of Liberal Arts, Nagasaki University, Natural Science*, 32(1): 7-165.
- Tankaya, W. (2001). *Implication of geochemistry on depositional environment in Mae Moh coal field Changwat Lampang*. Master's thesis, Chiang Mai University.
- Tapponnier, P., Peltzer, G., Le Dain, Y., and Armijo, R. 1982. Propagating extrusion tectonics in Asia: New insights from simple experiment with plasticine. *Geology*, 10: 611-616.
- Tapponnier, P., Peltzer, G., and Armijo, R. 1986. On the mechanics of the collision between India and Asia. *Geological Society Special Publication*, 19: 115-157.
- Tassy, P., Anupandhanant, P., Ginsburg, L., Mein, P., Ratanasthien, B., and Sutteethorn, V. 1992. A new *Stegolophodon* (Proboscidea, Mammalia) from the Early Miocene of northern Thailand. *Geobios*, 25(4): 511-523.
- Thiergart, T. von. 1937. Die pollenflora der Neiderlausitzer Braunkohle, besonders im profil der Grube Marga bei Senftenberg. *Jahrbuch der Preussischen geologischen Landesanstalt u Bergakademie zu Berlin*, 58: 282-356.
- Thomson, P.W., and Pflug, H. 1953. Pollen und Sporen des mitteleuropalschen Tertiars. *Palaeontographica*, B94: 1-138.
- Traverse, A. (1955). *Pollen analysis of the Brandon Lignite of Vermont*. U.S. Department of Interior, Report of Investigations, 5151: 1-107.
- Traverse, A. (1988). *Paleopalynology*. Unwin Hyman, Boston.
- Tryon, R.M., and Tryon, A.F. (1982). *Ferns and allied plants*. Springer-Verlag, New York.

- Tschudy, R.H., and Van Loenen, S. (1970). Illustrations of plant microfossils from the Yazoo Clay (Jackson Group, Upper Eocene) Mississippi. *U.S. Geological Survey Professional Paper*, 643E: 3 p.
- Tschudy, R.H., Tschudy, B.D., and Craig, L.G. (1984). Palynological evaluation of Cedar Mountain and Burro Canyon Formations, Colorado Plateau. *U.S. Geological Survey Professional Paper*, 1281: 1-21.
- Ukkakimapan, Y. (1992). Geological variation of Ban Pu, Ban Hong and Mae Long coal deposits, Lamphun. in: C. Piancharoen (Ed.), *National Conference on Geologic Resources of Thailand: Potential for Future Development* (pp. 574-583), Bangkok, Thailand.
- Uttamo, W., Elder, C., and Nichols, G. (2003). Sedimentology of Neogene sequences in the Mae Moh basin, northern Thailand. in B. Ratanasthien, S.L. Rieb, and S. Chantraprasert (Eds.), *The 8<sup>th</sup> International Congress on Pacific Neogene Stratigraphy: Pacific Neogene Paleoenvironments and Their Evolution* (pp. 31-48), Chiang Mai, Thailand.
- Van der Hammen, T. 1956. A palynological systematic nomenclature. *Boletín Geológico (Bogotá)*, 4(2-3): 63-101.
- Veevers, J.J., and Li, Z.X. 1991. Review of seafloor spreading around Australia. II. Marine magnetic anomaly modelling. *Australian Journal of Earth Science*, 38: 391-408.
- Von Koenigswald, G.H.R., von. (1959). *A mastodon and other fossil mammals from Thailand*. Report Investigation no. 2, Royal Department of Mines: 25-28.
- Wang, W., and Zhang, D. 1990. Tertiary sporo-pollen assemblages from the Shangdou-Huade basin, Inner Mongolia – with discussion on the formation of steppe vegetation in China. *Acta Micropalaeontologica Sinica*, 7(3): 239-252.

- Watanasak, M. (1988). *Mid-Tertiary palynology of onshore and offshore Thailand*.  
Doctor's thesis, University of Adelaide, Australia.
- Watson, P.V., (1996) *Ph1 and Ph2 well samples Phrae basin Thailand: biostratigraphy and paleoenvironments*. Core Laboratories, prepared for PTT Exploration and Production Public Company Ltd, File No. GSI-96015(B).
- Weir, G.H., and Thurston, E.L. 1977. Scanning electron microscopic identification of fossil Pinaceae pollen to species by surface morphology. *Palynology*, 1: 157-165.
- Willet, S.D., and Beaumont, C. 1994. Subduction of Asian lithospheric mantle beneath Tibet inferred from models of continental collision. *Nature*, 369: 642-645.
- Wilson, M.A. 1978. Palynology of three sections across the uppermost Cretaceous/Paleocene boundary in the Yukon territory and district of Mackenzie, Canada. *Palaeontographica*, Abt.B., 166: 99-183.
- Wilson, L.R., and Hoffmeister, W.S. 1953. Four new species of fossil *Pediastrum*. *American Journal of Science*, 251: 753-760.
- Wilson, L.R., and Webster, R.M. 1946. Plant microfossils from a Fort Union coal of Montana. *American Journal of Botany*, 33: 271-278.
- Wingate, F.H. 1983. Palynology and age of the Elko Formation (Eocene) near Elko, Nevada. *Palynology*, 7: 93-132.
- Wodehouse, R.P. 1933. Tertiary pollen II, The oil shales of the Eocene Green River formation. *Torrey Botanical Club Bulletin*, 60: 479-524.
- Wolff, H. 1934. Mikrofossilien des pliozänen Humodils der Grube Freigericht bei Dettingen a. Main und Vergleich mit älteren Schichten des Tertiärs sowie posttertiären Ablagerungen. *Arbeiten Institut für Paläobotanik und Petrographie des Brennsteine*, 5: 55-101.
- Wongratana, L. (2001) *Phytoplankton*. Kasetsart University Press.

- Wood, G.D., Gabriel, A.M., and Lawson, J.C. 1996. Chapter 3. Palynological techniques-processing and microscopy. In J. Jansonius and D.C. McGregor (eds.), *Palynology: Principles and Applications* (pp. 29-50), American Association of Stratigraphic Palynologists Foundation 1.
- Yabe, A. 2002. Paleoclimatic condition inferred from the Tertiary plant megafossil assemblages from northern Thailand. *Primate Research*, 18(2): 143-157 (in Japanese with English abstract).
- Yakzan, A.M., Morley, R.J., and Koraini, A.M. (1994). Freshwater algal palynomorphs as stratigraphic indicators in Malay basin. in P. Angsuwatana and others (Eds.), *Proceedings of the International Symposium on Stratigraphic Correlation of Southeast Asia* (pp. 207), Bangkok, Thailand.
- Yamanoi, T. 1984. Presence of sonneratiaceous pollen in Middle Miocene sediments, Central Japan. *Review of Palaeobotany and Palynology*, 40: 347-357.
- Zetter, R., and Ferguson, D.K. 2001. Trapaceae pollen in the Cenozoic. *Acta Palaeobotanica*, 41(2): 321-339.
- Zippi, P.A. 1998. Freshwater algae from the Mattagomi Formation (Albian), Ontario: Paleoecology, botanical affinities, and systematic taxonomy. *Micropaleontology*, 44: 1-78.