


TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	iii
ABSTRACT	iv
LIST OF TABLES	ix
LIST OF ILLUSTRATIONS	xi
CHAPTER I INTRODUCTION	1
Principles, theories, and rationale	1
Purposes of the study and hypotheses	2
Anticipated benefits	2
Scope of the study	3
Glossary of terms	3
CHAPTER II LITERATURE REVIEW	5
The permanent magnetic materials and magnetic fields	5
The applications of magnets in orthodontics	10
Biological implications of magnetic fields	12
Fibroblasts in cell culture	14
Investigation methods in the <i>in vitro</i> study	15
CHAPTER III MATERIALS AND METHODS	19
Materials	19
Methods	23
Statistical analyses	36
CHAPTER IV RESULTS	37
The viability and the growth of the cultured human gingival fibroblasts at 3 and 7 days in each experimental group	38

Comparison of the mean ranks of viability of the cultured human gingival fibroblasts among experimental groups	39
Comparison of the mean ranks of viability of the cultured human gingival fibroblasts between 3 days and 7 days	40
Comparison of mean ranks of viability of the cultured human gingival fibroblasts among experimental groups at 3 days and 7 days	40
The correlations between the viability of the cultured human gingival fibroblasts and the extent of magnetic field intensity	41
Comparison of mean ranks of growth (the rate of new DNA synthesis) of the cultured human gingival fibroblasts among experimental groups	43
Comparison of the mean ranks of the growth (the rate of new DNA synthesis) of the cultured human gingival fibroblasts between 3 days and 7 days	44
Comparison of mean ranks of the growth (the rate of new DNA synthesis) of the cultured human gingival fibroblasts among experimental group at 3 days and 7 days	44
The correlations between the growth (the rate of new DNA synthesis) of the cultured human gingival fibroblasts and the extent of magnetic field intensity	45
CHAPTER V DISCUSSION	48
Discussion on results of the study	48
Limitations of the research	51
Suggestion for further studies	51
CHAPTER VI CONCLUSION	53

BIBIOGRAPHY	54
APPENDIX	59
CURRICULUM VITAE	63



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

LIST OF TABLES

Table	Page
3.1 The distribution and the amount of magnetic field intensity of single magnet and the attractive magnets at the position 1 and the position 2	26
3.2 The mean ranks of growth (the rate of new DNA synthesis) of the cultured human gingival fibroblasts between the positive control group (TPA stimulation for 1 day) and the control group at days 3 and 7	30
3.3 Five experimental groups under different conditions of magnetic field	31
3.4 The number of repeated experiments for evaluating cell viability	35
3.5 The number of repeated experiments for evaluating cell growth	35
4.1 Medians (Mdn) and quartiles (P_{25}, P_{75}) of viability and growth (the percentage of new DNA synthesis) of the cultured human gingival fibroblasts at day 3 and day 7 in each experimental group	38
4.2 Comparison of mean ranks of the cultured human gingival fibroblast cell viability among experimental groups	39
4.3 Comparison of mean ranks of viability of the cultured human gingival fibroblasts between 3 days and 7 days	40
4.4 Comparison of the mean ranks of viability of the cultured human gingival fibroblasts among experimental group at 3 days and 7 days	41
4.5 The correlations between the viability of the cultured human gingival fibroblasts and the extent of magnetic field intensity	43
4.6 Comparison of the mean ranks of growth (the rate of new DNA synthesis) of the cultured human gingival fibroblasts among five experimental groups	43

- 4.7 Comparison of the mean ranks of growth (the rate of new DNA synthesis) of the cultured human gingival fibroblasts between 3 days and 7 days 44
- 4.8 Comparison of the mean ranks of growth (the rate of new DNA synthesis) of the cultured human gingival fibroblasts among experimental groups at 3 days and 7 days 45
- 4.9 The correlations between the growth (the rate of new DNA synthesis) of the cultured human gingival fibroblasts in each group and the extent of magnetic field intensity 47

LIST OF ILLUSTRATIONS

Figure		Page
2.1	Iron fillings sprinkled on a sheet of paper covering a bar magnet. The distribution of the fillings suggests the pattern of lines of the magnetic field	7
2.2	The magnetic field lines for a bar magnet. The lines form closed loops, leaving the magnet at its north pole and entering at its south pole	8
2.3	A two-dimensional computerized figure shows magnetic field of repelling magnets, (N) is the north pole and (S) is the south pole	9
2.4	A change in force magnitude with respect to the distance between magnets	10
2.5	The fibroblasts (under a microscope using a 10X objective lens)	14
2.6	A haemocytometer	16
2.7	The flow cytometric system	17
2.8	A flow cytometer	17
2.9	A cell cycle diagram	18
2.10	Sample graphs of cell cycle analyzed by flow cytometric system	
	A) A normal cell cycle distribution of stained cultured cells	
	B) An increase in S phase and G ₂	
	C) Cells arresting in G ₂	18
3.1	Magnet specimens in the dimension of 20 mm in diameter and 2 mm in thickness	20

3.3	Six-well flat bottom microtiter culture plates and culture flasks	21
3.4	A tesla meter 2000 with a Hall probe	21
3.5	A laminar flow biological cabinet	22
3.6	A carbondioxide incubator	22
3.7	A phase contrast microscope	23
3.8	The measurement of single magnetic field at two positions of culture plate: A) The position 1 was closest to the magnet, B) the position 2 was farthest from the magnet.	25
3.9	The measurement of attractive magnetic field at two positions of culture plate: A) The position 1 was closest to the magnets, B) the position 2 was farthest from the magnets.	25
3.10	A 7-AAD versus FITC anti-BrdU dot plot diagram and the percentages of cell population in each phase of cell cycle: R_2 is G_0+G_1 -phase, R_3 is S-phase, R_4 is G_2+M phase, and R_5 is apoptotic phase.	27
3.11	The dot plat diagrams of the BrdU-labeled cultured human gingival fibroblasts at different duration for BrdU uptake, A) 30 min B) 60 min C) 90 min D) 120 min	29
3.12	The cultured human gingival fibroblasts grown in DMEM with 10% FBS and 1% penicillin / streptomycin without exposure to static magnetic field (Group 1)	32
3.13	The cultured human gingival fibroblasts grown in DMEM with 10% FBS and 1% penicillin / streptomycin under high and low single static magnetic field (Group 2 and 3, respectively)	32
3.14	The cultured human gingival fibroblasts grown in DMEM with 10% FBS and 1% penicillin / streptomycin under high and low attractive static magnetic field (Group 4 and 5, respectively)	33

- 3.15 A standard haemocytometer chamber which are ruled into nine large squares. The circle represents an area of a large square. 34
- 4.1 Scatter diagrams between the viability of the cultured human gingival fibroblasts and the extent of magnetic field intensity, a) at low level of magnetic field, b) at high level of magnetic field 42
- 4.2 Scatter diagrams between the growth of the cultured human gingival fibroblasts and the extent of magnetic field intensity, a) at low level of magnetic field, b) at high level of magnetic field 46