REFERENCES

Acharya A, Sanyal SK, Moulik SP. Physicochemical investigations on microemulsification of eucalyptol and water in presence of polyoxyethylene (4) lauryl ether (Brij-30) and ethanol. Int J Pharm 2001; 229: 213-26.

Apinhasmit W, Sobhon P. *Opisthorchis viverrini*: effect of praziquentel on the adult tegument. Southeast Asian J Trop Med Public Health 1996; 27: 304-11.

Avancini RM, Silveira GA. Age structure and abundance in populations of muscoid flies from a poultry facility in Southeast Brazil. Mem Inst Oswaldo Cruz 2000; 95: 259-64.

Blust MH, Hopkins TL. Olfactory responses of a specialist and generalist grasshopper to volatiles of *Artemisia ludoviciana* Nuft. (Asteraceae). J Chem Ecol 1987; 13: 1893-902.

Bohart GE, Gressitt JL. Filth-inhabiting flies of Guam. Bull Bernice P Bishop Mus, 1951.

Brown AWA, Pal R. Insecticide resistance in arthropods. 2nd editions. World Health Organization, Geneva, 1971.

Brown JF, Adkins TR. Relationship of feeding activity of the facefly (*Musca autumnalis*) to production of keratoconjunctivitis in calves. Am J Vet Res 1972; 33: 2551-5.

Byrd JH, Castener JL. Forensic entomology: the utility of arthropods in legal investigations. CRC Press, USA, 2000.

Coats JR. Insecticide mode of action. Academic Press, USA, 1982.

De Vincenzi M, Silano M, De Vincenzi A, Maialetti F, Scazzocchio B. Constituents of aromatic plants: eucalyptol. Fitoterapia 2002; 73: 269-75.

Eldridge BF, Edman JD. Medical entomology. Kluwer Academic Publishers, Netherlands, 2000.

El Gebaly MW, El Nassery SF, El Azzouni MZ, Hammouda NA, Allam SR. Effect of mebendazole and ivermectin in experimental hapatic capillariasis: parasitological, scanning electron microscopy and immunological studies. J Egypt Soc Parasitol 1996; 26: 261-74.

El-Shazly MM, Nassar MI, El-Sherief HA. Toxic effect of ethanolic extract of *Nerium oleander* (Apocynaceae) leaves against different developmental stages of *Muscina stabulans* (Diptera-Muscidae). J Egypt Soc Parasitol 1996; 26: 461-73.

Faouzia h, Souad FT, A TE. Antimicrobial activity of twenty-one Eucalyptus essential oils. Fitoterapia 1993; 64: 71-7.

Ghisalberti EL. Bioactive acylphloroglucinol derivatives from *Eucalyptus* species. Phytochem 1996; 41: 7-22.

Graczyk TK, Knight R, Gilman RH, Cranfield MR. The role of non-biting flies in the epidemiology of human infectious diseases. Microbes Infect 2001; 3: 231-5.

Greenberg B. Flies and disease. Vol I. Ecology, Classification and Biotic Associations. Princeton, Princeton University Press, NJ, 1971.

Greenberg B. Flies and disease. Vol II. Biology and disease transmission. Princeton, Princeton University Press, NJ, 1973.

Hink WF, Fee BJ. Toxicity of D-limonene, the major component of citrus peel oil, to all life stages of the cat flea, *Ctenocephalides felis* (Siphonaptera: Pulicidae). J Med Entomol 1986; 23: 400-4.

Inder Pal S, Hideo E. Biological activity of phloroglucinol derivatives from *Eucalyptus* spp. Nat Proc Sci 1997; 3: 1-7.

Kawada H, Dohara K, Shingo G. Evaluation of larvicidal potency of insect growth regulator, 2 - [1- methyl – 2 (4 – phenoxy phenoxy) ethoxy] pyridine, against the house fly, *Musca domestica*. Jpn J Sanit Zool 1987; 38: 317-22.

Kelling FJ. "Olfaction in houseflies morphology and electrophysiology". Ph.D thesis, University of Groningen, Netherlands, 2001.

Kettle DS. Medical and veterinary entomology. 2nd edition. Cambridge: University Press, 1995.

Keiding J. The house fly: biology and control. World Health Organization, Geneva, 1986.

Kovacs Sz F, Medveczky I, Papp L, Gondar E. Role of prestomal teeth in feeding of the house fly, *Musca domestica* (Diptera; Muscidae). Med Vet Entomol 1990; 4: 331-5.

Kurahashi H, Chowanadisai L. Blow flies (Insecta: Diptera: Calliphoridae) from Indochina. Species Diversity 2001; 6: 185-242.

Lamiri A, Lhaloui S, Benjilali B, Berrada M. Insecticidal effects of essential oils against Hessian fly, *Mayetiola destructor* (Say). Field Crops Res 2001; 71: 9-15.

Laurent D, Vilaseca LA, Chantaraine J, Ballivian C, Saavedra G, Ibanez R. Insecticidal activity of essential oils on *Triatoma infestans*. Phytotherapy Res 1997; 11: 285-90.

Lee s, Peterson CJ, Coats JR. Fumigation toxicity of monoterpenoids to several stored product insects. J Stored Prod Res 2003; 39: 77-85.

Macchioni F, Cioni PL, Flamini G. Morelli I, Perrucci S, Franceschi A, Macchioni G, Ceccarini L. Acaricidal activity of pine essential oils and their main components against *Tyrophagus putrescentiae*, a stored food mite. J Agric Food Chem 2002; 50: 4586-8.

Matsumura F. Toxicology of insecticides. Plenum press, New York and London, 1985.

Maugh TH. To attract or repel, that is the question. Science 1982; 218: 278.

Medveczky I, Kovacs Sz F, Papp L. The role of the housefly, *Musca domestica*, in the spread of Aujeszky' s disease (pseudorabies). Med Vet Entomol 1988; 2: 81-6.

Monzon RB, Sanchez AR, Tadiaman BM, Najos OA, Valencia EG, de Rueda RR, Ventura JVM. A comparison of the role of *Musca domestica* (Linnaeus) and *Chrysomya megacephala* (Fabricius) as mechanical vectors of helminthic parasites in a typical slum area of metropolitan Manila. Southeast Asian J Trop Med Public Health 1991; 2: 222-8.

Morsy TA, Mazyad SA, el-Sharkawy IM. The larvicidal activity of solvent extracts of three medicinal plants against third instar larvae of *Chrysomyia albiceps*. J Egypt Soc Parasitol 1998; 28: 699-709.

Moursy LE. Insecticidal activity of *Calotropis procera* extracts of the flesh fly, *Sarcophaga haemorrhoidalis* Fallen. J Egypt Soc Parasitol 1997; 27: 505-14.

Nayduch D, Honko A, Noblet GP, Stutzenberger F. Detection of *Aeromonas caviae* in the common housefly *Musca domestica* by culture and polymerase chain reaction. Epidemiol Infect 2001; 127: 561-6.

Ndieqie IO, Budenburge WJ, Ofeno DO, Hassanali A. 1,8-cineole, an attractant for the banana weevil, *Cosmopolites sordidus*. Phytochemistry 1996; 42: 369-71.

Picken LG, Jaworski J, Kovac B, Mills JR GD. Traps and baits for flies (Diptera) on Pacific Islands. J Med Entomol 1994; 31: 826-32.

Prates HT, Santos JP, Waquil JM, Fabris JD, Oliveira AB, Foster JE. Insecticidal activity of monoterpenes against *Rhyzopeertha dominica* (F.) and *Tribolium castaneum* (Herbst). J Stored Prod Res 1998; 34: 243-9.

Rice PJ, Coats JR. Insecticidal properties of several monoterpenoids to the house fly (Diptera: Muscidae), Red flour beetle (Coleoptera: Tenebrionidae), and southern corn rootworm (Coleoptera: Chrysomelidae). J Econ Entomol 1994; 87: 1172-9.

Robacker DC. Specific hunger in, *Anastrepha ludens* (Dipt. Tephritidae): effects on attractiveness of proteinaceous and fruit derived lures. Environ Entomol 1991; 20: 1680-6.

Rozendaal JA. Vector control: methods for use by individuals and communities. World Health Organization, Geneva, 1997.

Saito K, Motoyama N, Dauterman WC. Effect of synergists on the oral and topical toxicity of Azamethiphos to organophosphate-resistant houseflies (Diptera: Muscidae). J Econ Entomol 1992; 85: 1041-5.

Sánchez-Ramos I, Castañera P. Acaricidal activity of natural monoterpenes on *Tyrophagus putrescentiae* (Schrank), a mite of store food. J Stored Prod Res 2001; 37: 93-101.

Scott JG, Geden CJ, Rutz DA, Liu NN. Comparative toxicity of seven insecticides to immature stages of *Musca domestica* (Diptera: Muscidae) and two of its important biological control agents, *Muscidifurax raptor* and *Spalangia cameroni* (Hymenoptera: Pteromalidae). J Econ Entomol 199; 8: 776-9.

Shoukry IFI. Toxicological deteriorations of two volatile oils of *Matricaria chamomilla* and *Clerodendron inerme* on the adult house fly *Musca domestica.* J Egypt Soc Parasitol 1997; 27: 893-904.

Smith KGV. Insects and other arthropods of medical importance. British museum (Natural History), London, 1973.

Sucharit S, Tumrasvin W. The survey of flies of medical and veterinary importance in Thailand. Jpn J Sanit Zool 1981; 32: 281-5.

Sucharit S, Tumrasvin W, Vutikes S. A survey of houseflies in Bangkok and neighboring provinces. Southeast Asian J Trop Med Public Health 1976; 7: 85-90.

Sukontason K, Bunchoo M, Sukontason K, Khantawa B, Piangjai S, Rongsriyam Y. Comparison between *Musca domestica* and *Chrysomya megacephala* as carriers of bacteria in Chiang Mai, Thailand. Med Vet Entomol 2003 (In press).

Tripathi AK, Prajapati V, Aggarwal KK, Kumar S. Toxicity, feeding deterrence, and effect of activity of 1,8-cineole from *Artemisia annua* on progeny production of *Tribolium castaneum* (Coleoptera: Tenebrionidae). J Econ Entomol 2001; 94: 979-83.

Tumrasvin W, Shinonaga S. Studies on medically important flies in Thailand. III. Report of species belonging to the genus *Musca* Linne, including the taxonomic keys (Diptera: Muscidae). Bull Tokyo Med Dent Univ 1977; 24: 209-18.

Tumrasvin W, Sucharit S, Kano R. Studies on medically important flies in Thailand. IV. Altitudinal distribution of flies belonging to Muscidae and Calliphoridae in Doi Indhanondh Mountain, Chiengmai, in early summer season. Bull Tokyo Med Dent Univ 1978; 25: 44-81.

Verheyen A, Vanparijs O, Borgers M, Thienpont D. Scanning electron microscopic observations of *Cysticercus fasciolaris* (=*Taenia taeniaeformis*) after treatment of mice with mebendazole. J Parasitol 1978; 64: 411-425.

Wells JD, Kurahashi H. *Chrysomya megacephala* (Fabricius) (Diptera: Calliphoridae) development: rate, variation and the implications for forensic entomology. Jpn J Sanit Zool 1994; 45: 303-9.

World Health Organization. Resistance of vectors of diseases to pesticides. Who Tech Rep Ser 655. Geneva: WHO, 1980.

World Health Organization. Vector control series: the house fly, training and information guide. Geneva: WHO, 1986.

Zumpt F. Myiasis in man and animals in the old world. Butterworth, London, 1965.