

REFERENCES

- Babu, T.D., Kuttan, G., and Padikkala, J. (1995). Cytotoxic and anti-tumour properties of certain taxa of Umbelliferae with special reference to *Centella asiatica* (L.) Urban. *J. Ethnopharmacol.*, 11, 48(1), 53-57.
- Baertschi, S.W., Raney, K.D., Stone, M.P. and Harris, T.M. (1988). Preparation of the 8,9-epoxide of the mycotoxin aflatoxin B₁-the ultimate carcinogenic species. *J. Am. Chem. Soc.*, 110, 7929-7931.
- Bailey, W.A., Iyer, R.S., Stone, M.P., Harris, T.M. and Essigmann, J.M. (1996). Mutational properties of the primary aflatoxin B₁-DNA adduct. *Proc. Natl Acad. Sci. USA*, 80, 2695-2698.
- Baliga, R., Zhang, Z., Baliga, M. and Shah, S.V. (1996). Evidence for cytochrome P-450 as a source of catalytic iron in myoglobinuric acute renal failure. *Kidney Int.*, 49(2), 362-369.
- Busby, W.F. and Wogan, G.N. (1984). Aflatoxins. in Searle, C.D. (ed.), *Chemical Carcinogenesis* (pp. 945-1136), Washington, D.C. : ACS Monograph 182
- Croy, R.G. and Wong, G.N. (1981). Temporal patterns of covalent DNA adducts in rat liver after single dose and multiple doses of aflatoxin B₁. *Cancer Res.*, 41, 197-203.
- Camilleri, P. (1998). *History and Development of Capillary Electrophoresis* (pp.1-22), New York: CRC Press.
- Chaudhuri, S. et al. (1978). Use of common Indian herb Mandukaparni in the treatment of leprosy (preliminary report). *Journal of the Indian Medical Association*, 70, 177-180.
- Chen, Y.J., Dai, Y.S., Chen, B.F., Chang, A., Chen, H.C., Lin, Y.C., Chang, K.H., Lai, Y.L., Chung, C.H. and Lai, Y.J. (1999). The effect of tetrrandrine and extracts of *Centella asiatica* on acute radiation dermatitis in rats. *Biol Pharm Bull.*, 22(7), 703-706.
- Coles, B., Meyer, D.J., Ketterer, B., Stanton, C.A. and Garner, R.C. (1985). Studies on the detoxication of microsomally-activated aflatoxin B₁ by glutathione and glutathione transferases in vitro. *Carcinogenesis*, 6(5), 693-697.

- Darnis, F., Orcel, L., de Saint-Maur, P.P. and Mamou, P. (1979). Use of a titrated extract of *Centella asiatica* in chronic hepatic disorders. *Semaine hopitaux de Paris*, 55, 1749–1750.
- Degan, G.H. and Neumann, H.G. (1978). The major metabolite of aflatoxin B₁ in the rat is a glutathione conjugate. *Chem. Biol. Interact.*, 22, 239-255.
- Department of Chemistry, University of Adelaide, Australia. 2004. "Capillary Electrophoresis" in *Stage 2 Chemistry Social Relevance Projects* [Online]. Available: <http://www.chemistry.adelaide.edu.au/external/soc-rel/content/cap-el.htm> [2004, Aug 28].
- Department of Chemistry, Faculty of Science, Masaryk University. 2004 "Capillary Electrophoresis" [Online]. Available: <http://chemi.muni.cz/~analytika/ce/cze.html> [2004, Aug 29].
- Dwivedi, Y., Rastogi, R., Garg, N.K. and Dhawan, B.N. (1992). Hepatoprotection by picroliv and iridoid glycoside components kutkoside and picroside-1 against galactosamine induced damage in rat liver. *Pharmacology & Toxicology*, 71, 383-387.
- Eaton, D.L. and Groopman, J.D. (1994). *The toxicology of aflatoxins, human health, veterinary, and agricultural significance*, USA: Academic press.
- Egner, P.A., Gange, S.J., Dolan, M.P., Groopman, J.D., Munoz, A. and Kensler, T.W. (1995). Levels of aflatoxin-albumin adduct biomarkers in rat plasma are modulated by both long-term and transient interventions with oltipraz. *Carcinogenesis*, 16, 1769-1773.
- Essigmann, J.M., Croy, R.G., Nadzan, A.M., Busby, W.F., Remholed, M., Buchi, G.B. and Wogan, G.N. (1977). Structural identification of the major DNA adduct formed by aflatoxin B₁ in vitro. *Proc.Natl. Acad. Sci. U.S.A.*, 74, 1870-1874.
- Farnsworth, N.R. and Bunyaphraphatsara, N. (1992). *Thai medicinal plants*, Bangkok: Prachachon.
- Floyd, R.A., Watson, J.J., Harris, J., West, M. and Wong, P.K. (1986). Formation of 8-hydroxydeoxyguanosine, hydroxyl free radical adduct of DNA in granulocytes exposed to the tumor promoter, tetradecanoylphorbolacetate. *Biochem. Biophys. Res. Commun.*, 137(2), 841-846.
- Gan, L-S., Skipper, P.L., Peng, X-C., Groopman, J.D., Chen, J-S., Wogan, G.N. and Tannenbaum, S.R. (1988). Serum albumin adduct in the molecular epidemiology of aflatoxin carcinogenesis: Correlation with aflatoxin B₁ intake and urinary excretion of aflatoxin M₁. *Carcinogenesis (Lond.)*, 9, 1323-1325.

- Geldart, S.E. and Brown, P.R. (1998). Analysis of nucleotides by capillary electrophoresis. *J. Chromatogr. A*, 828, 317-336.
- Gorelick, N.J. (1990). Risk assessment for aflatoxin: Metabolism of aflatoxin B1 by different species. *Risk Anal.*, 10(4), 539-559.
- Green, L.C., Skipper, P.L., Turesky, R.J., Bryant, M.S. and Tannenbaum, S.R. (1984). *In vivo* desitometry of 4-aminobiphenyl in rats via a cysteine adduct in hemoglobin. *Cancer Res.*, 44, 4254-4259.
- Groopman, J.D. (1994). Molecular dosimetry methods for assessing human aflatoxin exposure. in D.L. Eaton and J.D. Groopman (Eds.), *The toxicity of aflatoxins*, USA: Academic press INC.
- Groopman, J.D., Gan, L., Skipper, P., Chen, J., Tannenbaum S.R. and Wogan G.N. (1989). Dosimetry of aflatoxin DNA and serum albumin adducts in people living in Guangxi province. *Proc. Am. Assoc. Cancer Res.*, 30, 317.
- Groopman, J.D., Zhu, J., Donahue, P.R., Pikul, A., Zhang, L-S., Chen, J. and Wogan, G.N. (1992). Molecular dosimetry of urinary aflatoxin DNA adducts in people living in Guangxi Autonomous Region, People's Republic of China. *Cancer Res.*, 52, 45-51.
- Guengerich, F.P. (2001). Forging the links between metabolism and carcinogenesis. *Mutat Res.*, 488(3), 195-209.
- Guengerich, F.P., Johnson, W.W., Ueng, Y.F., Yamazaki, H. and Shimada, T. (1996). Involvement of cytochrome P450, glutathione S-transferase, and epoxide hydrolase in the metabolism of aflatoxin B1 and relevance to risk of human liver cancer. *Environ Health Perspect.*, 104 (3), 557-562.
- Guengerich, F.P., Johnson, W.W., Shimada, T., Ueng, Y.F., Yamazaki, H. and Langouet, S. (1998). Activation and detoxification of aflatoxin B₁. *Mutat. Res.*, 402, 121-128.
- Hadwiger, M.E., Torchia, S.R., Park,S., Biggin, M.E. and Lunte, C.E. (1996). A Study of the Pharmacokinetics of Racemic Isoproterenol using Microdialysis Sampling and Capillary Electrophoresis with Electrochemical Detection. *J. Chromatogr. B.*, 681, 241-249.
- Hayes, A.W. (1978). Bioloical activities of mycotoxins. *Mycopathologia*, 65, 99-105.
- Hayes, J.D., Judah, D.J., McLellan, L.I., Kerr, L.A., Peacock, S.D. and Neal, G.E. (1991a). Ethoxyquin-induced resistance to aflatoxin B₁ in the rat is associated with the expression of

- a novel alpha-class glutathione S-transferase subunit, Yc2, which possesses high catalytic activity for aflatoxin B1-8,9-epoxide. *Biochem. J.*, 279, 385-398.
- Hayes, J.D., Judah, D.J., McLellan, L.I., Neal, G.E. (1991b). Contribution of the glutathione S-transferases to the mechanisms of resistance to aflatoxin B₁. *Pharmacol Ther.*, 50(3), 443-472.
- Hayes, J.D., Judah, D.J., Neal, G.E. and Nguyen, T. (1992). Molecular cloning and heterologous expression of a cDNA encoding a mouse glutathione S-transferase Yc subunit possessing high catalytic activity for aflatoxin B1-8,9-epoxide. *Biochem J.*, 285 (1), 173-180.
- Hsieh, D.P.H. and Wong, J.J. (1982). Metabolism and toxicity of Aflatoxins. in: R. Snyder, D.V. Park, J.J. Kocsis, D.J. Jollow, G.G. Gibson and C.M. Witner (eds.), *Biological Reactive Intermediates II*, part B (pp. 847-863), New York : Plenum Publishers.
- IARC. (1993). *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Some Naturally Occurring Substances: Food Items and Constituents, Heterocyclic Aromatic Amines, and Mycotoxins*, 56, Lyon. France: IARC.
- Iwu, M.M. (1993). *Handbook of African medicinal plants*, Boca Raton F.L.: CRC Press.
- James, R.C. and Hrabison, R.D. (1982). Hepatic glutathione and hepatotoxicity. *Biochemical Pharmacology*, 31, 1829-1835.
- Johnson, W.W., Ueng, Y-F, Mannervick, B., Widersten, M., Hayes, J.D., Sherratt, P., Ketterer, B. and Guengerich, F.P. (1997). Conjugation of highly reactive aflatoxin B1 exo-8,9-epoxide catalyzed by rat and human glutathione transferases: estimation of kinetic parameters. *Biochemistry*, 36, 3056-3060.
- Kartnig, T. (1988). Clinical applications of *Centella asiatica* (L.) Urb. in L.E. Craker, J.E. Simon, (eds.), *Herbs, spices, and medicinal plants: recent advances in botany, horticulture, and pharmacology* (pp.145-173), Phoenix A.Z. : Oryx Press.
- Kensler, T.W., Davis, E.F. and Bolton, M.G. (1994). in D.L. Eaton and J.D. Groopman (Eds.), *The Toxicology of Aflatoxins: Human Health, Veterinary, and Agricultural Significance* (pp. 281-306), San Diego: Academic.
- Kensler, T.W., Egner, P.A., Davidson, N.E., Roebuck, B.D., Pikul, A. and Groopman, J.D. (1986). Modulation of aflatoxin metabolism, aflatoxin-N⁷-guanine formation, and hepatic

- tumorigenesis in rats fed ethoxyquin: role of induction of glutathione-S-transferases. *Cancer Res.*, 46, 3924-3931.
- Kensler, T.W., Gange, S.J., Egner, P.A., Dolan, P.M., Munoz, A., Groopman, J.D., Rogers, A.E. and Roebuck, B.D. (1997). Predictive value of molecular dosimetry: individual versus group effects of oltipraz on aflatoxin-albumin adducts and risk of liver cancer. *Cancer Epi.Bio.Prev.*, 6, 603-610.
- Kuhr, W.G. (1990). Capillary electrophoresis. *Anal. Chem.(Fundamental reviews)*, 62, 403R-414R.
- Klaunig, J.E., Xu, Y., Isenber, J.S., Bachowski, S., Kolaja, K.L., Jiang, J., Stevenson, D.E. and Walborg, E.F.Jr. (1998). The role of oxidative stress in chemical carcinogenesis. *Enviro. Health Perspect.*, 106 (suppl. 1), 289-295.
- Kulhanek, V. and Dimov, D.M. (1966). A new useful modification for the determination of glutamyltranspeptidase activity. *Clin. Chem. Acta.*, 14, 619-623.
- Leung, A.Y. and Steven, F. (1996). *Encyclopedia of Common Natural Ingredients used in food, drugs and cosmetics.* (2nd ed.) New York-Chichester: John Wiley
- Lin, H., Xu, D.K. and Chen, H.Y. (1997). Simultaneous determination of purine bases, ribonucleosides and ribonucleotides by capillary electrophoresis-electrochemistry with a copper electrode. *J. Chromatogr. A*, 760, 227-233.
- Lin, Y.C., Yang, T.I., Chen, J.Y and Yang, C.S. (1972). Search for biologically active substances in Taiwan medicinal plants: Screening for antitumor and antimicrobial substances. *Chinese J Microbiol.*, 5, 76-81.
- Linhardt, R.J. (1994). Capillary electrophoresis of oligosaccharides. *Methods Enzymol.*, 230, 265-280.
- Manson, M.M., Ball, H.W., Barrett, M.C., Clark, H.L., Judah, D.J., Williamson, G. and Neal, G.E. (1997). Mechanism of action of dietary chemoprotective agents in rat liver: induction of phase I and II drug metabolizing enzymes and aflatoxin B₁ metabolism. *Carcinogenesis*, 18(9), 1729-1738.
- Marnett, L.J. (2000). Oxyradicals and DNA damage. *Carcinogenesis*, 21, 361-370.
- McLellan, L.I., Kerr, L.A., Cronshaw, A.D., Hayes, J.D. (1991). Regulation of mouse glutathione S-transferases by chemoprotectors. Molecular evidence for the existence of three distinct

- alpha-class glutathione S-transferase subunits, Ya1, Ya2, and Ya3, in mouse liver. *Biochem J.*, 276 (Pt 2), 461-469.
- Mei, S.R., Yao, Q.H., Cai, L.S., Xing, J., Xu, G.W. and Wu, C.Y. (2003). Capillary electrophoresis with end-column amperometric detection of urinary 8-hydroxy-2'-deoxyguanosine. *Electrophoresis*, 24(9), 1411-1415.
- Miller, D. M., and Wilson, D. M. (1994). Veterinary Diseases Related to Aflatoxins. In D. L. Eaton and J. D. Groopman, (Eds.), *The Toxicology of Aflatoxins* (pp. 347-364), San Diego: Academic Press.
- Moller, L. and Hofer, T. (1997). [32P]ATP mediates formation of 8-hydroxy-2'-deoxyguanosine from 2'-deoxyguanosine, a possible problem in the 32P-postlabeling assay. *Carcinogenesis*, 18, 2415-2419.
- Monning, C.A. and Kenedy, R.T. (1994). Capillary electrophoresis. *Anal. Chem.*, 66, 280R-314R.
- Munoz, N., and Bosch,F.X. (1987). Epidemiology of hepatocellular carcinoma. In L.K. Okuda., and K.G. Ishak(eds.), *Neoplasms of the Liver* (pp. 3-19), Tokyo : Springer.
- Novotny, M. (1990). *Advances in Biomolecular Isolation and Structure Analysis* (p.7), New York: Aster.
- Novotny, M.V. and Sudor, J. (1993). High-performance capillary electrophoresis of glycoconjugates. *Electrophoresis*, 14, 373-389.
- O'Brien, K., Moss, E., Judah, D. and Neal, G. (1983). Metabolic basis of the species difference to aflatoxin B1 induced hepatotoxicity. *Biochem Biophys Res Commun.*, 114(2), 813-821.
- Oda, R.P., Lander, J.P. and Landers, J.P. (1997). Handbook of Capillary Electrophoresis (pp.2-73), London: CRC Press.
- Oefner, P.J. and Chiesa, C. (1994). Capillary electrophoresis of carbohydrates. *Glycobiology*, 4, 397-412.
- Oefner, P.J., Chiesa, C.G., Bonn and Horvath, C. (1994). Developments in capillary electrophoresis of carbohydrates. *J. Capillary Electrophoresis*, 1, 5-26.
- Orlowski, M. and Meister, A. (1963). γ -Glutamyl-p-nitroanilide: a new convenient substrate for determination and study of D- and L-gamma-glutamyl transpeptidase activities. *Biochim Biophys Acta.*, 73, 379-381.

- Palozza, P., Agostara, G., Piccioni, E. and Bartoli, G.M. (1994). Different role of lipid peroxidation in oxidative stress-induced lethal injury in normal and tumor thymocytes. *Arch. Biochem. Biophys.*, 312, 88-94.
- Park, S. and Lunte, C.E. (1995). A perfluorosulfonated ionomer end column electrical decoupler for capillary electrophoresis/electrochemical detection. *Anal. Chem.*, 67, 4366-4370.
- Park, S., Lunte, S.M. and Lunte, C.E. (1995). Voltammetric detection for capillary electrophoresis. *Anal. Chem.*, 67, 911-918.
- Patterson, D.S.P. (1993). Metabolism as a factor in determining the toxic action of aflatoxin in different animal species. *Fed Cosmet Toxicol.*, 11, 1009-1016.
- Peer, F.G. and Linsell, C.A. (1977). Dietary aflatoxins and human primary liver cancer. *Ann. Nutr. Aliment.*, 31, 1005-1017.
- Pereira, M.A., and Chang, L.W. (1980). Binding of chemicalcarcinogens and mutagens to rat hemoglobin. *Chem. Biol. Interact.*, 33, 301-305.
- Quinn, B.A., Crane, T.L., Kocal, T.E., Best, S.J., Cameron, R.G., Rushmore, T.H., Farber, E. and Hayes, M.A. (1990). Protective activity of different hepatic cytosolic glutathione S-transferases against DNA-binding metabolites of aflatoxin B₁. *Toxicol Appl Pharmacol.*, 105 (3), 351-363.
- Ramsdell, H.S. and Eaton, D.L. (1990). Mouse liver glutathione S-transferase isoenzyme activity toward aflatoxin B1-8,9-epoxide and benzo[a]pyrene-7,8-dihydrodiol-9,10-epoxide. *Toxicol Appl. Pharmacol.*, 105(2), 216-225.
- Raney, K.D., Coles, B., Guengerich, F.P. and Harris, T.M. (1992). The endo-8,9-epoxide of aflatoxin B₁: a new metabolite. *Chem. Res. Toxicol.*, 5, 333-335.
- Rastogi R., Saksena S., Garg N.K. and Dhawan B.N. (1995). Effect of picroliv on antioxidant system in liver of rats, after partial hepatectomy. *Phytotherapy Res.*, 9, 364-367.
- Richter, C., Park, J.W. and Ames, B.N. (1988). Normal oxidative damage to mitochondrial and nuclear DNA is extensive. *Proc. Natl. Acad. Sci. U.S.A.*, 85, 6465-6467.
- Rojanapo, W., Kupradinun, P., Tepsuwan, A. and Tanyakaset, M. (1993). Effect of varying the onset of exposure to DDT on its modulation of AFB₁-induced hepatocarcinogenesis in the rat. *Carcinogenesis*, 14, 663-667.

- Sabbioni, G., Skipper, P., Buchi, G. and Tannenbaum, S.R. (1987). Isolation and characterization of the major serum albumin adduct formed by aflatoxin B₁ *in vivo* in rats. *Carcinogenesis*, 8, 819-824.
- Sabbioni, G. and Wild, C.P. (1991). Identification of an aflatoxin G₁-serum albumin adduct and its relevance to the measurement of human exposure to aflatoxins. *Carcinogenesis*, 12, 97-103.
- Scholl, P.F., Musser, S.M., and Groopman, J.D. (1997). Synthesis and characterization of aflatoxin B₁ mercapturic acids and their identification in rat urine. *Chem. Res. Toxicol.*, 10 (10), 1144-1151.
- Schreiber, G., Urban, J., Ziihringer, J., Reutter, W., and Frosch, U. (1971). The secretion of serum protein and the synthesis of albumin and total protein in regenerating rat liver. *J. biol. Chem.*, 246 (14), 4531-4538.
- Shen, H.M., Ong, C.N., Lee, B.L. and Shi, C.Y. (1995). Aflatoxin B₁ induced 8-hydroxy-deoxyguanosine formation in rat hepatic DNA. *Carcinogenesis*, 16, 419-422.
- Shen, H.M., Shi, C.Y., Shen, Y. and Ong, C.N. (1996). Detection of elevated reactive oxygen species level in culture rat hepatocytes treated with aflatoxin B₁. *Free Radical Biol. Med.*, 21, 139-146.
- Shigenaga, M.K., Aboujaoude, E.N., Chen, Q. and Ames, B.N. (1994). Assays of oxidative DNA damage biomarker 8-oxo-2'-deoxyguanosine and 8-oxoguanosine in nuclear DNA and biological fluids by high-performance liquid chromatography with electrochemical detection. *Methods Enzymol.*, 234, 16-33.
- Shigenaga, M.K., Gimeno, C.J. and Ames, B.N. (1989). Urinary 8-hydroxy-2'-deoxy guanosine as a biologic marker of *in vivo* oxidative DNA damage. *Proc. Natl. Acad. Sci. U S A*, 86, 9697-9701.
- Shukla, A., Rasik, A.M. and Dhawan, B.N. (1999). Asiaticoside-induced elevation of antioxidant levels in healing wounds. *Phytother. Res.*, 13(1), 50-54.
- Singh, B. and Rastogi, R.P. (1969). A reinvestigation of the triterpenes of *Centella asiatica*. *Phytochemistry*, 8, 917-921.
- Skipper, P.L., Hutchins, D.H., Turesky, R.J., Sabbioni, G. and Tannenbaum, S.R. (1985). Carcinogen binding to serum albumin. *Proc. Am. Assoc. Cancer Res.*, 26, 356.

- Skipper, P.L. and Tannenbaum, S.R. (1990). The role of protein adducts in the study of chemical carcinogenesis. *Prog Clin Biol Res.*, 340C, 301-310.
- Smela, M.E., Currier, S.S., Bailey, E.A. and Essigmann, J.M. (2001). The chemistry and biology of aflatoxin B₁: from mutational spectrometry to carcinogenesis. *Carcinogenesis*, 22 (4), 535-545.
- Sporn, M.B., Dunlop, N.M., Newton, D.L., Smith, J.M. (1976). Prevention of chemical carcinogenesis by vitamin A and its synthetic analogs (retinoids). *Fed. Proc.*, 35, 1332-1338.
- Sukbunteung, J. (1996). *Effects of lemon grass extract on aflatoxin B₁ mutagenesis activated by liver microsomal enzymes*. M.S. Thesis, Chiang Mai University.
- Swenson, D.H., Miller, E.C. and Miller, J.A. (1974). Aflatoxin B₁-2,3-epoxide: evidence for its formation in rat liver in vivo and by liver human microsomes in vitro. *Biochem. Biophys. Res. Commun.*, 60, 1036-1043.
- Thompson, M., Bye, S.N. and Dutton, M.F. (1992). An investigation into the passage of natural toxin across the digestive tract wall using the everted sac technique. *J. Nat. Tox.*, 1, 9-6.
- Tietz, N.W., (1976). *Fundamentals of Clinical Chemistry*, 2nd ed. (p.622), Philadelphia P.A.: W.B. Saunders Co.
- Tietz, N.W., (1986). *Fundamentals of Clinical Chemistry*, 2nd ed. (p.723), Philadelphia P.A.: W.B. Saunders Co.
- Toyokuni, S., Okamoto, K., Yodoi, J. and Hiai, H. (1995). Persistent oxidative stress in cancer. *FEBS Lett.*, 358, 1-3.
- Tyler, V.E., Brady, L.R. and Robbers, J.E. (1988). *Pharmacognosy*, 9th ed. Philadelphia : Lea and Febiger.
- Veerendra, K.M.H. and Gupta, Y.K. (2003). Effect of Centella asiatica on cognition and oxidative stress in an intracerebroventricular streptozotocin model of Alzheimer's disease in rats. *Clin. Exp. Pharmacol. Physiol.*, 30(5-6), 336-42.
- Wang, J.S., Shen, X., He, X., Zhu, Y.R., Zhang, B.C., Wang, J.B., Qian, G.S., Kuang, S.Y., Zarba, A. and Egner, P.A. (1999). Protective Alterations in Phase 1 and 2 Metabolism of Aflatoxin B₁ by Oltipraz in Residents of Qidong, People's Republic of China. *J. Natl. Cancer Inst.*, 91, 347-354.

- Weiss, D.J. and Lunte, C.E. (2000). Detection of a urinary biomarker for oxidative DNA damage 8-hydroxydeoxyguanosine by capillary electrophoresis with electrochemical detection. *Electrophoresis*, 21(10), 2080-2085.
- Weston, A., Brown, P.R. (1997). *HPLC and CE: Principles and Practice* (p. 134), New York: Academic Press.
- Wild, C.P., Garner, R.C., Montesano, R., Tursi, F. (1986). Aflatoxin B₁ binding to plasma albumin and liver DNA upon chronic administration to rats. *Carcinogenesis*, 7, 853-858.
- Wild, C.P., Jiang, Y., Sabbioni, G., Chapot, G., Montesano, R. (1990a). Evaluation of methods for quantitation of aflatoxin-albumin adducts and their application to human exposure assessment. *Cancer Res.*, 50, 245-251.
- Wild, C.P., Jiang, Y., Allen, S.J., Jansen, L.A.M., Hall, A.J. and Montesano, R. (1990b). Aflatoxin-albumin adducts in human sera from different regions of the world. *Cancer Res.*, 50, 245-251.
- Wong, Z.A., Wei, C.I., Rice, D.W. and Hsieh, D.P.H. (1981). Effects of Phenobarbital pretreatment on the metabolism and toxicokinetics of aflatoxin B₁ in the rhesus monkey. *Toxicol. Appl. Pharmacol.*, 60, 387-397.
- Yaborough, A., Zhang, Y.J., Hsu, T.M., Santella, R.M., (1996). Immunoperoxidase detection of 8-hydroxydeoxyguanosine in aflatoxin B₁-treated rat liver and human oral mucosal cells, *Cancer Res.*, 56, 683-688.
- Yang, C.F., Lin, J., Wasser, S., Shen, H.M., Tan, C.E.L. and Ong, C.N. (2000). Inhibition of ebselen on aflatoxin B₁-induced hepatocarcinogenesis in Fischer 344 rats. *Carcinogenesis*, 21, 2237-2243.
- Zainol, M.K., Abd-Hamid, A., Yusof, S. and Muse, R. (2003). Antioxidative activity and total phenolic compounds of leaf, root and petiole of four accessions of *Centella asiatica* (L.) Urban. *Food Chemistry*, 81(4), 575-581.