


TABLE OF CONTENTS

	Page
Acknowledgements	iii
Abstract in Thai	vi
Abstract in English	ix
Table of Contents	xiii
List of tables	xvii
List of illustrations	xxi
CHAPTER 1 INTRODUCTION	1
1.1 Rationale of the study	1
1.2 Objectives	5
CHAPTER 2 LITERATURE REVIEW	7
2.1 Geography of Thailand	7
2.2.1 Thailand in Pre-Cambrian and Paleozoic	9
2.1.2 Thailand in Mesozoic	13
2.1.3 Thailand in Cenozoic	15
2.1.4 Thailand at present day	17
2.1.5 Climatic condition of recent Thailand	20
2.2 Geography of Northern Thailand	23
2.3 Biodiversity	26
2.3.1 The definition of biodiversity	26

2.3.2 The need of biodiversity	27
2.3.3 The lost of biodiversity; extinction	28
2.3.4 The way to maintain the biodiversity	30
2.4 The study of insects	31
2.5 The study of insects in Thailand	34
2.5.1 Trichoptera	34
2.5.2 Geometridae; Lepidoptera	39
2.6 Riparian zones	43
2.6.1 Terrestrial to aquatic ecosystems	44
2.6.2 Aquatic to terrestrial ecosystems	45
CHAPTER 3 MATERIALS AND METHODS	46
3.1 Study sites	46
3.2 Environmental parameters	49
3.2.1 Parameters in terrestrial ecosystem	50
3.2.2 Parameters in aquatic ecosystem	51
3.3 Riparian insects sampling	55
3.3.1 The sampling methods	55
3.3.2 Identification and preservation	58
3.4 The analysis of data	59
3.5 Research duration	60
3.6 Some important notes about this research	60
CHAPTER 4 RESULTS	61

4.1 Environmental parameters	61
4.2 The riparian insects	64
4.2.1 Pitfall traps	64
4.2.2 Beating tray method	68
4.2.3. Sweeping net	71
4.2.4 Malaise Trap	74
4.2.5 Trichoptera	75
4.2.6 Geometridae; Lepidoptera	84
4.3 Temporal pattern of Trichoptera	91
4.4 The endemic Trichoptera	97
CHAPTER 5 DISCUSSION	99
5.1 Environmental parameters	99
5.2 Ground dwelling insects from pitfall trap	99
5.3 The insects from beating tray	100
5.4 The insects from sweeping net	100
5.5 The discussion on Trichoptera	101
5.6 The discussion on Geometridae	103
5.7 Discussion on Trichoptera and Geometridae	106
5.8 The temporal pattern of Trichoptera	107
5.9 Riparian zone, the habitat of endemic species	108
5.10 Further studies of insects in Thailand	109
CHAPTER 6 CONCLUSION	112

REFERENCES	116
APPENDIX	134
APPENDIX A	134
APPENDIX B	143
APPENDIX C	152
APPENDIX D	167
CURRICULUM VITAE	177



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

LIST OF TABLES

Table	Page
2.1 The traveling steps of Indochina and Sibumasu blocks with geological and biological events in each period of Palaeozoic (References; Wiggins, 1984; Borrer <i>et al.</i> , 1989; William and Feltmate, 1992; Metcalfe, 1998).	12
2.2 The traveling of the Indochina and Sibumasu blocks with geological and biological events in each period of Mesozoic (References: Borrer <i>et al.</i> , 1989; William and Feltmate. 1992; Metcalfe, 1998; Mey, 1998).	13
2.3 Geological and biological events in each period of Cenozoic (References: Borrer <i>et al.</i> , 1989; William and Feltmate, 1992; Mey, 1998; Hall, 1998).	16
2.4 The distribution of Trichoptera in Chiang Mai (CM) and biogeographical region PA = Palearctic, OR = Oriental, NA = Nearctic, AU = Australasian, AF = Afrotropical and NT = Neotropical.	35
2.5 The approximated species number of Geometridae in each biogeographical region (Reference: Scoble, 1999).	41

- 3.1 The major rock type on Doi Suthep-Pui and Doi Inthanon. X to XXX indicate the composition of each rock type and their relative proportion from minor to major composition (Modified from: Rhodes *et al.*, 2000) 46
- 3.2 The specified program of spectrophotometers HACH DR2000 and wavelength for analysis the concentration of sulfate, phosphate, nitrate and ammonia in water sample. The specific reagents for developing chemical reaction from HACH Company are also mentioned. 54
- 4.1 Mean and standard deviation of some environmental parameters in Doi Suthep-Pui and Doi Inthanon at the elevation 600, 1,000 and 1,200. (* Indicate the significant difference at $p < 0.05$). 63
- 4.2 The family and percentage of insect yield captured from pitfall trap from Doi Suthep-Pui and Doi Inthanon at 600, 1,000 and 1,200 m. The results were arranged from the most abundance family to the lower. 65
- 4.3 The Arthropod taxa and their percentage yield that were captured by beating tray method from the lower reach canopy in riparian forest on Doi Suthep-Pui and Doi Inthanon at 600, 1,000 and 1,200 m. 69
- 4.4 The Arthropod taxa and their percentage yield that were captured by sweeping net from the shrub in riparian forest at Doi Suthep-Pui and Doi Inthanon at 600, 1,000 and 1,200. 72

4.5	The insects order and their abundance calculated in percentage that captured with Malaise trap from Doi Suthep-Pui and Doi Inthanon.	74
4.6	Trichoptera from Light trap from Doi Suthep-Pui and Doi Inthanon at 600, 1,000 and 1,200 m.	75
4.7	The site specific species of Trichoptera distribute in each study site on Doi Suthep-Pui and Doi Inthanon.	81
4.8	Species of Trichoptera family, found on Doi Suthep-Pui and Doi Inthanon with the number of total and sheared species.	83
4.9	The Geometridae (Order Lepidoptera) from Doi Suthep-Pui and Doi Inthanon.	85
4.10	Number of taxa present at particular localities, addressed in the Fig 4.10 "Z" mean Thailand recorded species, but unspecified the site.	88
4.11	Trichoptera that can be found throughout the year on Doi Suthep-Pui and Doi Inthanon National Park. The shaded cells indicate occurrence.	93
4.12	Trichoptera presenting during the southwestern moonsoon period on Doi Suthep-Pui and Doi Inthanon National Park. The shaded cells indicate occurrence.	94
4.13	Trichoptera presenting during the northeastern moonsoon period on Doi Suthep-Pui and Doi Inthanon National Park. The shaded cells indicate occurrence.	95

4.14	Trichoptera presenting on the pre and/or post southwestern moonsoon period on Doi Suthep-Pui and Doi Inthanon National Park. The shaded cells indicate occurrence.	96
4.15	The 42 species of Trichoptera from Doi Suthep-Pui National Park that were evaluated as the endemic species on the mountain.	97
4.16	The 60 species of Trichoptera from Doi Inthanon National Park that were evaluated as the endemic species on this mountain.	98
A.1	The family name with their amount of insects captured by pitfall trap from Doi Suthep-Pui and Doi Inthanon at 600, 1,000 and 1,200 m.	136
A.2	The family name with their amount of insects captured by beating tray from Doi Suthep-Pui and Doi Inthanon at 600, 1,000 and 1,200 m.	140
A.3	The family name with their amount of insects captured by sweeping net from Doi Suthep-Pui and Doi Inthanon at 600, 1,000 and 1,200 m.	142
A.4	Caddisflies from Doi Suthep-Pui and Doi Inthanon with the altitudinal preference, distribution, abundance, phenology and references.	145
A.5	Caddisflies from Doi Suthep-Pui and Doi Inthanon National Park capture with UV-dry light trap in 2002. The numbers of specimens are indicated follow by the location with the elevation in blanket.	168

LIST OF ILLUSTRATIONS

Figure	Page
2.1 Map of SE Asia showing microcontinents; Indochina and Shan-Thai (Sibumatsu) blocks that becoming the base of recent Thailand (Reference: Buffetaul and Suteethorn, 1999).	8
2.2 Distribution of land and sea in Cambrian and Ordovician (Cambro-Ordovician). NC = North China, SC = South China, T = Tarim, I = Indochina , QI = Qiangtang, L = Lhasa, S = Sibumatsu , WB = West Burma, GI = Greater India (Reference: Metcalfe, 1998).	10
2.3 Distribution of land and sea in Late Devonian. NC = North China, SC = South China, T = Tarim, I = Indochina , QI = Qiangtang, L = Lhasa, S = Sibumatsu . WB = West Burma (Reference: Metcalfe, 1998).	10
2.4 Distribution of land and sea in Late Permian. The Pangea was the combination continent between the Gondwana and Laurasia. Meanwhile, the microcontinents of SE Asia including Indochina and Sibumatsu blocks were rifted apart from the Gondwana. NC = North China, SC = South China, T = Tarim, I = Indochina , QI = Qiangtang, L = Lhasa, S = Sibumatsu , WB = West Burma (Reference: Metcalfe, 1998).	11


- 2.5 Distribution of land and sea in Triassic. NC = North China, SC = South China, T = Tarim, **I = Indochina**, QI = Qiangtang, L = Lhasa, **S = Sibusatsu**, WB = West Burma (Reference: Metcalfe, 1998). 14
- 2.6 Distribution of land and sea in Jurassic. NC = North China, SC = South China, T = Tarim, **I = Indochina**, QI = Qiangtang, L = Lhasa, **S = Sibusatsu**, WB = West Burma (Reference: Metcalfe, 1998). 14
- 2.7 Distribution of land and sea in Cretaceous. NC = North China, SC = South China, T = Tarim, **I = Indochina**, QI = Qiangtang, L = Lhasa, **S = Sibusatsu**, WB = West Burma (Reference: Metcalfe, 1998). 15
- 2.8 The simplified model for an explanation of the impact of India plate collision with the Eurasia that caused the development of major strike-slip faults on SE Asia (Reference: Hall, 1998). 16
- 2.9 Model shows the simulation of land and sea formation of SE Asia in the epoch; Oligocene, Miocene and Pliocene in Tertiary era. This plate tectonic movement simulation model was created by Hall, 1998. 17
- 2.10 Map of the recent world with major Biogeographic regions; Palaearctic, Oriental, Nearctic, Australasian, Afrotropical and Neotropical (Reference: William and Feltmate, 1992). 18
- 2.11 Map shows the division of Oriental region into 4 sub-regions; Indian, Indo-Chinese, Sundaic and Wallacean (Reference: Simpson and Bugna, 2001). 18

- 2.12 Map showing the mountain system of the connection and continuity of mountain ranges in Southeast Asia with the Great Himalayan range. 19
- 2.13 This map shows the 20% of the remaining forest patches in Thailand and nearby countries. The remaining forest patches are roughly indicated with the dark gray region in the Map. The number in the map indicate each provinces in Thailand (Reference: Parneli *et al.*, 2003). 21
- 2.14 Schematic of the Northeastern and Southwestern monsoons and seasons of Thailand. 23
- 2.15 Map presents the position of 4 major mountain range in Northern Thailand. A = Thanon Thong Chai, B = Dan Lao, C = Pee Pan Nam and D = Luang Pra Bang. 24
- 3.1 The geographical map of Doi Suthep-Pui and Doi Inthanon (Reference: Rhodes *et al.*, 2000). 47
- 3.2 The Topography map show the elevation of Doi Suthep-Pui and Doi Inthanon. 48
- 3.3 The model dry light trap. UV Lamp was operated under the DC battery 12 Volt. Chloroform in bottle was gradually evaporated pass through the wick in order to knock down the trapped insects. Egg package was applied for reduction the struggle of the trapped insects. 56

- 4.1 Cluster analysis (Flexible UPGMA) shows the classification of study sites based on the community of insect family captured from Pitfall traps. Su600 is Doi Suthep-Pui at 600 m; In600 is Doi Inthanon at 600 m. 67
- 4.2 Ordination (semi-strong hybrid multidimensional scaling) shows the relationship of study sites based on their insect community from pitfall trap. Stress = 0.74×10^{-1} . 67
- 4.3 Cluster analysis (Flexible UPGMA) shows the classification of study sites based on the community of Arthropod taxa captured from Beating tray method. 70
- 4.4 Ordination (semi-strong hybrid multidimensional scaling) shows the relationship of study sites based on their insect community from beating tray method. Stress = 0.11. 71
- 4.5 Cluster analysis (Flexible UPGMA) shows the classification of study sites based on the community of insects taxa captured from sweeping net. 73
- 4.6 Ordination (semi-strong hybrid multidimensional scaling) shows the relationship of study sites based on their insect community from sweeping net. Stress = 0.91×10^{-1} . 73
- 4.7 Cluster analysis (Flexible UPGMA) shows the classification of study sites based on the presence-absence of Trichoptera species. Su600 is Doi Suthep-Pui at 600 m; In600 is Doi Inthanon at 600 m. 79

- 4.8 Ordination (semi-strong hybrid multidimensional scaling) shows the relationship of study sites based on their Trichoptera community. *Kisasura surasa* (Philopotamidae) was identified by PCC as the most highly correlated species to the ordination. Stress = 0.3×10^{-5} . 79
- 4.9 TWINSpan summary shows the classification of study sites based on their Trichoptera community. The significant indicator species is *Ugandatrichia maliwan* (Hydroptilidae). 80
- 4.10 The locality record of Geometridae in Thailand; A = Doi Inthanon Doi Suthep-Pui (Chiang Mai), B = Doi Phukha (Nan), C = Phu Luang and Phu Rua (Loei), D = Phu Hin Rongkla and Thung Salaeng Luang (Phitsanulok), E = Nam Nao National Park (Phetchabun), F = Nam Proam Dam (Khon Kaen), G = Chulabhone Dam (Chaiyaphum), H = Khao Yai (Nakhon Nayok), I = Khao Soi Dow (Chanthaburi), J = Earawan and Tum Than Lod (Kanchanaburi), K = Khong Nahka (Ranong), L = Khao Sok (Surat Thani), M = Khao Luang (Nakhon Sri Thammarat) and N = Num Tok Ton Sai (Phuket). 87
- 4.11 Ordination of localities evaluated by geometrid moths 180 species. The meaning of each locality is showed on the Fig 4.6. 89
- 4.12 Dendrogram shows the classification of major localities on Geometridae in Thailand (180 species). 90
- 4.13 The species richness of Trichoptera in each month. The data includes species from Doi Suthep-Pui and Doi Inthanon National Park. 91

- 5.1 Model shows the simulation of the Geometrid locality positions on the plate tectonic movement of Southeast Asia since 30 Ma (Million years ago) until 5 Ma. The plate tectonic movement simulation model was created by Hall, 1998.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved