

BIBLIOGRAPHY

- Ang, K. K. H., Holmes, M. J., Higa, T., Hamann, M. T., Kara, U. A. K. (2000). *In vivo* antimalarial activity of the beta-carboline alkaloid manzamine A. *Antimicrob. Agents Chemother.*, 44(6), 1645-1649.
- Arbain, D., Sargent, M.V. (1987). The alkaloids of *Picrasma javanica*. *Aust. J. Chem.*, 9, 1527-1536.
- Bray, P.G., Ward, S.A. (1993). Malarial chemotherapy : resistance to quinoline containing drugs in *Plasmodium falciparum*. *FEMS Microbiol. Lett.*, 133, 1-7.
- Cain, M., Martei, R., Cook, J.M. (1982). Dichlorodicyanoquinone oxidations in the indole area. Synthesis of crenatine. *J. Org. Chem.*, 47(25), 4933-4936.
- Cumming, J.N., Wang, D., Park, S.B., Shapiro, T.A., Posner, G.H. (1998). Design, synthesis, derivatization, and structure-activity relationships of simplified, tricyclic, 1,2,3-trioxane analogues of the antimalarial artemisinin. *J. Med. Chem.*, 41(6), 952-964.
- Desjardins, R.E., Canfield, C.J., Haynes, D.E., Chulay, J.D. (1979). Quantitative assessment of antimalarial activity *in vitro* by a semiautomated microdilution technique. *Antimicrob. Agents Chemother.*, 16, 710-718.
- Frédérich, M., Tits, M., Hayette, M.-P., Brandt, V., Penelle, J., DeMol, P., Llabrés, G., Angenot, L. (1999). 10'-Hydroxyusambarensine, a new antimalarial bisindole alkaloid from the roots of *Strychnos usambarensis*. 62(4): 619-621.
- Fujimoto, Y., Sato, Y., Made, S. (1989). Isolation of anticancer germacrane derivatives from *Picrasma javanica*. *Patent-Japan Kokai Tokkyo Koho-01 228,979*: 6PP; NAPRALERT database 08/20/97.
- Hagen, T.J., Narayanan, K., Names, J., Cook, J.M. (1989). DDQ oxidations in the indole area. Synthesis of 4-alkoxy- β -carbolines including the natural products crenatine and 1-methoxycanthin-6-one. *J. Org. Chem.*, 54(9), 2107-2178.
- Ishii, K., Koike, K., Ohmoto, T. (1991). Javanicinosides D-H, quassinoid glucosides from *Picrasma javanica*. *Phytochemistry*, 30(12), 4099-4103.

- Johns, S.R., Lamberton, J.A., Sioumis, A.A. (1970). 4-Methoxy-1-vinyl- β -carboline, a new alkaloid from *Picrasma javanica* (Simaroubaceae). *Aust. J. Chem.*, 23, 629-630.
- Kardono, L. B S., Angerhofer, C. K., Tsauri, S., Padmawinata, K., Pezzuto, J. M., Kinghorn, A. D. (1991). Cytotoxic and antimalarial constituents of the roots of *Eurycoma longifolia*. *J. Nat. Prod.*, 54(5), 1360-1367.
- Ketusinh, O. (1948). Report on clinical antimalarial therapy of Thai medicinal plants. *Proceeding of the Siriraj 60 th Anniversary Meeting*, 275-281.
- Kim, H.-S., Yasuharu Shibata, Wataya, Y., Tsuchiya, K., Masuyama, A., Nojima, M. (1999). Synthesis and antimalarial activity of cyclic peroxides, 1,2,4,5,7-pentaxocanes and 1,2,4,5-tetroxanes. *J. Med. Chem.*, 42(14), 2604-2609.
- Kirsch, G., Kong, G. M., Wright, A. D., Kaminsky, R. (2000). A new bioactive sesterterpene and antiplasmodial alkaloids from the marine sponge *Hyrtios cf. erecta*. *J. Nat. Prod.*, 63, 825-829.
- Kitagawa, I., Mahmud, T., Simanjuntak, P., Hori, K., Uji, T., Shibuya, H. (1994). Indonesian medicinal plants. VIII. Chemical structures of three new triterpenoids, bruceajavanin A, dihydrobruceajavanin A, and bruceajavanin B, and a new alkaloidal glycoside, bruceacanthinoside, from the stem of *Brucea javanica* (Simaroubaceae). *Chem. Pharm. Bull.*, 42(7), 1416-1421.
- Koike, K., Ohmoto, T. (1990). Quassinoids from *Picrasma javanica*. *Phytochemistry*, 29(8), 2617-2621.
- Koike, K., Ishii, K., Mitsunaga, K., Ohmoto, T. (1990). New Quassinoids from Indonesian *Picrasma javanica*. Structures of Javanicine E, F, G and M. *Chem. Pharm. Bull.* 38(10), 2746-2749.
- Koike, K., Ishii, K., Mitsunaga, K., Ohmoto, T. (1991a). Quassinoids from *Picrasma javanica*. *Phytochemistry*, 30(3), 933-936.
- Koike, K., Ishii, K., Mitsunaga, K., Ohmoto, T. (1991b). New Des-4-methylpicrasane Quassinoids from *Picrasma javanica*. *J. Nat. Prod.* 54(3), 837-843.
- Koike, K., Ishii, K., Mitsunaga, K., Ohmoto, T. (1991c). Javanicins N, P, and Q, new quassinoids from *Picrasma javanica*. *Chem. Pharm. Bull.*, 39(4), 939-941.

- Koike, K., Ishii, K., Mitsunaga, K., Ohmoto, T. (1991d). New quassinoids from *Picrasma javanica*. Structures of javanicines U, V, W, X and Y. *Chem. Pharm. Bull.*, 39(8), 2021-2023.
- Koike, K., Ohmoto, T. (1992). New quassinoid glucosides, javanicinosides, I, J, K and L from *Picrasma Javanica*. *J. Nat. Prod.*, 55(4), 482-486.
- Koike, K., Ohmoto, T., Uchida, A., Onishi, I. (1994). Javacarboline, A new β -carboline alkaloid from the stem of *Picrasma javanica* in Java. *Heterocycles*, 38(6), 1413-1420.
- Koike, K., Yokoh, M., Furukawa, M., Ishii, S., Ohmoto, T. (1995). Picrasane quassinoids from *Picrasma javanica*. *Phytochemistry*, 40(1), 233-238.
- Komchonwongpaisan, S., Paitayatat, S., Thebtaranonth, Y., Wilairat, P., Yuthavong, Y. (1995). Mechanism-based development of new antimalarials: synthesis of derivatives of artemisinin attached to iron chelators. *J. Med. Chem.*, 38, 2311-2316.
- Kosin, J. (1996). Phytochemical study on the stem bark of *Garcinia atroviridis*. Bangkok: Chulalongkorn University, Thisis.
- Nosten, F., Price, R.N. (1995). New antimalarials drug. A risk-benefit analysis. *Drug-Saf.*, 12(4), 264-273.
- Ohmoto, T., Koike, K., Mitsunaga, K., Kagei, K. (1987). Alkaloids from *Picrasma javanica* growing in Indonesia. *Shoyakugaku Zasshi*, 41(4), 338-340; NAPRALERT database 08/20/97.
- Ohmoto, T., Koike, K., Mitsunaga, K., Fukuda, H., Kagei, K. (1989). Studies on the constituents of Indonesian *Picrasma javanica*. III. Structure of new quassinoids. Javanicins A, C and D. *Chem. Pharm. Bull.*, 37(11), 2991-2994.
- Old style Doctor Association. (1962). *Pra-muan-sappa-kun-ya-thai*. Part I. p-107. Bangkok: Amphon-pittaya.
- Olliaro, P., Cattani, J., Wirth, D. (1996). Malaria, the submerged disease. *J. Am. Med. Assoc.*, 275, 230-233.
- Pavanand, K., Yongvanitchit, K., Webster, H.K., Dechatiwongse, T., Nutakul, W., Jewwachdamrongkul, Y., Bansiddhi, J. (1988). *In vitro* antimalarial activity of a Thai medicinal plant *Picrasma javanica* Bl. *Phytotherapy Research*, 2, 33 – 36.

- Suzuki, H., Unemoto M., Hagiwara M., Ohyama T., Yokoyama Y., Murakami Y. (1999). Synthetic studies on indoles and related compounds. Part 46. First total synthesis of 4,8-dioxygenated β -carboline alkaloids. *J. Chem. Soc., Perkin Trans. 1*, 1717-1723.
- Willems, T.E. (1991). Molecular genetics of drug resistance in *Plasmodium falciparum* malaria. *Parasitol.*, 7, 110-112.
- World Health Organization. (1998). *The world health report 1998-Life in the 21 st century: a vision for all*. p 98. Geneva.
- Wright, C. W., Addac-Kyereme, J., Breen, A. G., Brown, J. E., Cox, M. F., Croft, S. L., Gokcek, Y., Kendrick, H., Phillips, R. M., Pollet, P. L. (2001). Synthesis and Evaluation of Cryptolepine Analogues for Their Potential as New Antimalarial Agents. *J. Med. Chem.*, 44(19), 3187-3194.
- Wright, C. W., Bray, D. H., O'Neill, M. J., Warhurst, D. C., Phillipson, J. D., Quetin-Lectercq, J., Angenot, L. (1991). Antiamoebic and antiplasmodial activities of alkaloids isolated from *Strychnos usambarensis*. *Planta Med.*, 57, 337-340.
- Yoshikawa, M., Harada, E., Aoki, S., Yamahara, J., Murakami, N., Shibuya, H., Kitagawa, I. (1993). Indonesia medicinal plants. VI. On the chemical constituents of the bark of *Picrasma javanica* Bl. (Simaroubaceae) from Flores island. Absolute stereostructures of picrajavanins A and B. *Chem. Pharm. Bull.*, 41(12), 2101-2105.