

**TABLE OF CONTENTS**

	<b>PAGE</b>
ACKNOWLEDGEMENT	iii
ENGLISH ABSTRACT	iv
THAI ABSTRACT	vi
LIST OF TABLE	ix
LIST OF ILLUSTRATIONS	xi
LIST OF ABBREVIATION AND SYMBOLS	xv
CHAPTER I INTRODUCTION	1
CHAPTER II MATERIALS AND METHODS	14
CHAPTER III RESULTS	26
CHAPTER IV DISCUSSION	59
CHAPTER V CONCLUSION	63
REFERENCE	64
CURRICULUM VITAE	70

## LIST OF TABLES

TABLES		PAGE
1	The body weight of male rats treated with 0.5, 1.0 and 1.5 mg/kg methyl parathion	27
2	The body weight of female rats treated with 0.5, 1.0 and 1.5 mg/kg methyl parathion	28
3	The body weight of male rats treated with 0.5, 1.0 and 1.5 %W/W ginger diet	29
4	The body weight of female rats treated with 0.5, 1.0 and 1.5 %W/W ginger diet	30
5	The body weight of male rats treated with 1.0 mg/kg methyl parathion and 1.5 %W/W ginger diet	31
6	The body weight of female rats treated with 1.0 mg/kg methyl parathion and 1.5 %W/W ginger diet	31
7	The amount of TBARS, AChE, GSH, GST and GR in male rats exposed to 0.5, 1.0 and 1.5 mg/kg methyl parathion	33
8	The amount of TBARS, AChE, GSH, GST and GR in female rats exposed to 0.5, 1.0 and 1.5 mg/kg methyl parathion	34
9	The amount of TBARS, AChE, GSH, GST and GR in male rats exposed to 0.5, 1.0 and 1.5%W/W ginger diet	41
10	The amount of TBARS, AChE, GSH, GST and GR in female rats exposed to 0.5, 1.0 and 1.5%W/W ginger diet	42

<b>TABLES</b>	<b>PAGE</b>
11      The amount of TBARS, AChE, GSH, GST and GR in male rats treated with 1.0 mg/kg methyl parathion and 1.5 %W/W ginger diet for 4 weeks	49
12      The amount of TBARS, AChE, GSH, GST and GR in female rats during with 1.0 mg/kg methyl parathion and 1.5 %W/W ginger diet for 4 weeks	49
13      The number of motor endplate positive AChE activity in muscle of male and female rats treated with 1.0 mg/kg methyl parathion and 1.5 %W/W ginger diet for 4 weeks	53

## LIST OF ILLUSTRATIONS

FIGURES	PAGE	
1	Structure of Methyl parathion	2
2	Metabolic pathways of methyl parathion	3
3	Active oxygen species	4
4	Oxidative injury can cause depletion of free radical scavenging antioxidants and antioxidant enzymes	4
5	The chain reaction causing lipid peroxidation	6
6	Structure of glutathione	6
7	Glutathione redox cycle; 1: Glutathione peroxidase, 2: Glutathione reductase, 3: G-6PD	7
8	Role of glutathione reductase	8
9	The action of acetylcholinesterase ( AChE)	9
10	Structure features of the neuromuscular junction	10
11	Structure of Gingerol and Shogaol	11
12	Ginger powder	15
13	Selection of the optimal dose of methyl parathion	16
14	Selection of the optimal dose of ginger diet	16
15	Percent reduction of ABTS by Trolox	15
16	The TBARS assay standard curve for lipid peroxidation	19
17	The GSH standard curve	20
18	The BCA assay standard curve for protein	21
19	Scheme of histological methods	24
20	Standard working sheet of Hippocratic screening test	32

FIGURE	PAGE
21 Red blood cell AChE levels in male rats exposed to 0.5, 1.0 and 1.5 mg/kg methyl parathion.	35
22 Red blood cell AChE levels in female rats exposed to 0.5, 1.0 and 1.5 mg/kg methyl parathion.	35
23 Serum TBARS concentration (MDA) in male rats exposed to 0.5, 1.0 and 1.5 mg/kg methyl parathion.	36
24 Serum TBARS concentration (MDA) in female rats exposed to 0.5, 1.0 and 1.5 mg/kg methyl parathion.	36
25 Red blood cell GSH content in male rats exposed to 0.5, 1.0 and 1.5 mg/kg methyl parathion.	37
26 Red blood cell GSH content in female rats exposed to 0.5, 1.0 and 1.5 mg/kg methyl parathion.	37
27 Liver GST activity in male rats exposed to 0.5, 1.0 and 1.5 mg/kg methyl parathion.	38
28 Liver GST activity in female rats exposed to 0.5, 1.0 and 1.5 mg/kg methyl parathion	38
29 Liver GR activity in male rats exposed to 0.5, 1.0 and 1.5 mg/kg methyl parathion.	39
30 Liver GR activity in female rats exposed to .5, 1.0 and 1.5 mg/kg methyl parathion.	39
31 Red blood cell AChE levels in male rats exposed to 0.5, 1.0 and 1.5%W/W ginger diet.	43
32 Red blood cell AChE levels in female rats exposed to 0.5, 1.0 and 1.5%W/W ginger diet.	43
33 Serum TBARS concentration (MDA) in male rats exposed to 0.5, 1.0 and 1.5%W/W ginger diet.	44

FIGURE	PAGE
34 Serum TBARS concentration (MDA) in female rats exposed to 0.5, 1.0 and 1.5%W/W ginger diet.	44
35 Red blood cell GSH content in male rats exposed to 0.5, 1.0 and 1.5%W/W ginger diet.	45
36 Red blood cell GSH content in female rats exposed to 0.5, 1.0 and 1.5%W/W ginger diet.	45
37 Liver GST activity in male rats exposed to 0.5, 1.0 and 1.5%W/W ginger diet.	46
38 Liver GST activity in female rats exposed to 0.5, 1.0 and 1.5%W/W ginger diet.	46
39 Liver GR activity in male rats exposed to 0.5, 1.0 and 1.5%W/W ginger diet.	47
40 Liver GR activity in female rats exposed to 0.5, 1.0 and 1.5%W/W ginger diet.	47
41 Red blood cell AChE level in male and female rats treated with 1.0 mg/kg methyl parathion and 1.5 %W/W ginger diet for 4 weeks.	50
42 Serum TBARS concentration (MDA) in male and female rats treated with 1.0 mg/kg methyl parathion and 1.5 %W/W ginger diet for 4 weeks.	50
43 Red blood cell GSH content in male and female rats treated with 1.0 mg/kg methyl parathion and 1.5 %W/W ginger diet for 4 weeks.	51


FIGURE		PAGE
44	Liver GST activity in male and female rats treated with 1.0 mg/kg methyl parathion and 1.5 %W/W ginger diet for 4 weeks.	51
45	Liver GR activity in male and female rats treated with 1.0 mg/kg methyl parathion and 1.5 %W/W ginger diet for 4 weeks.	52
46	The number of motor endplate positive AChE activity in muscle of male and female rats treated with 1.0 mg/kg methyl parathion and 1.5 %W/W ginger diet for 4 weeks.	54
47	Normal motor endplate showing positive AChE activity in muscle of rat treated with normal diet and 1.5%W/W ginger diet	55
48	Low AChE activity of motor endplate in muscle of rat treated with 1.0 mg/kg methyl parathion and 1.5%W/W ginger diet plus with 1.0 mg/kg methyl parathion	56
49	Percent reduction of ABTS by powder ginger	57
50	Percent reduction of ABTS by 1.5%W/W ginger diet	58

### ABBREVIATION AND SYMBOLS

A	Absorbance
ABTS	2,2'-azino-bis-(3-ethylbenzthiazoline-6-sulphonic acid)
Ach	Acetylcholine
°C	Degree celsius
cm <sup>2</sup>	Square centimeter
CDNB	1-chloro-2,4-dinitrobenzoic acid
AChE	Acetylcholinesterase
ε	Extinction coefficient
g	Gram
GR	Glutathione reductase
GSH	Glutathione
GSSG	Oxidized glutathione
GST	Glutathione-S-transferase
i.p.	Intraperitoneal
kg	Kilogram
L	Litre
LD50	Lethal dose 50
mg	Milligram
ml	Millilitre
mM	Millimolar
MP	Methyl parathion
nm	Nanometer
No.	Number
OFR	Oxygen free radical
%	Percentage



%W	Percent weight
TBARS	Thiobarbituric acid reactive substance
U	Unit



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่  
Copyright© by Chiang Mai University  
All rights reserved