

## REFERENCES

- Adaskaveg J.E., Blanchette, R.A. and Gilbertson, R.L. (1991). Decay of date palm wood by white-rot and brown-rot fungi. *Canadian Journal of Botany* 69: 615–629.
- Aggarwal R., Tewari, A.K., Srivastava, K.D. and Singh, D.V. (2004). Role of antibiosis in the biological control of spot blotch (*Cochliobolus sativus*) of wheat by *Chaetomium globosum*. *Mycopathologia* 157: 369–377.
- Agrios, G.N. (1988). *Plant Pathology*. Academic Press, San Diego, USA.
- Ahlick, K. and Sieber, T.N. (1996). The profusion of dark septate endophytic fungi in non-mycorrhizal fine roots of forest trees and shrubs. *New Phytologist* 132: 259–270.
- Ahrazem, O., Gómez-Miranda, B., Prieto, A., Bernabé, M. and Leal, J.A. (2000). Heterogeneity of the genus *Myrothecium* as revealed by cell wall polysaccharides. *Archives of Microbiology* 173: 296–302.
- Allen, M. (1993). *The Ecology of Mycorrhizae*. Cambridge Press, Cambridge, UK.
- Andrews, J.H., Hecht, E.P. and Bashirian, S. (1985). Association between the fungus *Acremonium curvulum* and Eurasian water milfoil, *Myriophyllum spicatum*. *Canadian Journal of Botany* 60: 1216–1221.
- Anononymous (1995). *JMP<sup>®</sup> Statistics and Graphics Guide, Version 3.1*. SAS Institute, Cary, NC, USA.

- Apisariyakul, A., Vanittanakom, N. and Buddhasukh, D. (1995). Antifungal activity of turmeric oil extracted from *Curcuma longa* (Zingiberaceae). *Journal of Ethnopharmacology* 49: 163–169.
- Araujo, A. and Ward, O.P. (1990). Extracellular mannanases and galactanases from selected fungi. *Journal of Industrial Microbiology* 6: 171–178.
- Arnold, A.E. and Herre, E.A. (2003). Canopy cover and leaf age affect colonization by tropical fungal endophytes: ecological pattern and process in *Theobroma cacao* (Malvaceae). *Mycologia* 95: 388–398.
- Arnold, A.E., Mejia, L.C., Kyllö, D., Rojas, E.I., Maynard, Z., Robbins, N. and Herre, E.A. (2003). Fungal endophytes limit pathogen damage in a tropical tree. *PNAS* 100: 15649–15654.
- Asuyama, H. (1965). Morphology, taxonomy, host range, and life cycle of *Piricularia oryzae*. In R.F.Jr. Chandler (ed.), *The Rice Blast Disease*, pp. 9–22. The Johns Hopkins Press, USA.
- Atlas, R.M. and Bartha, R. (1998). *Microbial Ecology, Fundamentals and Applications, 4<sup>th</sup> Edition*. Benjamin/Cummings Science Publishing, California, USA.
- Azevedo, J.L., Maccheroni, W.Jr., Pereira, J.O. and de Araújo, W.L. (2000). Endophytic microorganisms: a review on insect control and recent advances on tropical plants. *Electronic Journal of Biotechnology* 3: 40–65.

- Baayen, R.P., Bonants, P.J.M., Verkley, G., Carroll, G.C., van der Aa, H.A., de Weerd, M., van Brouwershaven, I.R., Schutte, G.C., Maccheroni, W.Jr., Glienke de Blanco, C. and Azevedo, J.L. (2002). Nonpathogenic isolates of the citrus black spot fungus, *Guignardia citricarpa*, identified as a cosmopolitan endophyte of woody plants, *G. mangiferae* (*Phyllosticta capitalensis*). *Phytopathology* 92: 464–477.
- Bacon, C.W. (1993). Abiotic stress tolerances (moisture, nutrients) and photosynthesis in endophyte infected tall fescue. *Agriculture, Ecology and Environment* 44: 123–141.
- Bacon, C.W. and de Battista, J. (1991). Endophytic fungi of grasses. In D.K. Arora, B. Rai, K.G. Mukerji and G.R. Knudsen (eds.), *Handbook of Applied Mycology: Soils and Plants*, pp. 231–256. Marcel Dekker, New York, USA.
- Bacon, C.W. and Hinton, D.M. (1996). Symptomless endophytic colonization of maize by *Fusarium moniliforme*. *Canadian Journal of Botany* 74: 1195–1202.
- Bacon, C.W., Bennett, R.M., Hinton, D.M. and Voss, K.A. (1992). Scanning electron microscopy of *Fusarium moniliforme* within asymptomatic corn kernel and kernels associated with equine leukoencephalomalacia. *Plant Disease* 76: 144–148.
- Bailey, A.G. and van Eijnatten, C. (1961). Corn gray spot caused by *Pyricularia grisea*. *Phytopathology* 51: 197–198.
- Bailey, J.A. and Jeger, M.J. (1992). *Colletotrichum: Biology, Pathology and Control*. CAB International, Wallingford, UK.

- Baimai, V. (2002). Biodiversity. In V. Baimai (ed.), *The Human Evolution and Biodiversity*, pp. 103–120. Jirawat Express, Bangkok, Thailand.
- Barklund, P. and Kowalski, T. (1996). Endophytic fungi in branches of Norway spruce with particular reference to *Tryblidiopsis pinastri*. *Canadian Journal of Botany* 74: 673–678.
- Barrow, J.R. (2004). Unique characteristics of a systemic fungal endophyte of native grasses in arid southwestern rangelands. *USDA Forest Service Proceedings RMRS-P-31*: 54–56.
- Bateman, R.P, Carey, M., Moore, D. and Prior, C. (1993). The enhanced infectivity of *Metarhizium flavoviride* in oil formulations to desert locusts at low humidities. *Annals of Applied Biology* 122: 145–152.
- Baura, G., Szaro, T.M. and Bruns, T.D. (1992). *Gastrosporella laricinus* is a recent derivative of *Suillus grevillei*: molecular evidence. *Mycologia* 84: 592–597.
- Bayman, B.P., Lebron, L.L., Tremblay, R.L. and Lodge, D.J. (1997). Variation in endophytic fungi from roots and leaves of *Lepanthes* (Orchidaceae). *New Phytologist* 135: 143–149.
- Beena, K.R., Ananda, K. and Sridhar, K.R. (2000). Fungal endophytes of three sand dune plant species of west coast of India. *Sydowia* 52: 1–9.
- Benhamou, N. and Brodeur, J. (2000). Evidence for antibiosis and induced host defense reactions in the interaction between *Verticillium lecanii* and *Penicillium digitatum*, the causal agent of green mold. *Phytopathology* 68: 4044–4060.

- Berbee, M.L. and Taylor, J.W. (2001). Fungal molecular evolution: gene trees and geologic time. In D.J. McLaughlin, E.G. McLaughlin and P.A. Lemke (eds.), *Systematics and Evolution, The Mycota VIIb*, pp. 229–245. Springer-Verlag, Berlin, Germany.
- Bernstein, M.E. and Carroll, G.C. (1977). Internal fungi in old growth Douglas fir foliage. *Canadian Journal of Botany* 55: 644–653.
- Bertoni, M.D. and Cabral, D. (1988). Phyllosphere of *Eucalyptus viminalis*. II: Distribution of endophytes. *Nova Hedwigia* 46: 491–502.
- Bettucci, L. and Saravay, M. (1993). Endophytic fungi of *Eucalyptus globulus*: a preliminary study. *Mycological Research* 97: 679–682.
- Bills, G., Dombrowski, A., Pelaez, F., Polishook, J. and An, Z. (2002). Recent and future discoveries of pharmacologically active metabolites from tropical fungi. In R. Watling, J.C. Frankland, A.M. Ainsworth, S. Issac and C.H. Robinson (eds.), *Tropical Mycology: Micromycetes, Volume 2*, pp. 165–194. CABI Publishing, New York, USA.
- Bills, G.F. (1996). Isolation and analysis of endophytic fungal communities from woody plants. In S.C. Redlin and L.M. Carris (eds.), *Endophytic Fungi in Grasses and Woody Plants. Systematics, Ecology and Evolution*, pp. 31–65. APS Press, St. Paul, Minnesota, USA.
- Bills, G.F. and Polishook, J.D. (1991). Microfungi from *Carpinus caroliniana*. *Canadian Journal of Botany* 69: 1477–1482.
- Bills, G.F. and Polishook, J.D. (1992). Recovery of endophytic fungi from

*Chamaecyparis thyroides*. *Sydowia* 44: 1–12.

Bills, G.F. and Polishook, J.D. (1994a). Abundance and diversity of microfungi in leaf litter of a lowland rain forest in Costa Rica. *Mycologia* 86: 187–198.

Bills, G.F. and Polishook, J.D. (1994b). Microfungi from decaying leaves of *Heliconia mariae* (Heliconiaceae). *Brenesia* 41–42: 27–43.

Bills, G.F., Rossman, A.Y. and Polishook, J.D. (1994). Rediscovery of *Albosynnema elegans* and *Solheimia costaspora*. *Sydowia* 46: 1–11.

Bissegger, M. and Sieber, T.N. (1994). Assemblages of endophytic fungi in coppice shoots of *Castanea sativa*. *Mycologia* 86: 648–655.

Blodgett, J.T., Swart, W.J., Louw, S.M. and Weeks, W.J. (2000). Species composition of endophytic fungi in *Amaranthus hybridus* leaves, petioles stems, and roots. *Mycologia* 92: 853–859.

Boddy, L. and Watkinson, S.C. (1995). Wood decomposition, higher fungi, and their role in nutrient redistribution. *Canadian Journal of Botany* 73: 1377–1383.

Bohn, M. (1993). *Myrothecium groenlandicum* sp. nov., a presumed endophytic fungus of *Betula nana* (Greenland). *Mycotaxon* 46: 335–341.

Bonde, M.R., Peterson, G.L., Emmett, R.W. and Menge, J.A. (1991). Isozyme comparisons of *Septoria* isolates associated with citrus in Australia and the United States. *Phytopathology* 81: 517–521.

Bonte, F., Noel-Hudson, M.S., Wepierre, J. and Meybeck, A. (1997). Protective effect of curcuminoids on epidermal skin cells under free oxygen radical stress. *Planta Medica* 63: 265–266.

- Borromeo, E.S., Nelson, R.J., Bonman, J.M. and Leung, H. (1993). Genetic differentiation among isolates of *Pyricularia* infecting rice and weed hosts. *Phytopathology* 83: 393–399.
- Braun, U. (2001). Taxonomic notes on some species of the *Cercospora* complex (VII). *Fungal Diversity* 8: 41–71.
- Breen, J.P. (1993). Enhanced resistance to three species of aphids (Homoptera: Aphididae) in *Acremonium* endophyte-infected turf grasses. *Journal of Economic Entomology* 86: 1279–1286.
- Breen, J.P. (1994). *Acremonium* endophyte interactions with enhanced plant resistance to insects. *Annual Reviews of Entomology* 39: 401–423.
- Brown, G.E. (1975). Factors affecting post-harvest development of *Colletotrichum gloeosporioides* in citrus fruits. *Phytopathology* 65: 404–409.
- Brown, K.B., Hyde, K.D. and Guest, D.I. (1998). Preliminary studies on endophytic fungal communities of *Musa acuminata* species complex in Kong Kong and Australia. *Fungal Diversity* 1: 27–51.
- Brown, K.B., Johnson, G.I. and Guest, D.I. (2003). Interactions between the endophytic fungi of durian (*Durio zibethinus*) and *Phytophthora palmivora*. *Proceedings of the 8<sup>th</sup> International Congress of Plant Pathology*, p. 41. Christchurch, New Zealand.
- Brunner, F. and Petrini, O. (1992). Taxonomic studies of *Xylaria* species and xylariaceous endophytes by isozyme electrophoresis. *Mycological Research* 96: 723–733.

- Bruns, T.D., White, T.J. and Taylor, J.W. (1991). Fungal molecular systematics. *Annual Reviews of Ecology and Systematics* 22: 525–564.
- Bryan, G.T., Daniels, M.J. and Osbourn, A.E. (1995). Comparison of fungi within the *Gaeumannomyces-Phialophora* complex by analysis of ribosomal DNA sequences. *Applied and Environmental Microbiology* 61: 681–689.
- Bu'lock, J.D. (1980). Mycotoxins as secondary metabolites. In P.S. Steyn (ed.), *The Biosynthesis of Mycotoxins, A Study in Secondary Metabolism*, pp. 1–16. Academic Press, New York, USA.
- Buchert, J., Salminen, M., Siika-aho, M., Ranua, M. and Viikari, L. (1993). The role of *Trichoderma reesei* xylanase and mannanase in the treatment of softwood kraft pulp prior to bleaching. *Holzforschung* 47: 473–478.
- Bultman, T.L., Borowicz, K.L., Schneble, R.M., Coudron, T.A. and Bush, L.P. (1997). Effect of fungal endophyte on the growth and survival of two *Euplectrus* parasitoids. *Oikos* 78: 170–176.
- Burt, B.L. and Smith, R.M. (1972). Tentative keys to the subfamilies, tribes and genera of the Zingiberales. *Notes from the Royal Botanic Garden Edinburgh* 31: 171–176.
- Bush, L.P., Wilkinson, H.H. and Schardl, C.L. (1997). Bioprotective alkaloids of grass-fungal endophyte symbioses. *Plant Physiology* 114: 1–7.
- Bussaban, B., Lumyong, P., McKenzie, E.H.C., Hyde, K.D. and Lumyong, S. (2002). Index of fungi described from the Zingiberaceae. *Mycotaxon* 83: 165–182.



- Bussaban, B., Lumyong, P., McKenzie, E.H.C., Hyde, K.D. and Lumyong, S. (2003a). *Xenosporium amomi* sp. nov. from Zingiberaceae in Thailand. *Fungal Diversity* 14: 61–66.
- Bussaban, B., Lumyong, P., McKenzie, E.H.C., Hyde, K.D. and Lumyong, S. (2004). Fungi on Zingiberaceae (ginger). In E.B.G. Jones, M. Tantichareon and K.D. Hyde (eds.), *Thai Fungal Diversity*, pp. 189–195. BIOTEC, Bangkok, Thailand.
- Bussaban, B., Lumyong, P., Seelanan, T., Park, D.C., McKenzie, E.H.C., Hyde, K.D. and Lumyong, S. (2005). Molecular and morphological characterization of *Pyricularia* and allied genera. *Mycologia* (In Press).
- Bussaban, B., Lumyong, S., Lumyong, P., Hyde, K.D. and McKenzie, E.H.C. (2001a). Two new species of endophytes (ascomycetes) from Zingiberaceae sporulating in culture. *Nova Hedwigia* 73: 487–493.
- Bussaban, B., Lumyong, S., Lumyong, P., Hyde, K.D. and McKenzie, E.H.C. (2003b). Three new species of *Pyricularia* isolated as zingiberaceous endophytes from Thailand. *Mycologia* 95: 519–524.
- Bussaban, B., Lumyong, S., Lumyong, P., McKenzie, E.H.C. and Hyde, K.D. (2001b). Endophytic fungi from *Amomum siamense*. *Canadian Journal of Microbiology* 47: 943–948.
- Bussaban, B., Lumyong, S., Lumyong, P., McKenzie, E.H.C. and Hyde, K.D. (2001c). A synopsis of the genus *Berkleasmium* with two new species and new records of *Canalisporium caribense* from Zingiberaceae in Thailand. *Fungal Diversity* 8: 73–85.

- Bussaban, B., Lumyong, S., Lumyong, P., McKenzie, E.H.C. and Hyde, K.D. (2001d). Endophytic fungi from Zingiberaceae: *Alpinia malaccensis*. *Phytopathology* 91: S104. Publication no. P-2001-0023-MSA.
- Cabral, D. (1985). Phyllosphere of *Eucalyptus viminalis*: dynamic of fungal populations. *Transactions of the British Mycological Society* 85: 501–511.
- Cabral, D., Stone, J.K. and Carroll, G.C. (1993). The internal mycobiota of *Juncus* spp.: microscopic and cultural observations of infection patterns. *Mycological Research* 97: 367–376.
- Cai, L., Tsui, C.K.M., Zhang, K. and Hyde, K.D. (2002). Aquatic fungi from Lake Fuxian, Yunnan, China. *Fungal Diversity* 9: 57–70.
- Callac, P. and Guinberteau, J. (2005). Morphological and molecular characterization of two novel species of *Agaricus* section *Xanthodermatei*. *Mycologia* 97: 416–424.
- Campbell, N.A., Reece, J.B., Mitchell, L.G. and Taylor, M.R. (2003). *Biology: Concepts and Connections, 4<sup>th</sup> Edition*. Benjamin/Cummings Publishing, USA.
- Campin, J. (2004). Guide to plant relationships, for food allergy and intolerance identification. <http://www.purr.demon.co.uk/Food/RelatedPlantList.html> [2005, February 13].
- Cannon, P.F. (1997). Strategies for rapid assessment of fungal diversity. *Biodiversity and Conservation* 6: 669–680.
- Cantoria, M.C. (2003). Asian herbal drugs from the family Zingiberaceae. *Proceedings of the 3<sup>rd</sup> Symposium on the Family Zingiberaceae*, pp. 208–221. Khon Kaen, Thailand,

- Cappellano, A., de Quartre, B., Valla, G. and Moiroudo, A. (1987). Root nodule formation by *Penicillium* sp. on *Alnus glutinosa* and *Alnus incana*. *Plant and Soil* 104: 45–51.
- Carbone, I. and Kohn, L.M. (1993). Ribosomal DNA sequence divergence within internal transcribed spacers 1 of the Sclerotiniaceae. *Mycologia* 85: 415–427.
- Carmichael, J.W., Kendrick, W.B., Connors, I.L. and Sigler, L. (1980). *Genera of Hyphomycetes*. The University of Alberta Press, Alberta, Canada.
- Carroll, F.E., Müller, E. and Sutton, B.C. (1977). Preliminary studies on the incidence of needle endophytes in some European conifers. *Sydowia* 29: 87–103.
- Carroll, G.C. (1986). The biology of endophytism in plants with particular reference to woody perennials. In N.J. Fokkema and J. van der Heavel, (eds.), *Microbiology of the Phylloplane*, pp. 205–222. Cambridge University Press, London, UK.
- Carroll, G.C. (1988). Fungal endophytes in stem and leaves: from latent pathogen to mutualistic symbiont. *Ecology* 69: 2–9.
- Carroll, G.C. and Carroll, F.E. (1978). Studies on the incidence of coniferous needle endophytes in the Pacific Northwest. *Canadian Journal of Botany* 56: 3034–3043.
- Carroll, G.C. and Wicklow, D.T. (1992). *The Fungal Community. Its Organization and Role in the Ecosystem*. Marcel Dekker, New York, USA.
- Cavara, F. (1892). Contribuzione alla micologia lombarda. *Attica Information Sciences Technology of Botanical University Laboratory (Crittogam Pavia)* 2:

207–292.

- Chaiprasert, A. (2004). Dermatophytes and pythiosis. In E.B.G. Jones, M. Tantichareon and K.D. Hyde (eds.), *Thai Fungal Diversity*, pp. 213–225. BIOTEC, Bangkok, Thailand.
- Chand, P., Aruna, A., Maqsood, A.M. and Rao, L.V. (2005). Novel mutation method for increased cellulase production. *Journal of Applied Microbiology* 98: 318–323.
- Chang, H.S. (1998). Three dematiaceous hyphomycetes from Taiwan. *Botanical Bulletin of Academic Sinica* 40: 247–250.
- Chapela, I.H. and Boddy, L. (1989). Fungal colonization of attached beech branches I. Early stages of development of fungal communities. *New Phytologist* 110: 39–45.
- Chapela, I.H. and Boddy, L. (1989). Fungal colonization of attached beech branches II. Spatial and temporal organization of communities arising from latent invaders in bark and functional sapwood, under different moisture regimes. *New Phytologist* 110: 47–57.
- Chase, M.W., Soltis, D.E., Soltis, P.S., Rudall, P.J., Fay, M.F., Hahn, W.H., Sullivan, S., Joseph, J., Molvray, M., Kores, P.J., Givnish, T.J., Sytsma, K.J. and Pires, J.C. (2000). Higher-level systematics of the monocotyledons: an assessment of current knowledge and a new classification. In K.L. Wilson and D.A. Morrison (eds.), *Monocots: Systematics and Evolution*, pp. 3–16. CSIRO Publishing, Collingwood, Australia.

- Cheeptham, N., Higasshiyama, T., Fukushi, E., Matsuura, H., Mikama, T., Yokota, T., Ichihara, A. and Tomita, F. (1999). Studies in an antifungal antibiotic from *Ellisiodothis ingiunans* L1588-A8. *Thai journal of Biotechnology* 1: 37–45.
- Chen, D., Zeigler, R.S., Leung, H. and Nelson, R.J. (1995). Population structure of *Pyricularia grisea* at two screening sites in the Philippines. *Phytopathology* 85: 1011–1020.
- Chen, Y.X., Wei, G. and Chen, W.P. (2002). New species of *Pestalotiopsis*. *Mycosystema* 21: 316–323.
- Cheplick, G.P. and Clay, K. (1988). Acquired chemical defences in grasses: The role of fungal endophytes. *Oikos* 52: 309–318.
- Chivers, A.H. (1915). A monograph of the genera *Chaetomium* and *Ascotricha*. *Memoirs of the Torrey Botanical Club* 14: 1–240
- Choi, Y.W., Hyde, K.D. and Ho, W.H. (1999). Single spore isolation of fungi. *Fungal Diversity* 3: 29–38.
- Chuakul, W. (2003). Ethnomedical uses of Thai zingiberaceous plants. *Proceedings of the 3<sup>rd</sup> Symposium on the Family Zingiberaceae*, pp. 201–207. Khon Kaen, Thailand.
- Clay, K. (1986). Grass endophytes. In N.J. Fokkema and J. van den Heuvel (eds.), *Microbiology of the Phyllosphere*. Cambridge University Press, Cambridge, UK, pp. 188–204.
- Clay, K. (1989). Clavicipitaceous endophytes of grasses: their potential as biocontrol agents. *Mycological Research* 92: 1–12.

- Clay, K. (1992) Fungal endophytes of plants: biological and chemical diversity. *Natural Toxins* 1: 147–149.
- Concepcion, G.P., Lazaro, J.E. and Hyde, K.D. (2001). Screening for bioactive novel compounds. In S.B. Pointing and K.D. Hyde (eds.), *Bio-Exploitation of Filamentous Fungi*, pp. 93–130. Fungal Diversity Press, Hong Kong.
- Cook, R.C. and Rayner, A.D.M. (1984). *Ecology of Saprophytic fungi*. Longman Group Ltd., Essex, UK.
- Couch, B.C. and Kohn, L.M. (2002). A multilocus gene genealogy concordant with host preference indicates segregation of a new species, *Magnaporthe oryzae*, from *M. grisea*. *Mycologia* 94: 683–693.
- Cuevas, W.A., Kantelinen, A., Tanner, P., Bodie, B. and Leskinen, S. (1996). Purification and characterization of novel mannanase used in pulp bleaching. In E. Srebotnik and K. Messner (eds.), *Biotechnology in The Pulp and Paper Industry*, pp. 123–126. Facultas-Universitätsverlag, Vienna, Austria.
- D'Souza, M., Singh, S.K. and Bhat, D.J. (2002). New conidial fungi from Andaman Islands, India. *Mycotaxon* 82: 133–143.
- Dahlgren, R.M.T., Clifford, H.T. and Yeo, P.F. (1985). *The Families of the Monocotyledons: Structure, Evolution and Taxonomy*. Springer-Verlag Berlin Heidelberg, Germany.
- Dahlman, D.L., Eichenseer, H. and Siegel, M.R. (1991). Chemical perspectives on endophyte-grass interactions and their implications to insect herbivory. In P. Barbosa, V.A. Krischik and C.G. Jones (eds.), *Microbial Mediation of Plant-*

- Herbivore Interactions*, pp. 227–252. John Wiley and Sons, New York, USA.
- de Barry, A. (1887). *Comparative Morphology and Biology of the Fungi, Mycetozoa and Bacteria*. Oxford, UK.
- de Hoog, G.S. (1985). Taxonomy of the *Dactylaria* complex, IV. *Dactylaria*, *Neta*, *Subulispora* and *Scolecobasidium*. *Studies in Mycology* 26: 1–60.
- de Hoog, G.S. and von Arx, J.A. (1973). Revision of *Scolecobasidium* and *Pleurophragmium*. *Kavaka* 1: 55–60.
- Denyer, C.V., Jackson, P. and Loakes, D.M. (1994). Isolation of antirhinoviral sesquiterpenes from ginger (*Zingiber officinale*). *Journal of Natural Products* 57: 658–662.
- Didukh, M., Vilgalys, R., Wasser, S.P., Isikhuemhen, O.S. and Nevo, E. (2005). Notes on *Agaricus* section *Duploannulati* using molecular and morphological data. *Mycological Research* 109: 729–740.
- Dillard, H.R. (1992). *Colletotrichum coccodes*: the pathogen and its hosts. In J.A. Bailey and M.J. Jeger, *Colletotrichum: Biology, Pathology and Control*, pp. 225–236. CAB International, Wallingford, UK.
- Dingle, J. and McGee, P.A. (2003). Some endophytic fungi reduce the density of pustules of *Puccinia recondita* f. sp. *tritici* in wheat. *Mycological Research* 107: 301–316.
- Dix, N.J. and Webster, J. (1995). *Fungal Ecology*. Chapman and Hall, London, UK.
- Dodd, J.C., Estrada, A. and Jeger, M.J. (1992). Epidemiology of *Colletotrichum gloeosporioides* in the tropics. In J.A. Bailey and M.J. Jeger (eds.), *Colletotrichum: Biology, Pathology and Control*, pp. 308–325. CAB

International, Wallingford, UK.

Doi, Y. (1977). *Protocreopsis*, a new genus of the Hypocreales. *Kew Bulletin* 31: 511–555.

Downie, B., Holhorst, H.W.M. and Bewley, J.D. (1994). A new assay for quantifying endo-beta-D-mannanase activity using Congo red dye. *Phytochemistry* 36: 829–835.

Dreyfuss, M. and Petrini, O. (1984). Further investigations on the occurrence and distribution of endophytic fungi in tropical plants. *Botanica Helvetica* 94: 33–40.

Dreyfuss, M.M. and Chapela, I.H. (1999). Potential of fungi in the discovery of novel, low-molecular weight pharmaceuticals. In V.C. Gullo (ed.), *The Discovery of Natural Products with Therapeutic Potential*, pp. 49–80. Butterworth-Heinemann, UK.

Edds, G.T. (1979). *Aflatoxins. Conference on Mycotoxins in Animal Feeds and Grain Related to Animal Health*, pp. 80–164. National Technical Information Service, Springfield, Virginia.

Edwards, M.J. (1988). *ATCC Microbes and Cells at Work: an Index to ATCC Strains with Special Applications*. American Type Culture Collections, Rockville, Maryland, USA.

Egger, K.N. (1995). Molecular analysis of ectomycorrhizal fungi communities. *Canadian Journal of Botany* 73: S1415–S1422.

Egger, K.N. and Sigler, L. (1993). Relatedness of the ericoid endophytes *Scytalidium vaccinii* and *Hymenoscyphus ericae* inferred from analysis of ribosomal DNA.



*Mycologia* 85: 219–230.

Elad, Y. and Kapat, A. (1999). The role of *Trichoderma harzianum* protease in the biocontrol of *Botrytis cinerea*. *European Journal of Plant Pathology* 105: 177–189.

Ellis, M.B. (1971). *Dematiaceous Hyphomycetes*. Commonwealth Mycological Institute, Kew, UK.

Ellis, M.B. (1976). *More Dematiaceous Hyphomycetes*. Commonwealth Mycological Institute, Kew, UK.

Espinosa-Gracia, F.J. and Longenheim, J.H. (1990). The leaf fungal endophytic community of a coastal red wood population—diversity and spatial patterns. *New Phytologist* 116: 89–98.

Evans, H.C., Fröhlich, J. and Shamoun, S.F. (2001). Biological control of weeds. In S.B. Pointing and K.D. Hyde (eds.), *Bio-Exploitation of Filamentous Fungi, Fungal Diversity Research Series 6*, pp. 349–401. Fungal Diversity Press, Hong Kong.

Farr, D.F., Bills, G.F., Chamuris, G.P. and Rossman, A.Y. (1989). *Fungi on Plants and Plant Products in the United States*. APS Press, Minnesota, USA.

Felsenstein, J. (1985). Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39: 783–791.

Ficker, C.E., Smith, M.L., Susiarti, S., Leaman, D.J., Irawati, C. and Arnason, J.T. (2003). Inhibition of human pathogenic fungi by members of Zingiberaceae used by the Kenyah (Indonesian Borneo). *Journal of Ethnopharmacology* 85: 289–293.

- Fisher, P.J. and Petrini, O. (1987). Location of fungal endophytes in tissues of *Suaeda fruticosa*: a preliminary study. *Transactions of the British Mycological Society* 89: 246–249.
- Fisher, P.J. and Petrini, O. (1990). A comparative study of fungal endophytes in xylem and bark of *Alnus* species in England and Switzerland. *Mycological Research* 94: 313–319.
- Fisher, P.J. and Petrini, O. (1992). Fungal saprobes and pathogens as endophytes of rice (*Oryza sativa* L.) *New Phytologist* 120: 137–143.
- Fisher, P.J., Anson, A.E. and Petrini, O. (1984). Novel antibiotic activity of an endophyte *Cryptosporiopsis* sp. isolated from *Vaccinium myrtillus*. *Transactions of the British Mycological Society* 83: 145–148.
- Fisher, P.J., Anson, A.E. and Petrini, O. (1986). Fungal endophytes in *Ulex europaeus* and *Ulex gallii*. *Transactions of the British Mycological Society* 86: 153–156.
- Fisher, P.J., Graf, F., Petrini, L.E., Sutton, B.C. and Wookey, P.A. (1995a). Fungal endophytes of *Dryas octopelata* from a high polar semidesert and from the Swiss Alps. *Mycologia* 81: 319–323.
- Fisher, P.J., Petrini, L.E., Sutton, B.C. and Petrini, O. (1995b). A study of fungal endophytes in leaves, stems and roots of *Gynoxis oleifolia* Muchler (Compositae) from Ecuador. *Nova Hedwigia* 60: 589–594.
- Fisher, P.J., Petrini, O. and Lappin-Scott, H.M. (1992). The distribution of some fungal and bacterial endophytes in maize (*Zea mays* L.). *New Phytologist* 122: 299–305.

- Fisher, P.J., Petrini, O. and Sutton, B.C. (1993). A comparative study of fungal endophytes in leaves, xylem and bark of *Eucalyptus* in Australia and England. *Sydowia* 45: 338–345.
- Fisher, P.J., Petrini, O., Petrini, L.E. and Sutton, B.C. (1994). Fungal endophytes from the leaves and twigs of *Quercus ilex* L. from England, Majorca, and Switzerland. *New Phytologist* 127: 133–137.
- Fox, F.M. (1993). Tropical fungi: their commercial potential. In S. Isaac, J.C. Frankland, R. Watling and A.J.S. Whalley (eds.), *Aspects of Tropical Mycology*, pp. 253–264. Cambridge University Press, Cambridge, UK.
- Frankland, J.C. (1992). Mechanisms in fungal succession. In G.C. Carroll and D.T. Wicklow (eds.), *The Fungal Community. Its Organization and Role in the Ecosystem, 2<sup>nd</sup> Edition*, pp. 383–401. Marcel Dekker, New York, USA.
- Franklin, T.J. and Snow, G.A. (2005). *Biochemistry and Molecular Biology of Antimicrobial Drug Action, 6<sup>th</sup> Edition*. Springer-Verlag, Berlin, Germany.
- Fravel, D.R. (1988). Role of antibiosis in the biocontrol of plant diseases. *Annual Reviews of Phytopathology* 26: 75–91.
- Freeman, S. and Rodriguez, R.J. (1993). Genetic conversion of a fungal plant pathogen to a non-pathogenic, endophytic mutualism. *Science* 260: 75–78.
- Fridborg, H., Jonsson, E., Nygren, P. and Larsson, R. (1999). Relationship between diagnosis-specific activity of cytotoxic of a human tumor colony-forming assay to new drug screening. *Cancer Research* 45: 2145–2153.

- Fröhlich, J. and Hyde, K.D. (1999). Biodiversity of palm fungi in the tropics: are global fungal diversity estimates realistic? *Biodiversity and Conservation* 8: 977–1004.
- Fröhlich, J., Hyde, K.D. and Petrini, O. (2000). Endophytic fungi associated with palms. *Mycological Research* 104: 1202–1212.
- Funk, C.R., Halisky, P.M., Johnson, M.C., Siegel, M.R., Stewart, A.V., Ahmad, S., Hurley, R.H. and Harvey, I.C. (1983). An endophytic fungus and resistance to sod webworms: association in *Lolium perenne*. *Bio/Technology* 1: 189–191.
- Gajaseneni, J. and Boonpragob, K. (1995). Methods for measurement of species diversity. In T.J.B. Boyle and B. Boontawee (eds.), *Measuring and Monitoring Biodiversity in Tropical and Temperate Forests*, pp. 303–307. Center for International Forestry Research, Indonesia.
- Gale, E.F., Cundliffe, E., Reynolds, P.E., Richmond, M.H. and Waring, M.J. (1981). *The Molecular Basis of Antibiotic Action, 2<sup>nd</sup> Edition*. Wiley & Sons, New York, USA.
- Gams, W. (2000). *Phialophora* and some similar morphologically little-differentiated anamorphs of divergent ascomycetes. *Studies in Mycology* 45: 187–199.
- Gardes, M. and Bruns, T.D. (1991). Rapid characterization of ectomycorrhizae using RFLP pattern of their PCR amplified-ITS. *Mycological Society Newsletter* 41: 14.
- George, M.L.C., Nelson, R.J., Zeigler, R.S. and Leung, H. (1998). Rapid population analysis of *Magnaporthe grisea* by using rep-PCR and endogenous repetitive

DNA sequences. *Phytopathology* 88: 223–229.

Goh, T.K. and Hyde, K.D. (1997). A revision of *Dactylaria*, with description of *D. tunicata* sp. nov. from submerged wood in Australia. *Mycological Research* 101: 1265–1272.

Goodwin, S.B., Levy, M., Cavaletto, J.R. and Tian, Y. (2003). Phylogenetic analyses place *Magnaporthe grisea* in the order Diaporthales and suggest rapid speciation. *Abstracts 8<sup>th</sup> International Congress of Plant Pathology*, p. 349. Christchurch, New Zealand.

Gravesen, S., Frisvad, J.C., Samson, R.A. and Munksgaard. (1994). *Microfungi*. Special-Trykkeriet Viborgals, Denmark.

Greuter, W. (2000). *International Code of Botanical Nomenclature (Saint Louis Code)*. International Association for Plant Taxonomy (Europe), Germany.

Griffiths, M. (1992). *Index of Garden Plants*. Timber Press, England.

Guadet, J., Julien, J., Lafay, J.F. and Brygoo, Y. (1989). Phylogeny of some *Fusarium* species as determined by large-subunit rRNA sequence comparison. *Molecular Biology and Evolution* 6: 227–242.

Guarro, J., Soler, L. and Rinaldi, M.G. (1995). Pathogenicity and antifungal susceptibility of *Chaetomium* species. *European Journal of Clinical Microbiology & Infectious Diseases* 14: 613–618.

Gübitz, G.M., Haltrich, D., Latal, B. and Steiner, W. (1997b). Mode of depolymerisation of hemicellulose by various mannanases and xylansases in relation to their ability to bleach softwood pulp. *Applied Microbiology and Biotechnology* 47: 658–662.

- Gübitz, G.M., Lischmig, T., Stebbing, D. and Saddler, J.N. (1997a). Enzymatic removal of hemicellulose from dissolving pulps. *Biotechnology Letters* 19: 491–495.
- Guo, L.D., Hyde, K.D. and Liew, E.C.Y. (1998). A method to promote sporulation in palm endophytic fungi. *Fungal Diversity* 1: 109–113.
- Guo, L.D., Hyde, K.D. and Liew, E.C.Y. (2000). Identification of endophytic fungi from *Livistona chinensis* based on morphology and rDNA sequences. *New Phytologist* 147: 617–630.
- Hajek, A.E., Wraight, S.P. and Vandenberg, J.D. (2001). Control of arthropods using pathogenic fungi. In S.B. Pointing and K.D. Hyde (eds.), *Bio-Exploitation of Filamentous Fungi, Fungal Diversity Research Series 6*, pp. 309–347. Fungal Diversity Press, Hong Kong.
- Halmschlager, E., Butin, H. and Donaubaue, E. (1993). Endophytische Pilze in Blättern und Zweigen von *Quercus pertraea*. *European Journal of Forest Pathology* 23: 51–63.
- Halpern, G.M. and Miller, A.H. (2002). *Medicinal Mushrooms, Ancient Remedies for Modern Ailments*. M. Evans and Company, New York, USA.
- Hammerli, U.A., Brandle, U.E., Petrini, O. and McDermott, J.M. (1992). Differentiation of isolates of *Discula umbrinella* (teleomorph : *Apiognomonina errabunda*) from beech, chestnut and oak using RAPD markers. *Molecular Plant-Microbe Interactions* 5: 479–483.
- Hammond, K.E. and Faeth, S.H. (1992). Ecology of plant-herbivore communities: a

fungal component? *Natural Toxins* 1: 197–208.

Hammond, P.M. (1992). Species inventory. In B. Groombridge (ed.), *Global Biodiversity, Status of the Earth's Living Resources*, pp. 17–39. Chapman & Hall, London, UK.

Hashioka, Y. (1971). Notes on *Pyricularia* I. Three species parasitic to Musaceae, Cannaceae and Zingiberaceae. *Transactions of the Mycological Society of Japan* 12: 126–135.

Hashioka, Y. (1973). Notes on *Pyricularia* II. Four species and one variety parasitic to Cyperaceae, Gramineae and Commelinaceae. *Transactions of the Mycological Society of Japan* 14: 256–265.

Hata, K. and Futai, K. (1993). Effect of the needle aging on the total colonization rates of *Pinus thunbergii* and *Pinus densiflora* needles. *Journal of Japanese Forest Society* 75: 338–341.

Hata, K. and Futai, K. (1995). Endophytic fungi associated with healthy pine needles and needles infested by the pine needle gall midge, *Thecodiplosis japonensis*. *Canadian Journal of Botany* 73: 384–390.

Hata, K., Atari, R. and Sone, K. (2002). Isolation of endophytic fungi from leaves of *Pasania edulis* and their within-leaf distributions. *Mycoscience* 43: 369–373.

Hawksworth, D.L. (1991). The fungal dimension of biodiversity: magnitude, significance, and conservation. *Mycological Research* 95: 641–655.

Hawksworth, D.L. (2001). The magnitude of fungal diversity: the 1.5 million species estimate revised. *Mycological Research* 105: 1422–1432.

- Hawksworth, D.L., Kirk, P.M., Sutton, B.C. and Pegler, D.N. (1995). *Ainsworth and Bisby's Dictionary of the Fungi, 8<sup>th</sup> Edition*. CAB International, Wallingford, UK.
- Hawksworth, D.L., Minter, D.W., Kinsey, J. and Cannon, P.F. (1996). Inventorying a tropical fungal biota: intensive and extensive approaches. In K.K. Janardhanan, C. Rajendran, K. Natacajan and D.L. Hawksworth (eds.), *Tropical Mycology*. Oxford & IBH Publishing, UK.
- Hebert, T.T. (1971). The perfect stage of *Pyricularia grisea*. *Phytopathology* 61: 83–87.
- Heywood, V.H. (1993). *Flowering Plants of the World*. Oxford University Press, UK.
- Hibbett, D.S. (1992). Ribosomal RNA and fungal systematics. *Transactions of the Mycological Society of Japan* 33: 533–556.
- Higashide, E., Asai, M., Ootsu, K., Tanida, S., Kozai, Y., Hasegawa, T., Kishi, T., Sugino, Y. and Yoneda, M. (1977). Ansamitocin, a group of novel maytansinoid antibiotics with antitumour properties from *Nocardia*. *Nature (London)* 270: 721–722.
- Higuchi, T., Saika, T., Senda, S., Mizobata, T., Kawata, Y. and Nagai, J. (1997). Development of biorational pest control formulation against longicorn beetles using a fungus, *Beauveria brongniartii* (Sacc.) Petch. *Journal of Fermentation and Bioengineering* 84: 236–243.



- Hirsch, G. and Braun, U. (1992). Communities of parasitic microfungi. In W. Winterhoff (ed.), *Handbook of Vegetation Science, Volume 19, Fungi in Vegetation Science*, pp. 225–250. Kluwer Academic, Dordrecht, The Netherlands.
- Ho, W.H., Yanna, Hyde, K.D. and Hodgkiss, I.J. (2002). Seasonality and sequential occurrence of fungi on wood submerged in Tai Po Kau Forest Stream, Hong Kong. *Fungal Diversity* 10: 21–43.
- Holdenrieder, O. and Sieber, T.N. (1992). Fungal associations of serially washed healthy non-mycorrhizal roots. *Mycological Research* 96: 151–156.
- Hooker, J.D. (1856). Decades LI.–LIV. Rio Negro fungi. *Hooker Journal of Botany* 8: 129–144.
- Hughes, S.J. (1958). Revisiones hyphomycetum aliquot cum appendice de nominibus rejiciendis. *Canadian Journal of Botany* 36: 727–836.
- Hyde, K.D. (1995). Diversity of microfungi in the wet tropics of north Queensland. In T.J.B. Boyle and B. Boontawee (eds.), *Measuring and Monitoring Biodiversity in Tropical and Temperate Forests*, pp. 271–286. Center for International Forestry Research, Indonesia.
- Hyde, K.D. (1997). Additions to the genus *Linocarpon* (Ascomycetes: Hyponectriaceae). *Botanical Journal of the Linnean Society* 123: 109–131.
- Hyde, K.D. (2001). Increasing the likelihood of novel compound discovery from filamentous fungi. In S.B. Pointing and K.D. Hyde (eds.), *Bio-Exploitation of Filamentous Fungi, Fungal Diversity Research Series 6*, pp. 77–92. Fungal

Diversity Press, Hong Kong.

Hyde, K.D. and Alias, S.A. (2000). Biodiversity and distribution of fungi associated with decomposing *Nypa fruticans*. *Biodiversity and Conservation* 9: 393–402.

Hyde, K.D. and Hawksworth, D.L. (1997). Measuring and monitoring the biodiversity of microfungi, In K.D. Hyde (ed.), *Biodiversity of Tropical Microfungi*, pp. 11–28. Hong Kong University Press, Hong Kong.

Hyde, K.D., Farrant, C.A. and Jones, E.B.G. (1987). Isolation and culture of marine fungi. *Botanica Marina* 30: 291–303.

Hyde, K.D., Fröhlich, J. and Taylor, J. (1997). Diversity of ascomycetes on palms in the tropics. In K.D. Hyde (ed.), *Biodiversity of Tropical Microfungi*, pp. 141–156. Hong Kong University Press, Hong Kong.

Hyde, K.D., Ho, W.H., McKenzie, E.H.C. and Dalisay, T. (2001). Saprobic fungi on bamboo culms. *Fungal Diversity* 7: 35–48.

Hyde, K.D., Taylor, J.E. and Fröhlich, J. (2000). *Genera of Ascomycetes from Palm, Fungal Diversity Research Series 2*. Fungal Diversity Press, Hong Kong.

Hyde, K.D., Wong, S.W., Lumyong, S. and Lumyong, P. (1997). *Apiosordaria striatispora*, an endophyte of *Mesua ferrea* and *Prunus arborea* from Thailand. *Mycoscience* 38: 437–439.

Hyde, K.D., Zhou, D.Q. and Dalisay, T.E. (2002a). Bambusicolous fungi: a review. *Fungal Diversity* 9: 1–14.

Hyde, K.D., Zhou, D.Q., McKenzie, E.H.C., Ho, W.H. and Dalisay, T.E. (2002b). Vertical distribution of saprobic fungi on bamboo culms. *Fungal Diversity* 11:

109–118.

Hywel-Jones, N.L. and Boonpratuang, T. (2001). *An updated preliminary checklist of fungi recorded from Thailand*. BIOTEC database.

Ingold, C.T. (1943). Further observations on aquatic hyphomycetes of decaying leaves. *Transactions of the British Mycological Society* 26: 105–115.

Ingold, C.T. and Hudson, H. (1993). *The Biology of Fungi, 6<sup>th</sup> Edition*. Chapman & Hall, London, UK.

Iwabe, N., Kuma, K., Hasegawa, M., Osawa, S. and Miyata, T. (1989). Evolutionary relationship of archaeobacteria, eubacteria and eukaryotes inferred from phylogenetic trees of duplicated genes. *Proceedings of the National Academy of Science USA* 86: 9355–9359.

Jackson, A.J., Walters, D.R. and Marshall, G. (1991). Biological control of chocolate spot on faba beans. *Aspects Applied Biology* 27: 117–122.

Jackson, A.J., Walters, D.R. and Marshall, G. (1997). Antagonistic interactions between the foliar pathogen *Botrytis fabae* and isolates of *Penicillium brevicompactum* and *Cladosporium cladosporioides* on faba beans. *Biological Control* 8: 97–106.

Jennings, D.H. and Lysek, G. (1996). *Fungal Biology: Understanding the Fungal Lifestyle*. BIOS Scientific Publishers, Oxford, UK.

Johnson, J.A. and Whitney, N.J. (1989). A study of fungal endophytes of needles of balsam fir (*Abies balsamea*) and red spruce (*Picea rubens*) in New Brunswick, Canada, using culture and electron microscope techniques. *Canadian Journal of Botany* 67: 3513–3516.

- Johnson, J.A. and Whitney, N.J. (1992). Isolation of fungal endophytes from black spruce (*Picea mariana*) dormant buds and needles from New Brunswick, Canada. *Canadian Journal of Botany* 70: 1754–1757.
- Jones, E.B.G. and Hyde, K.D. (2004). Introduction to Thai fungal diversity. In E.B.G. Jones, M. Tantichareon and K.D. Hyde (eds.), *Thai Fungal Diversity*, pp. 7–35. BIOTEC, Bangkok, Thailand.
- Jones, E.B.G., Wong, S.W., Sivichai, S., Au, D.W.T. and Hywel-Jones, N.L. (1999). Lignicolous freshwater Ascomycota from Thailand: *Micropeltopsis quinquecladiopsis* sp. nov. *Mycological Research* 103: 729–735.
- Jong, S.C. and Birmingham, J.M. (1993). Microbial diversity and its role in sustainable agriculture. *Proceeding of the Chinese American Academic and Profession Convention, United for a Prosperous 21<sup>st</sup> Century*, pp. 9.29–6.32.
- Jorquensen, J.H., Turnidge, J.D. and Washington, J.A. (1999). Antibacterial susceptibility test: Dilution and disk diffusion methods. In P.R. Murnay (ed.), *Manual of Clinical Microbiology, 7<sup>th</sup> Edition*, pp. 1526–1543. ASM Press, Washington DC, USA.
- Kar, A.K. and Mandal, M. (1969). New *Cercospora* spp. from West Bengal. I. *Transactions of the British Mycological Society* 53: 337–360.
- Kato, H., Yamaguchi, T. and Nishihara, N. (1976). The perfect stage of *Pyricularia oryzae* Cav. in culture. *Annals of the Phytopathological Society of Japan* 42: 507–510.

- Kato, H., Yamamoto, M., Yamaguchi-Ozaki, T., Kadouchi, H., Iwamoto, Y., Nakayashiki, H., Tosa, Y., Mayama, S. and Mori, N. (2000). Pathogenicity, mating ability and DNA restriction fragment length polymorphisms of *Pyricularia* populations isolated from Gramineae, Bambusideae and Zingiberaceae plants. *Journal of General Plant Pathology* 66: 30–47.
- Kawai, T., Kinoshita, K., Koyama, K. and Takahashi, K. (1994). Anti-emetic principles of *Magnolia abovata* bark and *Zingiber officinale* rhizome. *Planta Medica* 60: 17–20.
- Khurana, I.P.S. (1980). The Clavariaceae of India. XIV. The genus *Typhula*. *Mycologia* 72: 708–727.
- Killham, K. (1994). *Soil Ecology*. Cambridge University Press, Cambridge, UK.
- Kim, W.K., Mauthe, W., Hausner, G. and Klassen, G.R. (1990). Isolation of high molecular weight DNA and double-stranded RNAs from fungi. *Canadian Journal of Botany* 68: 1898–1902.
- Kirk, O., Borchert, T.V. and Claus Crone Fuglsang, C.C. (2002). Industrial enzyme applications. *Current Opinion in Biotechnology* 13: 345–351.
- Kirk, P.M., Cannon, P.F., David, J.C. and Stalpers, J.A. (2001). *Dictionary of the Fungi, 9<sup>th</sup> Edition*. CAB International, Surrey, UK.
- Kirk, T.K., Burgess, R.R. and Koning, J.W.Jr. (1993). Use of fungi in pulping wood: an overview of biopulping research. In G.F. Leatham (ed.), *Frontiers in Industrial Mycology*, pp. 99–111. Chapman & Hall, New York, USA.

- Kobayashi, M., Kanasaki, R., Sato, I., Abe, F., Nitta, K., Ezaki, M., Sakamoto, K., Hashimoto, M., Fujie, A., Hino, M. and Hori, Y. (2005). FR207944, an antifungal antibiotic from *Chaetomium* sp. No. 217 I. Taxonomy, fermentation and biological properties. *Bioscience, Biotechnology and Biochemistry* 69: 515–521.
- Kodsueb, R., Lumyong, S and Hyde, K.D. (2004). Terrestrial lignicolous microfungi. In E.B.G. Jones, M. Tantichareon and K.D. Hyde (eds.), *Thai Fungal Diversity*, pp. 155–161. BIOTEC, Bangkok, Thailand.
- Koo, K.L.K., Ammit, A.J., Tran, V.H., Duke, C.C. and Roufogalis, B.D. (2001). Gingerols and related analogues inhibit arachidonic acid-induced human platelet serotonin release and aggregation. *Thrombosis Research* 103: 3873–3897.
- Kotani, S. and Kurata, M. (1992). Black blotch of rhizome by *Pyricularia zingiberi* Nishikado. *Annals of the Phytopathological Society of Japan* 58: 469–472.
- Kowalski, T. and Kehr, R.D., (1992). Endophytic fungal colonization of branch bases in several forest tree species. *Sydowia* 44: 137–168.
- Kress, W.J. (1990). The phylogeny and classification of the Zingiberales. *Annals of Missouri Botanical Garden* 77: 698–721
- Kress, W.J. (1995). Phylogeny of the Zingiberanae: morphology and molecules. In PJ Rudall, PJ Cribb, DF Cutler and CJ Humphries (eds.), *Monocotyledons: Systematics and Evolution*, pp. 443–460. Royal Botanic Gardens, Kew, UK.
- Kress, W.J., Prince, L.M. and Williams, K.J. (2002). The phylogeny and a new classification of the gingers (Zingiberaceae): evidence from molecular data.

*American Journal of Botany* 89: 1682–1696.

Kress, W.J., Prince, L.M., Hahn, W.J. and Zimmer, E.A. (2001). Unraveling the evolutionary radiation of the families of the Zingiberales using morphological and molecular evidence. *Systematic Biology* 51: 926–944.

Kuhls, K., Lieckfeldt, E., Samuels, G.J., Meyer, W., Kubicek, C.P. and Börner, T. (1997). Revision of *Trichoderma* sect. *Longibrachiatum* including related teleomorphs based on analysis of ribosomal DNA internal transcribed spacer sequences. *Mycologia* 89: 442–460.

Kulik, M.M. (1984). Symptomless infection, persistence, and production of pycnidia in host and non-host plants by *Phomopsis batatae*, *P. phaseoli* and *P. sojae* and the taxonomic implications. *Mycologia* 76: 274–291.

Kulik, M.M. (1988). Observations by scanning electron and bright field microscopy on the mode of penetration of soy bean seedlings by *Phomopsis phaseoli*. *Plant Disease* 72: 115–118.

Kumaresan, V. and Suryanarayanan, T.S. (2001). Occurrence and distribution of endophytic fungi in a mangrove community. *Mycological Research* 105: 1388–1391.

Kumaresan, V. and Suryanarayanan, T.S. (2002). Endophytic assemblages in young, mature and senescent leaves of *Rhizophora apiculata*: evidence for the role of endophytes in mangrove litter degradation. *Fungal Diversity* 9: 81–91.

Kupchan, S.M., Komoda, Y., Court, W.A., Thomas, G.J., Smith, R.M., Karim, A., Gilmore, C.J., Haltiwanger, R.C. and Bryan, R.F. (1972). *Journal of the American Chemistry Society* 94: 1354–1356.

- Lambert, P.W. (1983). Industrial enzyme production and recovery from filamentous fungi. In J.E. Smith, D.R. Berry and B. Kristiansen (eds.), *The Filamentous Fungi, Volume 4. Fungal Technology*, pp. 210–237. Arnold, London, UK.
- Larone, D.H. (1995). *Medically Important Fungi: a Guide to Identification, 3<sup>rd</sup> Edition*. ASM Press, Washington D.C., USA.
- Larsen, K. (1980). Annotated key to the genera of Zingiberaceae of Thailand. *National History Bulletin of Siam Society* 28: 151–169.
- Larsen, K. (1996). A preliminary checklist of the Zingiberaceae of Thailand. *Thai Forest Bulletin (Botany)* 24: 35–49.
- Larsen, K., Lock, J.M., Mass, S. and Mass, P.J.M. (1998). Zingiberaceae. In K. Kubitzki (ed.), *The Families and Genera of Vascular Plants, Volume 4*, pp. 474–495. Springer-Verlag, Berlin, Germany.
- Latch, G.C.M. (1993). Physiological interactions of endophytic fungi and their hosts. Biotic stress tolerance imparted to grasses by endophytes. *Agriculture, Ecology and Environment* 44: 143–156.
- Lebrun, M.H., Capy, M.P., Garcia, N., Dutertre, M., Brygoo, Y., Notteghem, J.L. and Vales, M. (1991). Biology and genetics of *Pyricularia oryzae* and *P. grisea* populations: current situation and development of RFLP markers. *Proceedings of the 2<sup>nd</sup> International Rice Genetics Symposium*, pp. 487–497. IRRI, Los Banos, Laguna, The Philippines.



- Lechat-Vahirua, I., Francois, P., Menut, C., Lamaty, G. and Bessiere, J.M. (1993). Aromatic plants of French Polynesia. I. Constituents of the essential oils of rhizomes of three Zingiberaceae: *Zingiber zerumbet* Smith, *Hedychium coronarium* Koenig and *Etilingera cevuga* Smith. *Journal of Essential Oil Research* 5: 55–59.
- Lee, S.B. and Taylor, J.W. (1992). Phylogeny of five fungus-like protoctistan *Phytophthora* species, inferred from the internal transcribed spacers of ribosomal DNA. *Molecular Biology and Evolution* 9: 636–653.
- Legault, D., Dessureault, M. and Laflamme, G. (1989). Mycoflore des aiguilles de *Pinus banksiana* et *Pinus resinosa*. I. Champignons endophytes. *Canadian Journal of Botany* 67: 2056–2060.
- Lenné, J.M. (1992). *Colletotrichum* diseases of legumes. In J.A. Bailey and M.J. Jeger, *Colletotrichum: Biology, Pathology and Control*, pp. 134–166. CAB International, Wallingford, UK.
- Leuchtman, A. (1994). Isozyme characterization, persistence and compatibility of fungal and grass mutualists. In C.W. Bacon and J.Jr. White (eds.), *Biotechnology of Endophytic Fungi of Grasses*, pp. 21–33. CRC Press, Boca Raton, FL, USA.
- Leuchtman, A. and Clay, K. (1990). Isozyme variation in the *Acremonium/Epichloë* fungal endophyte complex. *Phytopathology* 80: 1133–1139.
- Leuchtman, A., Petrini, O., Petrini, L.E. and Carroll, G.C. (1992). Isozyme polymorphism in six endophytic *Phyllosticta* species. *Mycological Research* 96:

287–294.

Levy, M., Ramao, J., Marchetti, M.A. and Hamer, J.E. (1991). DNA fingerprinting with a dispersed repeated sequence resolves pathotype diversity in the rice blast fungus. *Plant Cell* 3: 95–102.

Li, J.Y., Harper, J.K., Grant, D.M., Tombe, B.O., Bashyal, B., Hess, W.M. and Strobel, G.A. (2001). Ambuic acid, a highly functionalized cyclohexenone with antifungal activity from *Pestalotiopsis* spp. and *Monochaetia* sp. *Phytochemistry* 56: 463–468.

Li, J.Y., Sidhu, R.S., Ford, E.J., Long, D.M., Hess, W.M. and Strobel, G.A. (1998). The induction of taxol production in the endophytic fungus, *Periconia* sp. from *Torreya grandifolia*. *Journal of Industrial Microbiology and Biotechnology* 20: 259–264.

Li, J.Y., Strobel, G.A., Harper, J.K., Lobkovsky, E. and Clardy, J. (2000). Cryptocin, a potent tetramic acid antimycotic from the endophytic fungus *Cryptosporiopsis cf. quercina*. *Organic Letters* 2: 767–770.

Li, W.H. (1997). Rates and patterns of nucleotide substitution. In W.H. Li (ed.), *Molecular Evolution*, pp. 177–213. Sinauer Associates, Sunderland, Massachusetts, USA.

Liou, G.Y. and Tzean, S.S. (1997). Phylogeny of the genus *Arthrobotrys* and allied nematode-trapping fungi based on rDNA sequences. *Mycologia* 89: 876–884.

Liu, C.H., Zou, W.X., Lu, H. and Tan, R.X. (2001). Antifungal activity of *Artemisia annua* endophyte against phytopathogenic fungi. *Journal of Biotechnology* 88: 277–282.

- Lodge, D.J., Fisher, P.J. and Sutton, B.C. (1996). Endophytic fungi of *Manilkara bidentata* leaves in Puerto Rico. *Mycologia* 88: 733–738.
- Lumyong, S., Lumyong, P., Boontim, N. and Hyde, K.D. (1998). *Distribution of Endophytic Fungi Among Indigenous Plant Species in Doi Suthep-Pui Area*. Research Report, Chiang Mai University, Thailand.
- Lumyong, S., Thongantha, S., Lumyong, P. and Tomita, F. (2000). Endophytic fungi from 13 bamboo species in Thailand. *Biotechnology for Sustainable Utilization of Biological Resources in the Tropics* 14: 96–101.
- Luo, J., Yin, J.F., Cai, L., Zhang, K.Q. and Hyde, K.D. (2004). Freshwater fungi in Lake Dianchi, a heavily polluted lake in Yunnan, China. *Fungal Diversity* 16: 93–112.
- Luz, A.I.R., Zoghbi, M.G.B., Ramos, L.S., Maia, J.G.S. and Silva, M.L. (1984). Essential oils of some Amazonian Zingiberaceae, III. Genera *Alpinia* and *Renealmia*. *Journal of Natural Products* 47: 907–908.
- Lynch, J.M. and Poole, N.J. (1979). *Microbial Ecology: a Conceptual Approach*. J. Wiley, New York, USA.
- Mabberley, D.J. (1987). *The Plant-Book: a Portable Dictionary of the Higher Plants*. Cambridge University Press, Cambridge, UK.
- Mabberley, D.J. (1992). *The Plant-Book: a Portable Dictionary of the Vascular Plants, 2<sup>nd</sup> edition*. Cambridge University Press, Cambridge, UK.
- Madigan, M.T., Martinko, J.M. and Parker, J. (2003). *Brock Biology of Microorganisms, 10<sup>th</sup> Edition*. Pearson Education.

- Malca, M.I. and Owen, J.H. (1957). The gray leaf spot disease of St. Augustine grass. *Plant Disease Reporter* 41: 874–875.
- Manner, J.M., Masel, A., Braithwaite, K.S. and Irwin, J.A.G. (1992). Molecular analysis of *Colletotrichum gloeosporioides* pathogenic on the tropical pasture legume *Stylosathes*. In J.A. Bailey and M.J. Jeger (eds.), *Colletotrichum: Biology, Pathology and Control*, pp. 250–268. CAB International, Wallingford, UK.
- Marshall, D., Tunali, B. and Nelson, L.R. (1999). Occurrence of fungal endophytes in species of wild *Triticum*. *Crop Science* 39: 1507–1512.
- Marvanová, L. and Descals, E. (1987). New taxa and new combinations of ‘aquatic hyphomycetes’. *Transactions of the British Mycological Society* 89: 499–507.
- Marvanová, L. and Hywel-Jones, N.L. (2000). *Sigmoidea contorta* sp. nov. and two rare species from waters in Thailand. *Cryptogamie Mykologie* 21: 13–26.
- Marx, D.H., Maul, S. B. and Cordell, C.E. (1993). Application of specific ectomycorrhizal fungi in world forestry. In G.F. Leatham (ed.), *Frontiers in Industrial Mycology*, pp. 78–98. Chapman & Hall, New York, USA.
- Matsushima, T. (1995). *Matsushima Mycological Memoirs* 8. Matsushima, Kobe, Japan.
- Matsuyama, N., Kato, H. and Yamaguchi, T. (1977). Comparison of the isozyme patterns of extracellular enzymes in *Pyricularia* strains from gramineous and zingiberaceous plants. *Annals of the Phytopathological Society of Japan* 43: 419–425.

- May, R.M. (1991). A fondness for fungi. *Nature* 352: 475–476.
- May, R.M. (1994). Conceptual aspects of the quantification of the extent of biological diversity. *Philosophical Transactions of the Royal Society of London, Section B* 345: 13–20.
- McCullough, M.J., Clemons, K.V., McCusker, J.H. and Stevens, D.A. (1998). Intergenic transcribed spacer PCR ribotyping for differentiation of *Saccharomyces* species and interspecific hybrids. *Journal of Clinical Microbiology* 36: 1035–1038.
- McKenzie, E.H.C. and Hyde, K.D. (1997). Microfungi on Pandanaceae. In K.D. Hyde (ed.), *Biodiversity of Tropical Microfungi*, pp. 157–177. Hong Kong University Press, Hong Kong.
- McKenzie, E.H.C., Whitton, S.R. and Hyde, K.D. (2002). The Pandanaceae: does it have a diverse and unique fungal biota? In R. Watling, J.C. Franklin, A.M. Ainsworth, S. Isaac and C.H. Robinson (eds.), *Tropical Mycology, Volume 2, Micromycota*, pp. 51–61. CAB International, Wallingford, UK.
- McLaughlin, D.J., McLaughlin, E.G. and Lemke, P.A. (2001). *Systematics and Evolution, The Mycota VIIa*. Springer-Verlag, Berlin, Germany.
- Menendez, A., Bertoni, M.D. and Cabral, D. (1995). Comparative study of occurrence of fungal endophytes in *Juncus* species of Argentina. *Nova Hedwigia* 60: 583–588.
- Menkis, A., Allmer, J., Vasiliauskas, R., Lygis, V., Stenlid, J. and Finlay, R. (2004). Ecology and molecular characterization of dark septate fungi from roots, living

stems, coarse and fine woody debris. *Mycological Research* 108: 965–973.

Meredith, D.S. (1963). *Pyricularia grisea* (Cooke) Sacc. causing pitting disease of bananas in central America. I. Preliminary studies on pathogenicity. *Annals of Applied Biology* 52: 453–463.

Millör, C.S. (1980). Infection processes on conifer needles. In J.P. Blakeman (ed.), *Microbial Ecology of the Phylloplane*, pp. 185–209. Academic Press, London, UK.

Miyoshi, N., Nakamura, Y., Ueda, Y., Abe, M., Ozawa, Y., Uchida, K. and Osawa, T. (2003). Dietary ginger constituents, galanals A and B, are potent apoptosis inducers in Human T lymphoma Jurkat cells. *Cancer Letters* 199: 113–119.

Moon, C.D., Scott, B., Schardl, C.L. and Christensen, M.J. (2000). The evolutionary origins of *Epichloë* endophytes from annual ryegrasses. *Mycologia* 92: 1103–1118.

Moore, D. and Chiu, S.W. (2001). Fungal products as food. In S.B. Pointing and K.D. Hyde (eds.), *Bio-Exploitation of Filamentous Fungi, Fungal Diversity Research Series 6*, pp. 223–251. Fungal Diversity Press, Hong Kong.

Morin, L., Hill, R.L. and Matayoshi, S. (1997). Hawaii's successful biological control strategy for mist flower (*Ageratina riparia*), can it be transferred to New Zealand? *Biocontrol News and Information* 18: 77–88.

Morris, E.F. (1967). Studies on the synnematosus fungi imperfecti II. *Mycopathologica et Mycologia Applicata* 33: 179–185.

Mullis, K.B. and Faloona, F.A. (1987). Specific synthesis of DNA *in vitro* via a polymerase-catalyzed chain reaction. *Methods in Enzymology* 155: 335–350.

- Murray, P.R., Baron, E.J., Pfaller, M.A., Tenover, F.C. and Tenover, R.H. (1999). *Manual of Clinical Microbiology 7<sup>th</sup> Edition*. ASM Press, Washington DC, USA.
- Muthappa, B.N. (1996). A new species of *Sphaceloma* on cardamom from India. *Sydowia* 19: 143–145.
- Nag Raj, T.R. (1995). What is *Myrothecium presonii*? *Mycotaxon* 53: 295–310.
- Narisawa, K., Currah, R.S. and Hashiba, T. (2003). The root endophytic fungus *Phialocephala fortinii* suppresses *Verticillium* yellow in Chinese cabbage. *Proceedings of the 8<sup>th</sup> International Congress of Plant Pathology*, p. 39. Christchurch, New Zealand.
- Newman, J.A., Abner, M.L., Dado, R.G., Gibson, D.J., Brookings A. and Parsons, A.J. (2003). Effects of elevated CO<sub>2</sub>, nitrogen and fungal endophyte-infection on tall fescue: growth, photosynthesis, chemical composition and digestibility. *Global Change Biology* 9: 425–437.
- Nicholas, K.B. and Nicholas, H.B.Jr. (1997). *GeneDoc: a Tool for Editing and Annotating Multiple Sequence Alignments*. <http://www.psc.edu/biomed/genedoc> [2002, August 5].
- Nicholson, R.L. (1992). *Colletotrichum graminicola* and the anthracnose diseases of maize and sorghum. In J.A. Bailey and M.J. Jeger, *Colletotrichum: Biology, Pathology and Control*, pp. 186–202. CAB International, Wallingford, UK.
- Nicklin, J., Graeme-Cook, K., Paget, T. and Killington, R. (1999). *Instant Notes in Microbiology*. BIOS Scientific Publishers, UK.

- Nisbet, L.J. and Fox, M.F. (1991). The importance of microbial diversity to biotechnology. In D.L. Hawksworth (ed.), *The Biodiversity of Microorganisms and Invertebrates: Its Role in Sustainable Agriculture*, pp. 229–244. CAB International, Wallingford, UK.
- Nishikado, Y. (1917). Studies on the rice blast fungus. I. *Berichte Ohara Institute Landwirtschaft Forschung* 1: 171–217.
- O'Donnell, J. and Dickinson, C.H. (1980). Pathogenicity of *Alternaria* and *Cladosporium* isolates on *Phaseolus*. *Transactions of the British Mycological Society* 74: 335–342.
- Ojumu, T.V., Solomon, B.O., Betiku, E., Layokun, S.K. and Amigun, B. (2003). Cellulase production by *Aspergillus flavus* Linn isolate NSPR 101 fermented in sawdust, bagasse and corncob. *African Journal of Biotechnology* 2: 150–152.
- Okane, I., Lumyong, S., Nakagiri, A. and Ito, T. (2003). Extensive host range of an endophytic fungus *Guignardia endophyllicola* (anamorph: *Phyllosticta capitalensis*). *Mycoscience* 44: 353–363.
- Okane, I., Nakagiri, A. and Ito, T. (1998). Endophytic fungi in leaves of ericaceous plants. *Canadian Journal of Botany* 76: 657–663.
- Okane, I., Nakagiri, A. and Ito, T. (2001). Assemblages of endophytic fungi on *Bruguiera gymnorrhiza* in the Shiira River Basin, Iriomote Is. *IFO Research Community* 20: 41–49.
- Ooi, V.E.C. (2001). In S.B. Pointing and K.D. Hyde (eds.), *Bio-Exploitation of Filamentous Fungi, Fungal Diversity Research Series 6*, pp. 191–222. Fungal



Diversity Press, Hong Kong.

Ou, S.H. (1987). *Rice Diseases, 2<sup>nd</sup> Edition*. CAB International, Kew, UK.

Oudemans, P. and Coffey, M.D. (1991). A revised systematics of twelve papillate *Phytophthora* species based on isozyme analysis. *Mycological Research* 95: 1025–1046.

Outtrup, H. and Boyce, C.O.L. (1990). Microbial proteases and biotechnology. In W.M. Fogarty and C.T. Kelly (eds.), *Microbial Enzymes and Biotechnology 2<sup>nd</sup> Edition*. Elsevier, Applied Science, New York, USA.

Pandey, A.K., Reddy, M.S. and Suryanarayanan, T.S. (2003). ITS-RFLP and ITS sequence analysis of foliar endophytic fungus *Phyllosticta* from different tropical trees in India. *Mycological Research* 107: 439–444.

Pappas, A.C. and Paplomatas, E.J. (1998). *Pyricularia* leaf spot: a new disease of ornamental plants of the family Marantaceae. *Plant Disease* 82: 465–469.

Park, J.Y., Okada, G. and Takahashi, M. (2002). Screening of fungi antagonists against yellows of cabbage caused by *Fusarium oxysporum* f. sp. *conglutinans*. *Mycoscience* 43: 447–451.

Paulus, B., Gadek, P. and Hyde, K.D. (2003). Two new species of *Dactylaria* (anamorphic fungi) and an update of species in *Dactylaria sensu lato*. *Fungal Diversity* 14: 157–170.

Pavgi, M.S. and Upadhyay, R. (1986). Some parasitic fungi on turmeric from India. *Sydowia* 21: 100–104.

- Peláez, F., Collado, J., Arenal, F., Basilio, A., Cabello, A., Díez Matas, M.T., García, J.B., González Del Val, A., González, V., Gorrochategui, J., Hernández, P., Martín, I., Platas, G. and Vicente, F. (1998). Endophytic fungi from plants living on gypsum soils as a source of secondary metabolites with antimicrobial activity. *Mycological Research* 102: 755–761.
- Penttilä, M.E., Andre, L., Salheimo, M., Lehtovaara, P. and Knowles, J.K.C. (1987). Expression of two *Trichoderma reesei* endoglucanases in the yeast *Saccharomyces cerevisiae*. *Yeast* 3: 175–185.
- Pereira, J.O., Azevedo, J.L. and Petrini, O. (1993). Endophytic fungi of *Stylosanthes*: a first report. *Mycologia* 85: 362–364.
- Petersen, O.G. (1889). Musaceae, Zingiberaceae, Cannaceae, Marantaceae. In A. Engler and K. Prantl (eds.), *Die Natürlichen Pflanzenfamilien, 1<sup>st</sup> Edition*, pp. 1–43. Verlag von Wilhelm Engelmann, Leipzig, Germany.
- Petrini, L.E. and Petrini, O. (1985). Xylariaceous fungi as endophytes. *Sydowia* 38: 216–234.
- Petrini, O. (1984). Endophytic fungi in British Ericaceae: a preliminary study. *Transactions of the British Mycological Society* 83: 510–512.
- Petrini, O. (1991). Fungal endophytes of tree leaves. In J.H. Andrew and S.S. Hirano (eds.), *Microbial Ecology of Leaves*, p. 179–187. Springer-Verlag, New York, USA.
- Petrini, O. (1993). Endophytes of *Pteridium* spp.: some considerations for biological control. *Sydowia* 45: 330–338.

- Petrini, O. (1996). Ecological and physiological aspects of host specificity in endophytic fungi. In S.C. Redlin and L.M. Carris (eds.), *Endophytic Fungi in Grasses and Woody Plants: Systematics, Ecology and Evolution*, pp. 87–100. APS Press, St. Paul, Minnesota, USA.
- Petrini, O. and Carroll, G.C. (1981). Endophytic fungi in foliage of some Cupressaceae in Oregon. *Canadian Journal of Botany* 59: 629–636.
- Petrini, O. and Fisher, P.J. (1986). Fungal endophytes of *Salicornia perennis*. *Transactions of the British Mycological Society* 91: 233–238.
- Petrini, O. and Fisher, P.J. (1988). A comparative study of fungal endophytes in xylem and whole stem of *Pinus sylvestris* and *Fagus sylvatica*. *Transactions of the British Mycological Society* 87: 647–651.
- Petrini, O. and Fisher, P.J. (1990). Occurrence of fungal endophytes in twigs of *Salix fragilis* and *Quercus robur*. *Mycological Research* 94: 1077–1080.
- Petrini, O. and Müller, E. (1979). Pilzliche Endophyten am Beispiel von *Juniperus communis* L. *Sydowia* 32: 224–251.
- Petrini, O., Fisher, P.J. and Petrini, L.E. (1992a). Fungal endophytes of bracken (*Pteridium aquilinum*), with some reflections on their use in biological control. *Sydowia* 44: 282–293.
- Petrini, O., Sieber, T.N., Toti, L. and Vivet, O. (1992b). Ecology, metabolite production and substrate utilization in endophytic fungi. *Natural Toxins* 1: 185–196.

- Petrini, O., Stone, J. and Carroll, F.E. (1982). Endophytic fungi in evergreen shrubs in western Oregon: a preliminary study. *Canadian Journal of Botany* 60: 789–796.
- Photita, W. (2003). *Survey and Isolation of Endophytic and Saprobiic Fungi of Wild Banana in Doi Suthep-Pui National Park*. Ph.D. Thesis, Chiang Mai University, Thailand.
- Photita, W., Lumyong, P., McKenzie, E.H.C., Hyde, K.D. and Lumyong, S. (2002). A new *Dictyosporium* species from *Musa acuminata* in Thailand. *Mycotaxon* 82: 415–419.
- Photita, W., Lumyong, P., McKenzie, E.H.C., Hyde, K.D. and Lumyong, S. (2003a). *Memnoniella* and *Stachybotrys* species from *Musa acuminata*. *Cryptogamie Mycologie* 24: 147–152.
- Photita, W., Lumyong, P., McKenzie, E.H.C., Hyde, K.D. and Lumyong, S. (2003b). Saprobiic fungi on dead wild banana. *Mycotaxon* 85: 345–356.
- Photita, W., Lumyong, S., Lumyong, P. and Hyde, K.D. (2001a). Fungi on *Musa acuminata* in Hong Kong. *Fungal Diversity* 6: 99–106.
- Photita, W., Lumyong, S., Lumyong, P. and Hyde, K.D. (2001b). Endophytic fungi of wild banana (*Musa acuminata*) at Doi Suthep-Pui National Park. *Mycological Research* 105: 1508–1513.
- Photita, W., Lumyong, S., Lumyong, P., McKenzie, E.H.C. and Hyde, K.D. (2004). Are some endophytes of *Musa acuminata* latent pathogens? *Fungal Diversity* 16: 131–140.

- Piercey-Normore, M.D. and Egger, K.N. (2001). Identification and genotypic characterization of filamentous fungi. In N.J. Talbot (ed.), *Molecular and Cell Biology of Filamentous Fungi: a Practical Approach*, pp. 1–22. Oxford University Press, New York, USA.
- Pinnoi, A. (2004). *Biodiversity and Antifungal Production by Fungi on the Palm Eleiodoxa confera in Siridhorn Peat Swamp Forest, Narathiwat, Thailand*. M.Sc. Thesis, Chiang Mai University, Thailand.
- Pinnoi, A., Jones, E.B.G., McKenzie, E.H.C. and Hyde, K.D. (2003a). Aquatic fungi from peat swamp palms: *Unisetosphaeria penguinoides* gen. et sp. nov., and three new *Dactylaria* species. *Mycoscience* 44: 377–382.
- Pinnoi, A., McKenzie, E.H.C., Jones, E.B.G. and Hyde, K.D. (2003b). Palm fungi from Thailand: *Custingophora undulatistipes* sp. nov. and *Vanakripa minutiellipsoidea* sp. nov. *Nova Hedwigia* 77: 213–219.
- Pinnoi, A., Pinruan, U., Hyde, K.D., McKenzie, E.H.C. and Lumyong, S. (2004). *Submersisphaeria palmae* sp. nov. with a key to species and notes on *Helicoubisia*. *Sydowia* 56: 72–78.
- Pinruan, U., Jones, E.B.G. and Hyde, K.D. (2002). Aquatic fungi from peat swamp palms: *Jahnula appendiculata* sp. nov. *Sydowia* 54: 242–247.
- Pinruan, U., Lumyong, S., McKenzie, E.H.C., Jones, E.B.G. and Hyde, K.D. (2004a). Three new species of *Craspedodidymum* from palm in Thailand. *Mycoscience* 45: 177–180.
- Pinruan, U., McKenzie, E.H.C., Jones, E.B.G. and Hyde, K.D. (2004b). Two new

species of *Stachybotrys*, and a key to the genus. *Fungal Diversity* 17: 145–157.

Pinruan, U., Sakayaroj, J., Jones, E.B.G. and Hyde, K.D. (2004c). Aquatic fungi from peat swamp palms: *Phruensis brunnespora* gen. et. sp. nov. and its hyphomycete anamorph. *Canadian Journal of Botany* 96: 1161–1181.

Pointing, S.B. (1999). Qualitative methods for the determination of lignocellulolytic enzyme production by tropical fungi. *Fungal Diversity* 2: 17–33.

Pointing, S.B. and Hyde, K.D. (2001). *Bio-Exploitation of Filamentous Fungi*. Fungal Diversity Press, Hong Kong.

Polishook, J.D., Dombrowski, A.W., Tsou, N.N., Salituro, G.M. and Curotto, J.E. (1993). Preussomerin D from the endophyte *Hormonema dematioides*. *Mycologia* 85: 62–64.

Poon, M.O.K. and Hyde, K.D. (1998). Evidence for the vertical distribution of saprophytic fungi on senescent *Phragmites australis* at Mai Po Marshes, Hong Kong. *Botanica Marina* 41: 285–292.

Poulsen, A.D. (1996). Species richness and density of ground herbs within a plot of lowland rainforest in north-west Borneo. *Journal of Tropical Ecology* 12: 177–190.

Powell, R.G., Smith, C.R.Jr., Plattner, R.D. and Jones, B.E. (1983). Additional new maytansinoids from *Trewia nudiflora*: 10-epitrewiasine and nortrewiasine. *Journal of Natural Products* 46: 660–666.

Prasada, R. and Goyal, J.P. (1970). A new species of *Pyricularia* on Bajra. *Current Science* 39: 287–288.

- Prompttha, I., Hyde, K.D., Lumyong, P., McKenzie, E.H.C. and Lumyong, S. (2003). *Dokmaia monthadangii* gen. et sp. nov., a synnematos anamorphic fungus on *Manglietia garetii*. *Sydowia* 55: 99–103.
- Prompttha, I., Hyde, K.D., Lumyong, P., McKenzie, E.H.C. and Lumyong, S. (2005a). Fungi on *Magnolia liliifera*: *Cheiromyces magnoliae* sp. nov. from dead branches. *Nova Hedwigia* (In press).
- Prompttha, I., Lumyong, S., Lumyong, P., McKenzie, E.H.C. and Hyde, K.D. (2002). Fungal succession on senescent leaves of *Manglietia garrettii* in Doi Suthep-Pui National Park, Northern Thailand. *Fungal Diversity* 10: 89–100.
- Prompttha, I., Lumyong, S., Lumyong, P., McKenzie, E.H.C. and Hyde, K.D. (2004a). A new species of *Pseudohalonectria* gen. nov. from Thailand. *Cryptogamie Mycologie* 25: 43–47.
- Prompttha, I., Lumyong, S., Lumyong, P., McKenzie, E.H.C. and Hyde, K.D. (2004b). Fungal saprobes on dead leaves of *Magnolia liliifera* (Magnoliaceae) in Thailand. *Cryptogamie Mycologie* 25: 315–321.
- Prompttha, I., Lumyong, S., Lumyong, P., McKenzie, E.H.C. and Hyde, K.D. (2005b). A new species of *Anthostomella* on *Magnolia liliifera* from northern Thailand. *Mycotaxon* (In press).
- Pujol, I., Guarro, J., Llop, C., Soler, L. and Fernandez-Ballart, J. (1996). Comparison study of broth macrodilution and microdilution antifungal susceptibility tests for the filamentous fungi. *Antimicrobial Agents and Chemotherapy* 40: 2106–2110.

- Pulici, M., Sugawara, F., Koshino, H., Okada, G., Esumi, Y., Uzawa, J. and Yoshida, S. (1997). Metabolites of *Pestalotiopsis* spp., endophytic fungi of *Taxus brevifolia*. *Phytochemistry* 46: 313–319.
- Quimio, T.H. and Quimio, A.J. (1975). Notes of Philippine grape and guava anthracnose. *Plant Disease Reporter* 59: 221–224.
- Quintiliani, R., Sahm, D.F. and Courvalin, P. (1999). Mechanisms of resistance to antimicrobial agents. In P.F. Murray, E.J. Baron M.A. Pfaller, F. C. Tenover and R.H. Tenover (eds.), *Manual of Clinical Microbiology*, 7<sup>th</sup> Edition, pp. 1505–1525. ASM Press, Washington DC, USA.
- Ramakrishnan, T.S. (1942). Leaf spot disease of *Zingiber officinale* caused by *Phyllosticta zingiberi* sp. nov. *Proceedings of Indian Academic Science, Section B* 15: 167–171.
- Rao, V.G. (1962). Some new records of fungi-imperfecti from India. *Sydowia* 16: 41–45.
- Rathaiah, Y. (1980). Leaf spot of turmeric. *Plant Disease* 64: 104–105.
- Rathaiah, Y. (1981). A new species of *Cercoseptoria* causing a leaf spot of ginger. *Mycologia* 73: 774–777.
- Raven, P.H., Evert, R.F. and Eichhorn, S.E. (1992). *Biology of Plants*, 5<sup>th</sup> Edition. Worth Publishers, New York, USA.
- Rehner, S.R. and Samuels, G.J. (1994). Taxonomy and phylogeny of *Gliocladium* analyzed by large subunit rDNA sequences. *Mycological Research* 98: 625–634.



- Rehner, S.R. and Samuels, G.J. (1995). Molecular systematics of the Hypocreales: a teleomorph gene phylogeny and the status of their anamorphs. *Canadian Journal of Botany* 73: S816–S823.
- Reissinger, A., Winter, S., Steckelbroeck, S., Hartung, W. and Sikora, R.A. (2003). Infection of barley roots by *Chaetomium globosum*: evidence for a protective role of the exodermis. *Mycological Research* 107: 1094–1102.
- Richardson, K.A. and Currah, R.S. (1995). The fungal community associated with the roots of some rainforest epiphytes of Costa Rica. *Selbyana* 16: 49–73.
- Riesen, T.K. and Close, R.C. (1987). Endophytic fungi in propiconazole-treated and untreated barley leaves. *Mycologia* 79: 546–552.
- Riesen, T.K. and Sieber, T. (1985). *Endophytic Fungi in Winter Wheat (Triticum aestivum L.)*. Mikrobiologisches Institut, Swiss Federal Institute of Technology, Zurich.
- Rinehart, K.L.Jr. and Shield, L.S. (1976). Chemistry of the ansamycin antibiotics. *Fortsch der Chem Org Naturst* 33: 231–307.
- Rodrigues, K.F. (1994). The foliar fungal endophytes of the Amazonian palm *Euterpe oleracea*. *Mycologia* 86: 376–385.
- Rodrigues, K.F. and Petrini, O. (1997). Biodiversity of endophytic fungi in tropical regions. In K.D. Hyde (ed.), *Biodiversity of Tropical Microfungi*, pp. 57–69. Hong Kong University Press, Hong Kong.
- Rodrigues, K.F. and Samuels, G.J. (1990). Preliminary study of endophytic fungi in a tropical palm. *Mycological Research* 94: 827–830.

- Rodrigues, K.F., Hesse, M. and Werner, C. (2000). Antimicrobial activities of secondary metabolites produced by endophytic fungi from *Spondias mombin*. *Journal of Basic Microbiology* 40: 261–267.
- Rodrigues, K.F., Leuthmann, A. and Petrini, O. (1993). Endophytic species of *Xylaria*: cultural and isozymic studies. *Sydowia* 45: 116–138.
- Rodrigues, K.F., Sieber, T.N., Grünig, C.R. and Holdenrieder, O. (2004). Characterization of *Guignardia mangiferae* isolated from tropical plants based on morphology, ISSR-PCR amplifications and ITS1-5.8S-ITS2. *Mycological Research* 108: 45–52.
- Romero, A., Carrión, G. and Rico-Gray, V. (2001). Fungal latent pathogens and endophytes from leaves of *Parthenium hysterophorus* (Asteraceae). *Fungal Diversity* 7: 81–87.
- Rossman, A.Y. (1994). A strategy for an all-taxa inventory of fungal biodiversity. In C.I. Peng and C.H. Chou (eds.), *Biodiversity and Terrestrial Ecosystems*. *Monograph Series No 14*, pp. 169–194. Institute of Botany, Academia Sinica, Taiwan.
- Rossman, A.Y. (1997). Biodiversity of tropical microfungi: an overview. In K.D. Hyde (ed.), *Biodiversity of Tropical Microfungi*, pp. 1–10. Hong Kong University Press, Hong Kong.
- Rossman, A.Y., Farr, D.F. and Castlebury, L.A. (2001). Phylogenetic relationships among obscure microfungi: why does it matter? *Phytopathology* 91: S164. Publication no. P-2001-0119-MSA.

- Rossmann, A.Y., Howard, R.J. and Valent, B. (1990). *Pyricularia grisea*, the correct name for the rice blast disease fungus. *Mycologia* 82: 509–512.
- Rubini, M.R., Silva-Ribeiro, R.T., Pomella, A.W.V., Maki, C.S., Araújo, W.L., dos Santos, D.R. and Azevedo, J.L. (2005). Diversity of endophytic fungal community of cacao (*Theobroma cacao* L.) and biological control of *Crinipellis perniciosus*, causal agent of witches' broom disease. *International Journal of Biological Science* 1: 24–33.
- Sabini, E., Schubert, H., Murshudov, G., Wilson, K.S., Siika-aho, M. and Penttiä, M. (2000). The three-dimensional structure of a *Trichoderma reesei*  $\beta$ -mannanase from glycoside hydrolase family 5. *Acta Crystallographica, Section D* 56: 3–13.
- Saccardo, P.A. (1880). Fungorum extra-europaeorum Pugillus. *Michelia* 2: 136–149.
- Saccardo, P.A. (1886). *Sylloge Fungorum* 4: 751.
- Sadaba, R.B., Vrijmoed, L.L.P., Jones, E.B.G. and Hodgkiss, I.J. (1995). Observations on vertical distribution of fungi associated with standing senescent *Acanthus ilicifolius* stem at Mai Po Mangrove, Hong Kong. *Hydrobiologia* 295: 119–126.
- Sahashi, N., Kubono, T., Miyasawa, Y. and Ito, S. (1999). Temporal variations in isolation frequency of endophytic fungi of Japanese beech. *Canadian Journal of Botany* 77: 197–202.
- Sahr, T., Ammer, H., Besl, H. and Fischer, M. (1999). Infrageneric classification of the boleticolous genus *Sepedonium*: species delimitation and phylogenetic relationships. *Mycologia* 91: 935–943.

- Sakai, K., Ichikawa, T., Yamada, K., Yamashita, M., Tanimoto, M., Hikita, A., Ijuin, H. and Kondo, K. (1988). Antitumor principles in mosses: the first isolation and identification of maytansinoids, including a novel 15-methoxyansamitocin P-3. *Journal of Natural Products* 51: 845–850.
- Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989). *Molecular Cloning: a Laboratory Manual, 2<sup>nd</sup> Edition*. Cold Spring Harbor Press, New York, USA.
- Samuels, G.J. (1989). *Nectria and Sesquillium*. *Memoirs of New York Botanical Garden* 49: 266–285.
- Sangthong, S. (2002). *Effect of Endomycorrhizal Fungus on Growth of Dendrobium scabrilingue Lindl. and its Hybrids*. M.Sc. Thesis, Chiang Mai University, Thailand.
- Sanmee, R. (2004). *Biodiversity, Host Range, and Growth of Edible Ectomycorrhizal Fungi in Upper Northern Thailand*. Ph.D. Thesis, Chiang Mai University, Thailand.
- Sanmee, R., Yang, Z.L., Lumyong, P. and Lumyong, S. (2003). *Amanita siamensis*, a new species of *Amanita* from Thailand. *Mycotaxon* 87: 225–228.
- Sarbajna, K.K. (1990). New species of *Mycovellosiella* and *Pyricularia* from West Bengal. *Journal of Mycopathological Research* 28: 159–164.
- Sardsud, V., Sittigul C., Sardud, U., Chantrasri, P. and Promin, S. (1998). Endophytic fungi in longan. In L.M. Friut Coates, P.J. Hofman and G.I. Johnson (eds.), *Disease Control and Storage Life Extension*, pp.147–152. Australian Centre for International Agricultural Research, Australia.

- Schulz, B., Römmert, A., Dammann, U., Aust, H. and Strack, D. (1999). The endophyte-host interaction: a balanced antagonism? *Mycological Research* 103: 1275–1283.
- Schulz, B., Wan, K.L.U., Draeger, S. and Sust, H.J. (1993). Endophytes from herbaceous plants and shrubs: effectiveness of surface sterilisation methods. *Mycological Research* 97: 1447–1450.
- Schumann, K. (1904). Zingiberaceae. In A. Engler (ed.), *Das Pflanzenreich*, pp. 1–458. Leipzig, Germany.
- Seifert, K.A. (1985). A monograph of *Stilbella* and some allied genera. *Studies in Mycology* 27: 1–235.
- Seifert, K.A. and Gams, W. (2001). The taxonomy of anamorphic fungi. In D.J. McLaughlin, E.G. McLaughlin and P.A. Lemke (eds.), *Systematics and Evolution, the Mycota VIIa*, pp. 307–347. Springer-Verlag, Berlin, Germany.
- Seifert, K.A., Louis-Seize, G. and Sampson, G. (2003). *Myrothecium acadense*, a new hyphomycete isolated from the weed *Tussilago farfara*. *Mycotaxon* 87: 317–327.
- Senechal, Y., Sanier, C., Gohet, E. and Auzac, J.D. (1987). Different modes de penetration de *Colletotrichum gloeosporioides* dans les feuilles d'*Hevea braziliensis*. *Comptes Rendues de l'Academie Sciences Paris* 305: 537–542.
- Shamoun, S.F. and Sieber, T.N. (2000). Colonisation of leaves and twigs of *Rubus parviflorus* and *R. spectabilis* by endophytic fungi in a reforestation site in British Columbia. *Mycological Research* 104: 841–845.

Shannon, C.E. and Weaver, W. (1949). *The Mathematical Theory of Communication*.

University of Illinois Press, Urbana, USA.

Shull, V. and Hamer, J.E. (1994). Genomic structure and variability in *Pyricularia grisea*. In R. Zeigler, S.A. Leong and P.S. Teng (eds.), *Rice Blast Disease*, pp. 65–86. CAB International, Wallingford, UK.

Sieber, T.N. (1989). Endophytic fungi in twigs of healthy and diseased Norway spruce and white fir. *Mycological Research* 92: 322–326.

Sieber, T.N. and Dorworth, C.E. (1994). An ecological study of endophytic fungi in *Acer macrophyllum* in British Columbia, in search of candidate mycoherbicides. *Canadian Journal of Botany* 72: 1397–1402.

Sieber, T.N. and Hugentobler, C. (1987). Endophytische Pilze in Blättern und Ästen gesunder und geschädigter Buchen (*Fagus sylvatica* L.). *European Journal of Forest Pathology* 17: 411–425.

Sieber, T.N., Riesen, T.K., Müller, E. and Fried, P.M. (1988). Endophytic fungi in four winter wheat cultivars (*Triticum aestivum* L.) differing in resistance against *Stagonospora nodorum* (Berk.) Cast. & Germ. = *Septoria nodorum* Berk. *Journal of Plant Phytopathology (Berlin)* 122: 289–306.

Sieber, T.N., Sieber-Canavesi, F. and Dorworth, C.E. (1991). Endophytic fungi of red alder (*Alnus rubra*) leaves and twigs in British Columbia. *Canadian Journal of Botany* 69: 407–411.

Sieber, T.N., Sieber-Canavesi, F., Petrini, O. and Dorworth, C.E. (1989). The possible use of endophytic fungi in controlling red alder. *Canadian Journal of Plant*

*Pathology* 11: 198.

Sieber-Canavesi, F., Petrini, O. and Seiber, T.N. (1991). Endophytic *Leptostroma* species on *Picea albies*, *Albies alba* and *Abies balsamea*: a cultural, biochemical and numerical study. *Mycologia* 83: 89–96.

Siegel, M.R. and Schardl, C.L. (1991). Fungal endophytes of grasses: detrimental and beneficial associations. In J.H. Andrews and S.S. Hirano (eds.), *Microbial Ecology of Leaves*, pp. 198–221. Springer-Verlag, Berlin, Germany.

Siegel, M.R., Latch, G.C., Bush, L.P., Fannin, N.F., Rowen, D.D., Tapper, B.A., Bacon, C.W. and Johnson, J.M. (1991). Alkaloids and insecticidal activity of grasses infected with different endophytes. *Journal of Chemical Ecology* 16: 3301–3315.

Sinclair, J.B. and Cerkaukas, R.F. (1996). Latent infection vs. endophytic colonisation by fungi. In S.C. Redlin and L.M. Carris (eds.), *Endophytic Fungi in Grasses and Woody Plants: Systematics, Ecology and Evolution*. APS Press, St. Paul, Minnesota, USA, pp. 3–30.

Singh, D.V., Aggarwal, R. and Srivastava, K.D. (2003). The antagonism by *Chaetomium globosum* against spot blotch of wheat caused by *Drechslera sorokiniana*. *Proceedings of the 8<sup>th</sup> International Congress of Plant Pathology*, p. 32. Christchurch, New Zealand.

Siriruga, P. (1998). Thai Zingiberaceae: species diversity and their uses. *Pure and Applied Chemistry* 70: 2111–2118.

Sivichai, S. and Hywel-Jones, N.L. (1999). *Biflagellospora* (aero-aquatic hyphomycetes) from submerged wood in Thailand. *Mycological Research* 103:

908–914.

Sivichai, S., Goh, T.K., Hyde, K.D. and Hywel-Jones, N.L. (1998). The genus *Brachydesmiella* from submerged wood in the tropics, including a new species and a new combination. *Mycoscience* 39: 239–247.

Sivichai, S., Jones, E.B.G. and Hywel-Jones, N.L. (2000). Fungal colonisation of wood in a freshwater stream at Khao Yai National Park, Thailand. *Fungal Diversity*. 5: 71–88.

Sivichai, S., Jones, E.B.G. and Hywel-Jones, N.L. (2002a). Fungal colonisation of wood in a freshwater stream at Tad Ta Phu, Khao Yai National Park, Thailand. *Fungal Diversity* 10: 113–129.

Sivichai, S., Jones, E.B.G. and Hywel-Jones, N.L. (2002b). Lignicolous freshwater higher fungi with reference to their teleomorph and anamorph stages. In R. Watling, J.C. Frankland, A.M. Ainsworth, S. Isaac and C.H. Robinson (eds.), *Tropical Mycology*, pp. 41–49. CAB International, Wallingford, UK.

Smith, J.F., Kress, W.J. and Zimmer, E.A. (1993). Phylogenetic analysis of the Zingiberales based on *rbcL* sequences. *Annals of Missouri Botanical Garden* 80: 620–630.

Smitinand, T. (1995). Overview of the status of biodiversity in tropical and temperate forests. In T.J.B. Boyle and B. Boontawee (eds.), *Measuring and Monitoring Biodiversity in Tropical and Temperate Forests*, pp. 1–4. Center for International Forestry Research, Indonesia.



- Solomon, B.O., Amigun, B., Betiku, E., Ojumu, T.V. and Layokun, S.K. (1999). Optimization of cellulase production by *Aspergillus flavus* Linn isolate NSPR 101 grown on bagasse. *Journal of the Nigerian Society of Chemical Engineers* 16: 61–68.
- Somrithipol, S., Jones, E.B.G. and Hywel-Jones, N.L. (2002). Fungal diversity and succession on pods of *Delonix regia* (Leguminosae) exposed in a tropical forest in Thailand. *Fungal Diversity* 10: 131–139.
- Sontirat, P., Pitakprivan, P., Khamhangritdirong, T., Choobamroong, W. and Kueprakone, U. (1994). *Host Index of Plant Diseases in Thailand*. Mycology Section, Plant Pathology and Microbiology Division, Department of Agriculture, Bangkok, Thailand.
- Soytong, K. (1992). Biological of tomato wilt caused by *Fusarium oxysporum* f. sp. *lycopersici* using *Chaetomium cupreum*. *Kasetsart Journal (Natural Science)* 26: 310–313.
- Soytong, K. and Quimio, T.H. (1989). Antagonism of *Chaetomium globosuum* to the rice blast pathogen, *Pyricularia oryzae*. *Kasetsart Journal (Natural Science)* 23: 198–203.
- Soytong, K., Kanokmedhakul, S., Kukongviriyapa, V. and Isobe, M. (2001). Application of *Chaetomium* species (Ketomium®) as a new broad spectrum biological fungicide for plant disease control: a review article. *Fungal Diversity* 7: 1–15.

- Specht, C.D., Kress, W.J., Stevenson, D.W. and DeSalle, R. (2001). A molecular phylogeny of Costaceae (Zingiberales). *Molecular Phylogeny and Evolution* 21: 333–345.
- Sprague, R. (1950). *Disease of Cereals and Grasses in North America*. Ronald Press, New York, USA.
- Sreenivasaprasad, S. and Mills, P.R. (1998). Future directions for PCR in mycology. In P.D. Bridge, D.K. Arora, C.A. Reddy and R.P. Elander (eds.), *Applications of PCR in Mycology*, pp. 325–345. CAB International, UK.
- Sridhar, K.R. and Bärlocher, F. (1992a). Aquatic hyphomycetes in spruce roots. *Mycologia* 84: 580–584.
- Sridhar, K.R. and Bärlocher, F. (1992b). Endophytic aquatic hyphomycetes of roots of spruce, birch and maple. *Mycological Research* 96: 305–308.
- Stierle, A., Strobel, G. and Stierle, D. (1993). Taxol and taxane production by *Taxomyces andreanae*, an endophytic fungus of Pacific yew. *Science* 260: 214–216.
- Stone, J.K. (1987). Initiation and development of latent infections by *Rhabdocline parkeri* on Douglas fir. *Canadian Journal of Botany* 65: 2614–2621.
- Stone, J.K. (1988). Fine structure of latent infections by *Rhabdocline parkeri* on Douglas fir, with observations on uninfected epidermal cells. *Canadian Journal of Botany* 66: 45–54.
- Stone, J.K., Bacon, C.W. and White, J.F.Jr. (2000). An overview of endophytic microbes: endophytism defined. In C.W. Bacon and J.F.Jr. White (eds.),

*Microbial Endophytes*, pp. 1–29. Marcel Dekker, New York, USA.

Strobel, G. and Daisy, B. (2003). Bioprospecting for Microbial Endophytes and Their Natural Products. *Microbiology and Molecular Biology Reviews* 67: 491–502.

Strobel, G., Daisy, B., Castillo, U. and Harper, J. (2004). Natural products from endophytic microorganisms. *Journal of Natural Products* 67: 257–268.

Strobel, G.A., Hess, W.M., Ford, E., Sidhu, R.S. and Yang, X. (1996a). Taxol from fungal endophytes and the issue of biodiversity. *Journal of Industrial Microbiology and Biotechnology* 17: 412–423.

Strobel, G.A., Miller, R.V., Miller, C., Condron, M., Teplow, D.B. and Hess, W.M. (1999). Cryptocandin, a potent antimycotic from the endophytic fungus *Cryptosporiopsis* cf. *quercina*. *Microbiology* 145: 1919–1926.

Strobel, G.A., Torczynski, R. and Bollon, A. (1997). *Acremonium* sp.: a leucinostatin a producing endophyte of European yew (*Taxus baccata*). *Plant Science* 128: 97–108.

Strobel, G.A., Yang, X., Sears, J., Kramer, R., Sidhu, R.S. and Hess, W.M. (1996b). Taxol from *Pestalotiopsis microspora*, an endophytic fungus of *Taxus wallachiana*. *Microbiology* 142: 435–440.

Summerbell, R.C. (1989). Microfungi associated with the mycorrhizal mantle and adjacent microhabitats within the rhizosphere of black spruce. *Canadian Journal of Botany* 67: 1085–1095.

Suryanarayanan, T.S. and Thennarasan, S. (2002). Temporal variation in endophyte assemblages of *Plumeria rubra*. *Fungal Diversity* 15: 197–204.

Suryanarayanan, T.S. and Vijaykrishna, D. (2001). Fungal endophytes of aerial roots

of *Ficus benghalensis*. *Fungal Diversity* 8: 155–161.

Suryanarayanan, T.S., Senthilarasu, G. and Muruganandam, V. (2000). Endophytic fungi from *Cuscuta reflexa* and its host plants. *Fungal Diversity* 4: 117–123.

Suske, J. and Acker, G. (1987). Internal hyphae in young, symptomless needles of *Picea abies*: electron microscopic and cultural investigations. *Canadian Journal of Botany* 65: 2098–2103.

Sutton, B.C. (1980). *The Coelomycetes*. Commonwealth Mycological Institute, Kew, UK.

Sutton, D.A., Fothergill, A.W. and Rinaldi, M.G. (1998). *Guide to Clinically Significant Fungi, 1<sup>st</sup> Edition*. Williams & Wilkins, Baltimore, Maryland, USA.

Suwanborirux, K., Chang, C.J., Spjut, R.W. and Cassady, J.M. (1990). Ansamitocin P-3, a maytansinoid, from *Claopodium crispifolium* and *Anomodon attenuatus* or associated actinomycetes. *Experientia* 46: 117–120.

Swart, L., Crous, P.W., Petrini, O. and Taylor, J.E. (2000). Fungal endophytes of Proteaceae, with particular emphasis on *Botryosphaeria proteae*. *Mycoscience* 41: 123–127.

Swofford, D.L. (2002). *PAUP\*: Phylogenetic Analysis Using Parsimony (\* and Other Methods), Beta Version 4.0b10*. Sinauer Associates, Sunderland, Massachusetts.

Takamatsu, S. (1998). PCR applications in fungal phylogeny. In P.D. Bridge, D.K. Arora, C.A. Reddy and R.P. Elander (eds.), *Applications of PCR in Mycology*, pp. 125–151. CAB International, UK.

- Tan, B.K.H. and Vanitha, J. (2004). Immunomodulatory and antimicrobial effects of some traditional Chinese medicinal herbs: a review. *Current Medicinal Chemistry* 11: 1423–1430.
- Tang, W., Yang, H. and Ryder, M. (2001). Research and application of *Trichoderma* spp. in biological control of plant pathogens. In S.B. Pointing and K.D. Hyde (eds.), *Bio-Exploitation of Filamentous Fungi, Fungal Diversity Research Series* 6, pp. 403–435. Fungal Diversity Press, Hong Kong.
- Tassi, F. (1899). Novae micromycetum species descriptae et iconibus illustratae. VI. *Bollettino del Laboratorio Orto Botane Reale Universe Siena* 2: 139–163.
- Taylor, J.E., Hyde, K.D. and Jones, E.B.G. (1999). Endophytic fungi associated with the temperate palm, *Trachycarous fortunei*, within and outside its natural geographic range. *New Phytologist* 142: 335–346.
- Templeton, A.R. (1995). Biodiversity at the molecular genetic level: experiences from disparate macroorganisms. In D.L. Hawksworth (ed.), *Biodiversity Measurement and Estimation*, pp. 59–64. The Royal Society Alden Press, Oxford, UK.
- Templeton, G.E. (1992). Use of *Colletotrichum* strains as mycoherbicides. In J.A. Bailey and M.J. Jeger (eds.), *Colletotrichum: Biology, Pathology and Control*, pp. 358–380. CAB International, Wallingford, UK.
- Tenover, F.C. and Rasheed, (1999). Genetic methods for detecting antibacterial and antiviral resistance genes. In P.F. Murray, E.J. Baron M.A. Pfaller, F.C. Tenover and R.H. Tenover (eds.), *Manual of Clinical Microbiology*, 7<sup>th</sup> Edition, pp.

1578–1592. ASM Press, Washington DC, USA.

- Thaithatgoon, P., Thaithatgoon, B., Boonkemthong, C., Tantichareon, M. and Jones, E.B.G. (2004). Edible mushrooms. In E.B.G. Jones, M. Tantichareon and K.D. Hyde (eds.), *Thai Fungal Diversity*, pp. 241–251. BIOTEC, Bangkok, Thailand.
- Thompson, J.D., Higgins, D.G. and Gibson, T.J. (1994). CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position specific gap penalties and weight matrix choice. *Nucleic Acids Research* 22: 4673–4680.
- Thomson, B.D., Robson, A.D. and Abbott, L.K. (1986). Effects of phosphorus and the formation of mycorrhizas by *Gigaspora calospora* and *Glomus fasciculatum* in relation to root carbohydrates. *New Phytologist* 103: 751–765.
- Thongkantha, S., Lumyong, S., Lumyong, P., Whitton, S.R., McKenzie, E.H.C. and Hyde, K.D. (2003). Microfungi on the *Pandanaceae*: *Linocarpon lamiae* sp. nov., *L. siamensis* sp. nov. and *L. suthepensis* sp. nov. are described with a key to *Linocarpon* species from the *Pandanaceae*. *Mycologia* 95: 360–367.
- Tian, X.L., Cao, L.X., Tan, H.M., Zeng, Q.G., Jia, Y.Y., Han, W.Q. and Zhou, S.N. (2004). Study on the communities of endophytic fungi and endophytic actinomycetes from rice and their antipathogenic activities *in vitro*. *World Journal of Microbiology and Biotechnology* 20: 303–309.
- Tode, H.F. (1790). *Fungi Mecklenbergensis Selecti* 1: 25–28. Luneberg.
- Tomita, F. (2003). Endophytes in Southeast Asia and Japan: their taxonomic diversity and potential applications. *Fungal Diversity* 14: 187–204.

- Tomschy, A., Tessier, M., Wyss, M., Brugger, R., Broger, C., Schnoebelen, L., van Loon, A.P.G.M. and Pasamontes, L. (2000). Optimization of the catalytic properties of *Aspergillus fumigatus* phytase based on the three-dimensional structure. *Protein Science* 9: 1304–1311.
- Toofanee, S.B. and Dulyamamode, R. (2002). Fungal endophytes associated with *Cordemoya integrifolia*. *Fungal Diversity* 11: 169–175.
- Toti, L., Viret, O., Horat, G. and Petrini, O. (1993). Detection of the endophyte *Discula umbrinella* in buds and twigs of *Fagus sylvatica*. *European Journal of Forest Pathology*. 23: 147–152.
- Trakoontivakorn, G., Nakahara, K., Shinmoto, H., Takenaka, M., Onishi-Kameyama, M., Ono, H., Yoshida, M., Nagata, T. and Tsushida, T. (2001). Structural analysis of a novel antimutagenic compound, 4-hydroxypanduratin A, and the antimutagenic activity of flavonoids in a Thai spice, fingerroot (*Boesenbergia pandurata* Schult.) against mutagenic heterocyclic amines. *Journal of Agriculture and Food Chemistry* 49: 3046–3050.
- Trinci, A.P.J. (1992). Myco-protein: a twenty-year overnight success story. *Mycological Research* 96: 1–13.
- Tsoumis, G. (1991). *Science and Technology of Wood: Structure, Properties, Utilization*. Van Nostrand Reinhold, New York, USA.
- Tuchinda, P., Reutrakul, V., Claeson, P., Pongprayoon, U., Sematong, T., Santisuk, T. and Taylor, W.C. (2002). Anti-inflammatory cyclohexenyl chalcone derivatives in *Boesenbergia pandurata*. *Phytochemistry* 59: 169–173.

- Tulloch, M. (1972). The genus *Myrothecium* Tode ex Fr. *Mycological Papers* 130: 1–42.
- Turnidge, J.D. and Jorgensen, J.H. (1999). Antimicrobial susceptibility testing: general considerations. In Murray, P.R., Baron, E.J., Tenover, F.C. and Tenover, F.C. and Tenover, F.C. and Tenover, F.C. and Tenover, F.C. and Tenover, F.C. and Tenover, F.C. and Tenover, F.C. and Tenover, F.C. and Tenover, F.C. (eds.), *Manual of Clinical Microbiology 7<sup>th</sup> Edition*, pp. 1469–1473. ASM Press, Washington DC, USA.
- Tyler, V.E. (1996). Natural products and medicine: an overview. In M.J. Balick, E. Elisabetsky and S.A. Laird (eds.), *Medicinal Resources of the Tropical Forest: Biodiversity and its Importance to Human Health*, pp. 3–10. Columbia University Press, New York, USA.
- Tzanov, T., Calafell, M., Guebitz, G.M. and Cavaco-Paulo, A. (2001). Biolpreparation of cotton fabrics. *Enzyme and Microbial Technology* 29: 357–362.
- Umali, T.E., Quimio, T.H. and Hyde, K.D. (1999). Endophytic fungi in leaves of *Bambusa tuldoidea*. *Fungal Science* 14: 11–18.
- Valla, G., Cappellano, A., Huguency, R. and Moiroudo, A. (1989). *Penicillium nodositatum* Valla, a new species including myconodules on *Alnus* roots. *Plant and Soil* 114: 142–146.
- van der Aa, H.A. (1973). Studies in *Phyllosticta* I. *Studies in Mycology* 5: 1–100.
- Vannacci, G. and Harman, G.E. (1987). Biocontrol of seed borne *Alternaria raphani* and *A. brassicola*. *Canadian Journal of Microbiology* 34: 850–856.
- Varghese, K.I.M. and Rao, V.G. (1977). *Solheimia kamatii* sp. nov. (hyphomycete) a



new generic record to India. *Current Science* 46: 23–24.

Verhoeff, K. (1974). Latent infections by fungi. *Annual Reviews of Phytopathology* 12: 99–110.

Viikari, L., Tenkanen, M., Buchert, J., Rättö, M., Bailey, M., Siika-aho, M. and Linko, M. (1993). Hemicellulases for industrial applications. In J.N. Saddler (ed.), *Bioconversion of Forest and Agricultural Plant Residues*, pp. 255–262. CAB International, Wallingford, UK.

Viret, O. and Petrini, O. (1994). Colonisation of beech leaves (*Fagus sylvatica*) by the endophyte *Discula umbrinella* (teleomorph: *Apiognomonina errabunda*). *Mycological Research* 98: 423–432.

Viret, O., Scheidegger, C. and Petrini, O. (1993). Infection of beech leaves (*Fagus sylvatica*) by the endophyte *Discula umbrinella* (teleomorph, *Apiognomonina errabunda*): low temperature scanning electron microscopy studies. *Canadian Journal of Botany* 71: 1520–1527.

Vittal, B.P.R. (1981). *Xenosporium intermedium* sp. nov. from India. *Transactions of the British Mycological Society* 76: 513–515.

von Arx, J.A. (1981). *The Genera of Fungi Sporulating in Pure Culture*. A.R. Gantner, Germany.

Walker, J. (1980). *Gaeumannomyces*, *Linocarpon*, *Ophiobolus* and several other genera of scolecospored ascomycetes and *Phialophora* conidial states, with a note on hyphopodia. *Mycotaxon* 11: 1–129.

Waller, J.M. (1992). *Colletotrichum* diseases of perennial and other cash crops. In J.A. Bailey and M.J. Jeger, *Colletotrichum: Biology, Pathology and Control*, pp.

167–185. CAB International, Wallingford, UK.

Walther, D. and Gindrat, D. (1988). Biological control of damping-off of sugar-beet and cotton with *Chaetomium globosum* and a fluorescent *Pseudomonas* sp. *Canadian Journal of Microbiology* 34: 631–637.

Ward, E. and Bateman, G.L. (1999). Comparison of *Gaeumannomyces*- and *Phialophora*-like fungal pathogens from maize and other plants using DNA methods. *New Phytologist* 141: 323–331.

Watanabe, T., Watanabe, Y. and Nakamura, K. (2003). *Myrothecium dimorphum*, sp. nov., a soil fungus from beach sand in the Bonin (Ogasawara) Islands, Japan. *Mycoscience* 44: 283–286.

Webber, J. (1981). A natural control of Dutch elm disease. *Nature (London)* 292: 449–451.

Wells, H.D., Burton, G.W. and Powell, J.B. (1969). *Piricularia* leaf spot of pearl millet. *Phytopathology* 59: 1057.

Whalley, A.J.S. (1993). Tropical Xylariaceae: their distribution and ecological characteristics. In S. Isacc, J.C. Frankland, R. Watling and A.J.S. Whalley (eds.), *Aspects of Tropical Mycology*, pp. 103–119. Cambridge University Press, Cambridge, UK.

Whalley, A.J.S. (1996). The xylariaceous way of life. *Mycological Research* 100: 897–922.

Whalley, A.J.S. (1997). Xylariaceae. In K.D. Hyde (ed.), *Biodiversity of Tropical Microfungi*, pp. 279–296. Hong Kong University Press, Hong Kong.

White, T.J. (1996). The future of PCR technology: diversification of technologies and

applications. *Trends in Biotechnologies* 14: 478–483.

White, T.J., Bruns, T.D., Lee, S. and Taylor, J.W. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In M.A. Innis, D.H. Gelfand, J.J. Sninsky and T.J. White (eds.), *PCR Protocols: a Guide to Methods and Applications*, pp. 315–322. Academic Press, San Diego, USA.

Widler, B. and Müller, E. (1984). Untersuchungen über endophytische Pilze von *Arctostaphylos uva-ursi* (L.) Sprengel (Ericaceae). *Botanica Helvetica* 94: 307–337.

Wildman, H.G. (1997). Potential of tropical microfungi within the pharmaceutical industry. In K.D. Hyde (ed.), *Biodiversity of Tropical Microfungi*, pp. 29–46. Hong Kong University Press, Hong Kong.

Willetts, K.E., Ekgangaki, A. and Eden, J.A. (2003). Effect of a ginger extract on pregnancy-induced nausea: a randomised controlled trial. *Australia and New Zealand Journal of Obstetric Gynaecology* 43: 139–144.

Wilson, D. (1993). Fungal endophytes: out of sight but should not be out of mind. *Oikos* 68: 379–384.

Wilson, D. (1995). Endophyte: the evolution of a term, and clarification of its use and definition. *Oikos* 73: 247–276.


Wilson, D. and Carroll, G.C. (1994). Infection studies of *Discula quercina*, an endophyte of *Quercus garryana*. *Mycologia* 86: 635–647.

Wilson, E.O. (1992). *The Diversity of Life*. Belknap Press, Cambridge, UK.

- Wiyakrutta, S., Sriubolmas, N., Panphut, W., Thongon, N., Danwisetkanjana, K., Ruangrunsi, N. and Meevootisom, V. (2003). Endophytic fungi with anti-microbial, anti-cancer and anti-malarial activities isolated from Thai medicinal plants. *World Journal of Microbiology and Biotechnology* 20: 265–272.
- Woese, C.R. (1994). There must be a prokaryote somewhere: microbiology's search for itself. *Microbiological Reviews* 58: 1–9.
- Wong, M.K.M. and Hyde, K.D. (2001). Diversity of fungi on six species of Gramineae and one species of Cyperaceae in Hong Kong. *Mycological Research* 105: 1485–1491.
- Wood, T. (1992). VA mycorrhizal fungi: challenges for commercialization. In D.K. Arora, R.P. Elander and K.G. Mukerji (eds.), *Handbook of Applied Mycology, Volume 4, Fungal Biotechnology*, pp. 823–847. Marcel Dekker, New York, USA.
- Wraight, S.P., Carruthers, R.I., Jaronski, S.T., Bradley, C.A., Garza, C.J. and Galaini-Wraight, S. (2000). Evaluation of the entomopathogenic fungi *Beauveria bassiana* and *Paecilomyces fumosoroseus* for microbial control of the silverleaf whitefly, *Bemisia argentifolii*. *Biological Control* 17: 203–217.
- Wu, T.-L. (1981). *Flora Reipublicae Popularis Sinicae, Volume 16*. Science Press, Beijing, China.
- Wuthi-udomlert, M., Grisanapan, W., Luanratana, O. and Caichompoo, W. (2000). Antifungal activity of *Curcuma longa* grown in Thailand. *Southeast Asian Journal of Tropical Medicine and Public Health* 31: 178–182.

- Yaegashi, H. (1977). On the sexuality of blast fungi, *Pyricularia* spp. *Annals of the Phytopathological Society of Japan* 43: 432–439.
- Yamahara, J., Huang, Q., Li, Y., Xu, L. and Fujimura, H. (1990). Gastrointestinal motility-enhancing effect of ginger and its active constituents. *Chemical and Pharmaceutical Bulletins* 38: 430–431.
- Yanna (2001). *Biodiversity, Ecology and Taxonomy of Saprobic Fungi on Palm Fronds*. Ph.D. Thesis, The University of Hong Kong, Hong Kong.
- Ying, J.Z., Mao, X.L., Ma, Q.M., Zong, Y.C. and Wen, H.A. (1987). *Icones of Medicinal Fungi from China*. Science Press, Beijing, China.
- Youssef, N.N. and Dugan, F.M. (2000). Location of an endophytic *Neotyphodium* sp. within various leaf tissues of wild barley (*Hordeum brevisubulatum* subsp. *violaceum*). *Plant Genetic Resources Newsletter* 124: 17–19.
- Yu, T.W., Bai, L., Clade, D., Hoffmann, D., Toelzer, S., Trinh, K.Q., Xu, J., Moss, S.J., Leistner, E. and Floss, H.G. (2002). The biosynthetic gene cluster of the maytansinoid antitumor agent ansamitosisin from *Actinosynnema pretiosum*. *PNAS* 99: 7968–7973.
- Zarr, J.H. (1999). *Biostatistical Analysis, 4<sup>th</sup> Edition*. Prentice Hall, New Jersey, USA.
- Zhou, D.Q. and Hyde, K.D. (2001). Host-specificity, host-exclusivity and host-recurrence in saprobic fungi. *Mycological Research* 105: 1449–1457.
- Zhu, H., Jiang, W., Wang, J., Lei, C., Ling, Z. and Zhu, L. (1992). DNA fingerprint and pathotype characterization of *Pyricularia oryzae* isolates. *Mycosystema* 5: 89–98.

Zucconi, L., Onofri, S. and Persiani, A.M. (1984). Rare or interesting hyphomycetes from tropical forest litter. II. *Pyricularia fusispora* comb. nov., a new species combination for the species *Nakataea fusispora*. *Micologica Italica* 2: 7–10.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่  
Copyright © by Chiang Mai University  
All rights reserved