

TABLE OF CONTENTS

	Page
Acknowledgement	iii
English abstract	iv
Thai abstract	vi
List of tables	viii
List of illustrations	x
Chapter 1 Introduction	1
1. 1. Principle and rationale	1
1. 2. Objectives	2
1. 3. Usefulness of research	2
1. 4. Experiment scope	2
Chapter 2 Theory and Literature reviews	3
2. 1. Pineapple	3
2. 1. 1. Botany	3
2. 1. 2. Inflorescence and flower	4
2. 1. 3. Fruit development	5
2. 1. 4. Physico-chemical changes during fruit maturation and ripening.	6
2. 2. Quality of pineapple fruit and environmental factors affected quality	16
2. 2. 1 Quality of pineapple fruit	16
2. 2. 2 Environmental factors affected fruit quality	17
2. 2. 3 Translucency in pineapple	18
Chapter 3 Change of morphological attributes in relation to maturity and cropping seasons of "Smooth Cayenne" pineapple fruit	20
3. 1. Introduction	20
3. 2. Materials and methods	21
3. 3. Results and discussion	25
3. 4. Conclusion	39

TABLE OF CONTENTS (CONTINUE)

	Page
Chapter 4 Influence of maturity and cropping seasons on physico-chemical and sensory attributes of “Smooth Cayenne” pineapple fruit	41
4. 1. Introduction	41
4. 2. Experiment 1: Assessment of color quality, color pigments and flesh translucency of pineapple fruits	42
4. 2. 1. Materials and methods	42
4. 2. 2. Results and discussion	44
4. 2. 3. Conclusion	59
4. 3. Experiment 2: Assessment of physico-chemical attributes of pineapple fruits	63
4. 3. 1. Materials and methods	63
4. 3. 2. Results and discussion	65
4. 3. 3. Conclusion	78
4. 4. Experiment 3: Change of sensory quality and ripening of pineapple fruit	80
4. 4. 1. Materials and methods	80
4. 4. 2. Results and discussion	80
4. 4. 3. Conclusion	87
Chapter 5 Overall conclusions	88
References	90
Appendices	98
Appendix A Data of morphological attributes of pineapple fruits of all crops for the years 2002 and 2003	99
Appendix B Data of physico-chemical attributes and sensory attributes of pineapple fruits of all crops for the years 2002, 2003 and 2004	102
Curriculum vitae	115

LIST OF TABLES

Tables		Page
2.1	Proximate analysis of “Smooth Cayenne” fruit	9
2.2	Structure component of cell wall	12
3.1	Fruit shape as determined by ratio of base and top diameters and fruit length.	24
3.2	Environmental factors during four months (0-110 days) and the correlation to fruit weight of different crops of years 2002 and 2003	28
3.3	Morphological attributes of pineapple fruits harvested at 130 DAFB from all crops for the years 2002 and 2003	29
4.1	Physico-chemical properties of pineapple fruits harvested at 130 DAFB from all crops for the years 2002 and 2003	47
4.2	Flesh firmness of all parts and positions of pineapple fruit from all crops for the year 2002	67
4.3	Flesh firmness of all parts and positions of pineapple fruit from all crops for the year 2003	68
4.4	Chemical attributes of pineapple fruits harvested at 130 DAFB from all crops for the years 2002 and 2003	73
4.5	Chemical properties and sugars content of pineapple fruits at 130 DAFB in the regular crop for the year 2004	76
4.6	Moisture and crude fiber content (%) of pineapple fruits from all crop for the year 2002	79
4.7	Sensory quality of pineapple fruit at each harvesting date from all crops for the years 2002 and 2003	82
4.7	Sensory quality of pineapple fruit at each harvesting date from all crops for the years 2002 and 2003 (continue)	83

LIST OF ILLUSTRATIONS

Figure	Page
2.1 Pineapple flower and fruit	4
2.2 Changes in sugar contents and activities of sugar metabolic enzymes in developing pineapple flesh. (A) sucrose (Suc), glucose (Glu), and fructose (Fru) contents. (B) Activities of sucrose synthase (SS) and sucrose phosphate synthase (SPS). (C) Activities of acid invertase (AI), neutral invertase (NI), cell-wall invertase (CWI). Fruit were harvested every 2 weeks from 12 weeks before harvest. Symbols represent mean \pm SD of three replications.	8
2.3 Chemical structure (a) cellulose, (b) hemicellulose, (c) pectic and (d) idealized cell wall model.	14
3.1 Time schedule of harvesting and total rainfall (mm), relative humidity (%), maximum and minimum temperatures ($^{\circ}$ C) during flowering induction, fruit development and harvesting of pineapple fruit produced in Lampang Province, Northern, Thailand. The early, regular and late crops were harvested, during the summer, rainy season and winter, respectively.	23
3.2 Fruit shape of pineapple	24
3.3 Fruit weight, fruit length, fruit diameter and fruitlet number For the early, regular and late crops for the years 2002 and 2003	30
3.4 Shape distribution of pineapple fruit at different cropping seasons (A) crop year 2002 and (B) crop year 2003	31
3.5 Relations between fruit weight and shape distribution of each shape. (a), (b) early crop, (c), (d) regular crop, and (e), (f) late crop of years 2002 and 2003.	32
3.6 Characteristics of pineapple fruit of different cropping seasons harvested at 130 DAFB.	33
3.7 Crown weight, crown length and crown diameter of pineapple fruits harvested in the early, regular and late crops of years 2002 and 2003	35
3.8 Relationship of fruit and crown weights in all crops for the year 2002	36
3.9 Relationship of fruit and crown weights in all crops for the year 2003	37

LIST OF ILLUSTRATIONS (CONTINUE)

Figure	Page
3.10 Regression of fruit and crown weight in relation to solar radiation and temperatures (T day, T night).	40
4.1 Shell color score of pineapple	42
4.2 Average of shell color score changes during harvesting period in the early, regular and late crops of years 2002 and 2003.	48
4.3 Percentage of fruit with different shell color score of all crops at harvesting date 110-160 DAFB (A-F) for the year 2002	49
4.4 Percentage of fruit with different shell color score of all crops at harvesting date 110-160 DAFB (A-F) for the year 2003	50
4.5 Comparative changes of total chlorophyll (A) and carotenoids (B) of pineapple peel of crop year 2003	51
4.6 Carotenoids of pineapple flesh from all crops at harvesting time for the year 2003 at absorption 420 nm (A) and 447 nm (B)	53
4.7 L* value of pineapple flesh from all crops at harvesting time for the years 2002 (A) and 2003 (B)	54
4.8 a* value of pineapple flesh from all crops at harvesting time for the years 2002 (A) and 2003 (B)	55
4.9 b* value of pineapple flesh from all crops at harvesting time of crop years 2002 (A) and 2003 (B)	56
4.10 Hue angle of pineapple flesh from all crops at harvesting time for the years 2002 (A) and 2003 (B)	57
4.11 Chroma value of pineapple flesh from all crops at harvesting time for the years 2002 (A) and 2003 (B)	58
4.12 Percentage translucency and normal pineapple fruit at each harvesting time of all crops for the year 2002	60
4.13 Percentage translucency and normal fruit at each harvesting time of all crops for the year 2003	61
4.14 Percentage of translucency and normal fruit of all crops for the years 2002 and 2003	62

LIST OF ILLUSTRATIONS (CONTINUE)

Figure	Page
4.15 Flesh firmness of 3 parts (basal, medial, top) and 3 positions (inner, middle, outer) of all crops for the year 2002	69
4.16 Flesh firmness of 3 parts (basal, medial, top) and 3 positions (inner, middle, outer) of all crops for the year 2003	70
4.17 Total soluble solids and titratable acidity of flesh pineapple of all crops for the years 2002 and 2003	74
4.18 TSS/TA ratio and pH of flesh pineapple fruit of all crops for the years 2002 and 2003	75
4.19 Changes in sugar contents of the regular crop for the year 2004	77
4.20 Spider plots of five sensory attributes (flavor) of pineapple fruit harvested at 130 DAFB of all crops for the years 2002 and 2003	84
4.21 Spider plots of five sensory attributes (texture) of pineapple fruit harvested at 130 DAFB of all crops for the years 2002 and 2003	85