

## REFERENCES

1. Ahmad, I. and Beg, A. Z. (2001). Antimicrobial and phytochemical studies on 45 Indian medicinal plants against multi-drug resistant human pathogens. *J. Ethnopharmacol.*, 74, 113-123.
2. Akiyama, H., Fujii, K., Yamasaki, O., Oono, T. and Iwatsuki, K. (2001). Antibacterial action of several tannins against *Staphylococcus aureus*. *J. Antimicrobiol. Chemother.*, 48, 487-491.
3. Amarowicz, R. and Shahidi, F. (1996). A rapid chromatographic method for separation of individual catechins from green tea. *Food Res. Int.*, 29, 71-76.
4. Azis, N. H., Farag, S. E., Mousa, L. A. and Abo-Zaid, M. A. (1998). Comparative antibacterial and antifungal effects of some phenolic compounds. *Microbios.*, 93, 43-54.
5. Barry, A. L. and Thornsberry, C. (1991). Susceptibility tests: diffusion test procedures. in A. Balows, W. J. Hausler, K. L. Herrmann, H. D. Isenberg and S. H. Jean, *Manual of Clinical Microbiology* (pp. 1117-1125), Washington, DC.: American Society for Microbiology.
6. Basílico, M. Z. and Basílico, J. C. (1999). Inhibitory effects of some spice essential oils on *Aspergillus ochraceus* NRRL 3174 growth and ochratoxin A production. *Lett. Appl. Microbiol.*, 29, 238-241.

7. Bauer A. W., Kirby, W. M. M., Sherris, J. C. and Truck, M. (1966). Antibiotic susceptibility testing by a standardized single disc method. *Am. J. Clin. Pathol.*, 45, 493-496.
8. Betina, V. (1973). Bioautography in paper and thin-layer chromatography and its scope in the antibiotic field. *J. Chromatogr.*, 78, 41-51.
9. Binutu, O. A., Adesogan, K. E. and Okogun, J. I. (1996). Antibacterial and antifungal compounds from *Kigelia pinnata*. *Planta Med.*, 62, 352-353.
10. Borris, R. P. (1996). Natural products research: perspectives from a major pharmaceutical company. *J. Ethnopharmacol.*, 51, 29-38.
11. Boskabady, M. H. and Ramazani-Assari, M. (2001). Relaxant effect of *Pimpinella anisum* on isolated guinea pig tracheal chains and its possible mechanism(s). *J. Ethnopharmacol.*, 74, 83-88.
12. Brukhardt, G., Reichling, I., Martin, R. and Becker, H. (1986). Terpene hydrocarbons in *Pimpinella anisum*. *Pharm. Weekly Sci.*, 8, 190-193.
13. Chanwitheesuk, A., Rakariyatham, N., Teerawutgulrag, A., Wongchanapiboon, T. (2002). Screening of antioxidants from some Thai vegetables and herbs. *Chiang Mai J. Sci.*, 29, 1-5.
14. Cowan, M. M. (1999). Plant products as antimicrobial agents. *Clin. Microbiol. Rev.*, 12, 564-582.
15. Croft, K. D. (1999). Antioxidant effects of plant phenolic compounds. in T. K. Basu, N. J. Temple and M. L. Grag, *Antioxidants in Human Health and Disease* (pp. 109-120), Oxford: CABI Publishing.

16. Davidson, P. M. (1993). Parabens and phenolic compounds. in P. M. Davidson and A. L. Branen, *Antimicrobials in Foods* (pp. 263-306), New York: Marcel Dekker.
17. Duffy, C. F. and Power, R. F. (2001). Antioxidant and antimicrobial properties of some Chinese plant extracts. *Int. J. Antimicrobiol. Agents*, 17, 527-529.
18. Fernández, M. A., García, M. D. and Sáenz, M. T. (1996). Antibacterial activity of the phenolic acids fractions of *Scrophularia frutescens* and *Scrophularia sambucifolia*. *J. Ethnopharmacol.*, 53, 11-14.
19. Fernández, M. A., Sáenz, M. T. and García, M. D. (1998). Anti-inflammatory activity in rats and mice of phenolic acids isolated from *Scrophularia frutescens*. *J. Pharm. Pharmacol.*, 50, 1183-1186.
20. Flück, H. (1988). *Medicinal Plants and Their Uses* (p. 107), London: W. Foulsham.
21. Gennaro, A. R., Nora, A. H., Nora, J. J., Stander, R. W. and Weiss, L. (1979). *Blakiston's Gould Medical Dictionary: A Modern Comprehensive Dictionary of the Terms Used in All Branches of Medicine and Allied Sciences; with Illustrations and Table* (p. 94), New York: McGraw-Hill.
22. Goodwin, T. W. and Mercer, E. I. (1983). *Introduction to Plant Biochemistry* (pp. 528-566), New York: Pergamon Press.
23. Gordon, M. H. (1990). The mechanism of antioxidant action *in vitro*. in B. J. F. Harborne, *Food Antioxidants* (pp. 1-18), London: Elsevier Science.

24. Grieve, M. (1974). *A Modern Herbal: The Medicinal, Culinary, Cosmetic and Economic Properties, Cultivation, and Folklore of Herbs, Grasses, Fungi, Shrubs, and Trees with All Their Modern Scientific Uses* (pp. 41-43), New York: Hafner Press.
25. Grover, J. K., Yadav, S. and Vats, V. (2002). Medicinal plants of India with anti-diabetic potential. *J. Ethnopharmacol.*, 81, 81-100.
26. Gülçin, İ., Oktay, M., Kireççi, E., İrfan, Ö. and Küfrevioğlu, İ. (2003). Screening of antioxidant and antimicrobial activities of anise (*Pimpinella anisum* L.) seed extractd. *Food Chem.*, 83, 371-382.
27. Hammerschmidt, P. A. and Pratt, D. E. (1978). Phenolic antioxidants of dried soybeans. *J. Food Sci.*, 43, 556-559.
28. Harborne, J. B. (1979). Variation in and functional significance of phenolic conjugation in plants. in T. Swan, J. B. Harborne and C. F. van Sumere, *Biochemistry of Plant Phenolics* (pp. 457-474), New York: Plenum Press.
29. Harborne, J. B. (1998). *Phytochemical Methods: A Guide to Modern Techniques of Plants Analysis* (pp. 40-106), London: Chapman & Hall.
30. Haslan, E. (1998). *Practical Polyphenolics: From Structure to Molecular Recognition and Physiological Action* (pp. 138-177), Cambridge: Cambridge University Press.
31. Hayashi, K., Kamiya, M. and Hayahi, T. (1995). Virucidal effect of the steam distillate from *Houttuynia cordata* and its components on HSV-1, influenza virus, and HIV. *Planta Med.*, 61, 237-241.
32. Ibrahim, D. and Osman, H. (1995). Antimicrobial activity of *Cassia alata* from Malaysia. *J. Ethnopharmacol.*, 45, 151-156.

33. Jacquat, C. (1990). *Plants from the Market of Thailand* (pp. 43-44), Bangkok: Duang Kamol.
34. Kar, A., Choudhary, B. K. and Bandyopadhyay, N. G. (2003). Comparative evaluation of hypoglycemic activity of some Indian medicinal plants in alloxan diabetic rats. *J. Ethnopharmacol.*, 84, 105-108.
35. Kochhar, S. P. and Rossell, J. B. (1990). Detection, estimation and evaluation of antioxidants in food systems. in B. J. F. Harborne, *Food Antioxidants* (pp. 19-65), London: Elsevier Science.
36. Kreydiyyeh, S. I., Usta, J., Knio, K., Markossian, S. and Dagher, S. (2003). Aniseed oil increases glucose absorption and reduces urine output in the rat. *Life Sci.*, 74, 663-673.
37. Kroes, B. H., van den Berg, A. J. J., Quarles van Ufford, H. C., van Dijk, H. and Labadie, R. P. (1992). Anti-inflammatory activity of gallic acid. *Planta Med.*, 58, 499-504.
38. Kubo, I., Fugita, K. and Nihei, K. (2003). Molecular design of multifunctional antibacterial agents against methicillin resistant *Staphylococcus aureus* (MRSA). *Bioorg. & Med. Chem.*, 11, 4255-4262.
39. Kubo, I., Xiao, P. and Fugita, K. (2002). Anti-MRSA activity of alkyl gallates. *Bioorg. & Med. Chem.*, 12, 113-116.
40. Larson, R. A. (1988). The antioxidants of higher plants. *Phytochem.*, 27, 969-978.
41. Liégeois, C., Lermusieau, G. and Collin, S. (2000). Measuring antioxidant efficiency of wort, malt, and hops against the 2,2'-azobis (2-amidinopropane) dihydrochloride-induced oxidation of an aqueous dispersion of linoleic acid. *J. Agric. Food Chem.*, 48, 1129-1134.

42. Lin Hsu, F., Chueh, Chen, Y. and Tang Cheng, J. (2000). Caffeic acid as active principle from the fruit of *Xanthium strumarium* to lower plasma glucose in diabetic rats. *Planta Med.*, 66, 228-230.
43. Ma, W. G., Fukushi, Y., Tahara, S. and Osawa, T. (2000). Fungitoxic alkaloids from Hokkaido Papaveraceae. *Fitoter.*, 71, 527-534.
44. Masuda, T., Inazumi, A., Yamada, Y., Padolima, W. G., Kikuzaki, H. and Nakatani, N. (1991). Antimicrobial phenylpropanoids from *Piper sarmentosum*. *Phytochem.*, 30, 3227-3228.
45. McMakin, P. D. (1988). *A Field Guide to the Flowering Plants of Thailand* (p76), Bangkok: White Lotus.
46. Mihopoulos, N., Vagias, C., Chinou, I., Roussakis, C., Scoullou, M. Harvala, C. and Roussis, V. (1999). Antibacterial and cytotoxic natural and synthesized hydroquinone from sponge *Ircinia spinosula*. *J. Biosci.*, 54, 417-423.
47. Miki, K., Yasushi, O., Yunmo, R., Tetsuro, I., Hideyuki, Y., Toshihiro, A., Motohiro, W., Norio, Y., Seigou, A., Genzou, T., Shinya, M., Kohshi, G., Hisayoshi, F. and Kazunori, F. (2001). Antitumor effect of gallic acid on LL-2 lung cancer cells transplanted in mice. *Anti-Cancer Drugs*, 12, 847-852.
48. Moreira, A. S., Spitzer, V., Schapoval, E. E. and Schenkel, E. P. (2000). Anti-inflammatory activity of extracts and fractions from the leaves of *Gochnatia polymorpha*. *Phytother. Res.: PTR.*, 14, 638-640.
49. Moreman, D. E. (1996). An analysis of the food plants and drug plants of native North America. *J. Ethnopharmacol.*, 52, 1-22


50. Naczki, M., Wanasundara, P. K. J. P. D. and Shahadi, F. (1992). Facile spectrophotometric quantification method of sinapic acid in hexane-extracted and methanol-ammonia-water-treated mustard and rapeseed meals. *J. Agric. Food Chem.*, 40, 444-448.
51. Naidu, A. S. (2000). Phytoantimicrobial (PAM) agents as multifunctional food additives. in W. R. Bidlack, S. T. Omaye, M. S. Meskin and D. K. W. Topham, *Phytochemicals As Bioactive Agents* (pp. 105-129.), Lancaster: Technomic Publishing.
52. Nakamura, C. V., Ueda-Nakamura, T., Bando, E., Melo, A. F. N., Cortez, D. A. G. and Filho, B. P. D. (1999). Antibacterial activity of *Ocimum gratissimum* L. essential oil. *Mem. Inst. Oswaldo Cruz, Rio de Janeiro*, 94, 675-678.
53. Nasir, E. and Ali, S. I. (1981). *Flora of Pakistan* (pp. 19-20), Karachi: Department of Botany, University of Karachi.
54. Nychas, G. J. E. (1995). Natural antimicrobials from plants. in G. W. Gould, *New Methods of Food Preservation* (pp. 58-89), London: Academic Professional.
55. Panizzi, L., Caponi, C., Catalano, S., Cioni, P. L. and Morelli, I. (2002). *In vitro* antimicrobial activity of extracts and isolated constituents of *Rubus ulmifolius*. *J. Ethnopharmacol.*, 79, 165-168.
56. Penna, C., Marino, S., Vivot, E., Cruaños, M. C., Muñoz, J. de D., Cruaños, J., Ferraro, G., Gutkind, G. and Martino, V. (2001). Antimicrobial activity of Argentine plants used in the treatment of infectious diseases: isolation of active compounds from *Sebastiania brasiliensis*. *J. Ethnopharmacol.*, 77, 37-40.

57. Pietta, P. G. (2000). Flavonoids as antioxidants. *J. Nat. Prod.*, 63, 1035-1042.
58. Ponce, A. G., Fritz, R., del Valle, C. and Roura, S. I. (2003). Antimicrobial activity of essential oils on the native microflora of organic Swiss chard. *Lebensm.-Wiss. u.-Technol.*, 36, 679-684.
59. Pourgholami, M. H., Majzoob, S., Javadi, M., Kamalinejad, M., Fanace, G. H. R. and Sayyah, M. (1999). The fruit essential oil of *Pimpinella anisum* exerts anticonvulsant effects in mice. *J. Ethnopharmacol.*, 66, 211-215.
60. Prodr, R. B. (1984). Gymnema. *Flora of China*, 16, 239-241.
61. Puupponen-Pimiä, R., Nohynek, L., Meier, C., Kähkönen, M., Hopia, A. and Oksman-Caldentey, K. M. (2001). Antimicrobial properties of phenolic compounds from berries. *J. Appli. Microbiol.*, 90, 494-507.
62. Rama Raje Urs, N. V. and Dunleavy, J. M. (1975). Enhancement of the bactericidal activity of a peroxidase system by phenolic compounds. *Phytopathol.*, 65, 686-690.
63. Raymond, J., Rakariyatham, N. and Azanza, J. L. (1984). Polyphenol-protein interactions in sunflower seed (*Helianthus annuus*). *Sci. Des Ali.*, 4, 95-108.
64. Rice-Evan, C. A. and Miller, N. J. (1996). Antioxidant activities of flavonoids as bioactive components of food. *Biochem. Soc. Trans.*, 24, 790-795.
65. Rodrigues, V. M., Rosa, P. T., Marques, M. O., Petenate, A. J. and Meireles, M. A. (2003). Supercritical extraction of essential oil from aniseed (*Pimpinella anisum* L.) using CO<sub>2</sub>: solubility, kinetics, and composition data. *J. Agric. Food Chem.*, 51, 1518-1523.
66. Sakorn, P. (2002). *Production, Purification and Characterization of Myrosinase from Aspergillus sp. NR-4201*. Ph.D. Thesis, Chiang Mai University.

67. Santos, P. M., Figueiredo, A. C., Oliveira, M. M., Barroso, J. G., Pedro, L. G., Deans, S. G., Younus, A. K. M. and Scheffer, J. J. C. (1998). Essential oils from hairy root cultures and from fruits and roots of *Pimpinella anisum*. *Phytochem.*, 48, 455-460.
68. Saxena, G., Mc Cutcheon, A., Farmer, S., Towers, G. and Hancock, R. (1994). Antimicrobial constituents of *Rhus glabrata*. *J. Ethnopharmacol.*, 42, 95-99.
69. Scalbert, A. (1991). Antimicrobial properties of tannins. *Phytochem.*, 30, 3875-3883.
70. Senji, S., Lekh Raj, J. and Makoto, T. (2000). Antimicrobial effects of green tea polyphenols on thermophilic spore-forming bacteria. *J. Biosci. Bioeng.*, 90, 81-85.
71. Shetty, K. (2004). Role of proline-linked pentose-phosphate pathway (PLPPP) in phenolic phytochemical biosynthesis and mechanism of action for human health and environmental applications. *Asia-Pacific J. Clin. Nut.*, 13, 1-24.
72. Shimizu, K., Ozeki, M., Iino, A., Nakajyo, S., Urakawa, N. and Atsuchi, M. (2001). Structure-activity relationships of triterpenoid derivatives extracted from *Gymnema iodorum* leaves on glucose absorption. *Jpn. J. Pharmacol.*, 86, 223-229.
73. Shimizu, K., Ozeki, M., Tanaka, K., Itoh, K., Nakajyo, S., Urakawa, N. and Atsuchi, M. (1997). Suppression of glucose absorption by extracts from the leaves of *Gymnema inodorum*. *J. Vet. Med. Sci.*, 59, 753-757.
74. Shukla, Y. N., Srivastava, A., Kumar, S. and Kumar, S. (1999). Phytotoxic and antimicrobial constituents of *Argyreia speciosa* and *Oenothera biennis*. *J. Ethnopharmacol.*, 67, 241-245.

75. Smitinand, T. (1984). *Flora of Thailand; Volume Four Part One* (pp. 67-69), Bangkok: The Forest Herbarium, Royal Forest Department.
76. Stapleton, P. D., Shah, S., Anderson, J. C., Hara, Y., Hamilton-Miller, J. M. T. and Taylor, P. W. (2004). Modulation of  $\beta$ -lactam resistance in *Staphylococcus aureus* by catechins and gallates. *Int. J. Antimicrob. Agents*, 23, 462-467.
77. Stobart, T. (1977). *Herbs, Spices and Flavourings* (pp. 46-47), Harmondsworth: Penguin.
78. Strack, D. (1997). Phenolic metabolism. in P. M. Dey and J. B. Harborne, *Plant Biochemistry* (pp. 387-416), London: Academic Press.
79. Tringali, C., Spatafora, C. and Longo, O. D. (2000). Bioactive constituents of the bark of *Parkia biglobosa*. *Fitoter.*, 71, 118-125.
80. Uhl, S. R. (2000). *Handbook of Spices, Seasonings, and Flavorings* (pp. 62-64), Lancaster, Pa.: Technomic Publishing.
81. Van der Watt, E. and Pretorius, J. C. (2001). Purification and identification of active antibacterial components in *Carpobrotus edulis* L. *J. Ethnopharmacol.*, 76, 87-91.
82. Venkateswaran, S. and Pari, L. (2003). Effect of *Coccinia indica* leaves on antioxidant status in streptozotocin-induced diabetic rats. *J. Ethnopharmacol.*, 84, 163-168.
83. Vijaya, M., Cass, I., Judy, G., Nadeem, T., Bobby, W. E., Debasis, B. and Harry, G. P. (2001). Antifungal activities of origanum oil against *Candida albicans*. *Molec. Cell Biochem.*, 228, 111-117.

84. Wanasundara, U., Amarowicz, R. and Shahidi, F. (1994). Isolation and identification of an antioxidative component in canola meal. *J. Agric. Food Chem.*, 42, 1285-1290.
85. Waterman, P. G. and Mole, S. (1994). *Method in Ecology: Analysis of Phenolic Plant Metabolites* (pp. 2-3), London: Blackwell Scientific.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่  
Copyright © by Chiang Mai University  
All rights reserved