

REFERENCES

- Al-Aidross, K. and Seifert, A.M. (1980). Polysaccharide and protein degradation, germination, and virulence against mosquitoes in the entomopathogenic fungus *Metarhizium anisopliae*. *J. Inv. Pathol.*, 36: 29-34.
- Adelberg, E.A., Mandel, M. and Chen, G.C.C. (1965). Optimal conditions for mutagenesis by N-methyl-N'-nitro-N-nitroso guanidine in *Escherichia coli* K12. *Biochem. Biophys. Res. Commun.*, 18:788-795.
- Barcelo, S., Gardiner, J.M., Gascher, A. and Chipman, J.K. (1996). CYPE2E1-mediated mechanisms of anti-genotoxicity of the broccoli constituent sulforaphane. *Carcinogenesis.*, 17: 277-282.
- Benn, M. (1977). Glucosinolates. *Pure Appl. Chem.*, 49: 197-210.
- Bennett, R.N. and Wallsgrove, R.M. (1994) Secondary metabolites in plant defense-mechanisms. *New Phytol.*, 127: 617-633.
- Bestwick, L.A., Grønning, L.M., James, D.C., Bones, A.M. and Rossiter, J.T. (1993). Purification and characterization of a nitrilase from *Brassica napus*. *Physiol Plant.*, 89(4): 811-816.
- Bjerg, B. and Sørensen, H. (1987). Isolation of intact glucosinolates by column chromatography and determination of their purity. In: World Crops: Production, Utilization, Description. Glucosinolates in Rapeseeds: Analytical Aspects., (Ed. J.-P. Wathélet). Martinus Nijhoff Publishers, Kluwer Academic Publishers Group., 13: 59-75.
- Bjorkman, R. and Janson, J.C. (1972). Studies on myrosinases I: Purification and characterization of a myrosinase from white mustard seed (*Sinapis alba* L.). *Biochem. Biophys. Acta.*, 276: 508-518.
- Bjorkman, R. and Lonnerdal, B. (1973). Studies on myrosinases. III. Enzymatic

properties of myrosinases from *Sinapis alba* and *Brassica napus* seeds.

Biochim. Biophys. Acta., 327: 121-131.

Bjorkman, R. (1976). Properties and function of plant myrosinases. In The biology and chemistry of *Cruciferae* (Vaughan, J.G., A.J. MacLeod and B.M.G. Jones, eds.). Academic Press, London., 191-205.

Block, G., Patterson, B. and Subar, A. (1992). Fruit, vegetables and cancer prevention: A review of the epidemiological evidence. *Nutr. Cancer.*, 18: 1-29.

Bones, A.M. (1991). Compartmentation and molecular properties of thioglucoside glucohydrolase (myrosinase). Dr-thesis. Unigen Center for Molecular Biology, Department of Botany, University of Trondheim., 1-195.

Bones, A.M. and Rossiter, J.T. (1996). The myrosinase-glucosinolate system, its organization and biochemistry. *Physiologia Plantarum.*, 97: 194-208.

Bones, A.M., Visvalingam, S. and Thangstad, O.P. (1994). Sulphate can induce differential expression of thioglucoside glucohydrolases (myrosinase). *Planta.*, 193: 558-566.

Bones, A.M. and Slupphaug, G. (1989). Purification, characterization and partial amino acid sequencing of β -thioglucosidase from *Brassica napus* L. *J. Plant Physiol.*, 134: 722-729.

Brabban, A.D. and Edwards, C. (1995). The effects of glucosinolates and their hydrolysis products on microbial growth. *J. Appl. Bacteriol.*, 79: 171-177.

Buchwaldt, L., Larsen, L.M., Plöger, A. and Sørensen, H. (1986). Fast polymer liquid chromatography isolation and characterization of plant myrosinase, β -thioglucoside glucohydrolase, isoenzymes. *J. Chromatography.*, 363: 71-80.

Burmeister, W.P., Cottaz, S., Driquez, H., Iori, R., Palmieri S. and Henrissat, B.

- (1997). The crystal structures of *Sinapis alba* myrosinase and a covalent glycosyl-enzyme intermediate provide insights into the substrate recognition and active-site machinery of an S-glycosidase. *Structure.*, 5: 663-675.
- Bussy, A. (1840). Sur la formation de l'huile essentielle de moutarde. *J. Pharmacol.*, 27: 464-471.
- Carlile, M.J. and Watkinson, S.C. (1994). *The Fungi*. Academic Press Ltd, London., 329 - 340.
- Chadchawan, S., Bishop, J., Thangstad, O.P., Bones, A.M., Mitchell-Olds, T. and Bradley, D. (1993). *Arabidopsis* cDNA sequence encoding myrosinase. *Plant Physiol.*, 103: 671-672.
- Chadha, K.C., Dembinski, W.E., Dunn, C.B., Ardi, J., Bardos, T.J., Dunn, J.A. and Ambrus, Sr. J.L. (2004). Effect of increasing thiolation of the polycytidylic acid strand of poly I: poly C on the alpha, beta and gamma interferon inducing properties, antiviral and antiproliferative activities. *Antiviral Res.*, 64: 171-177.
- Chisholm, H. (1973). Biosynthesis of 3-Methoxycarbonylpropyl-Glucosinolate in an *Erysimum* species. *Phytochemistry.*, 12: 605-608.
- Christensen, B.W., KJÆR, A., Øgaard, M.J., Olsen, C.E., and Sørensen, H. (1982). Mass-Spectrometric Characteristics of some Pertrimethylsilylated Desulfoglucosinolates. *Tetra.*, 38: 353-357.
- Doll, R. (1992). The lessons of life: Keynote address to the nutrition and cancer conference. *Cancer Res.*, 52: 2024s-2029s.
- Duncan, A.J. and Milne, J.A. (1989). Glucosinolates. *Aspects of Applied Biology.*, 19: 75-92.
- Ettlinger, M.G. and Lundeen, A.J. (1956). The structures of sinigrin and sinalbin; an

enzymatic rearrangement. *Journal of the American Chemical Society.*, 78: 4172-4173.

Fahey, J.W., Zhag, Y. and Talalay, P. (1997). Broccoli sprouts; an exceptionally rich source of inducers of enzymes that protect against chemical carcinogens. *Proceedings of the National Academy of Science of the USA.*, 94: 10367-10372.

Fahey, J.W., Zalcmann, A.T. and Talalay, P. (2001). The chemical diversity and distribution of glucosinolates and isothiocyanates among plants. *Phytochem.*, 56: 5-51.

Falk, A., Xue, J., Lenman, M. and Rask, L. (1992). Sequence of a cDNA clone encoding the enzyme myrosinase and expression of myrosinase in different tissues of *Brassica napus*. *Plant Sc.*, 83: 181-186.

Falk, A., Taipalensu, J.E.B., Lenman, M. and Rask, L. (1995). Characterization of rapeseed myrosinase-binding protein. *Planta.*, 195: 387-395.

Gadamer, J. (1897). Uber das sinigrin. *Bericthe der Deutche Chemie.*, 30: 2322.

Gamet-Payraastre, L., Li, P. and Lumeau, S. (2000). Sulforaphane, a naturally occurring isothiocyanate, induces cell cycle arrest and apoptosis in Ht29 human colon cancer cells. *Cancer Res.*, 60: 1426-1433.

Getahun, S.M. and Chung, F.L. (1999). Conversion of glucosinolates to isothiocyanates in humans after ingestion of cooked watercress. *Cancer Epidemiol Biomarkers Prev.*, 8: 447-51.

Glendening, T.M., Poulton, J.E., (1988). Glucosinolate biosynthesis. Sulfation of desulfoglucosinolate by cell-free extracts of cress (*Lepidium sativum* L.) seedlings. *Plant Physiology.*, 86: 319-321.

Goodman, I., Fouts, J.R. Bresnick, E., Menegas, R. and Hitchings, G.H. (1959). A

- mammalian thioglycosidase. *Science.*, 130: 450-451.
- Giamoustaris, A., and Mithen, R. (1996). Genetics of aliphatic glucosinolates. IV. Side-chain modification in *Brassica oleracea*. *Theoretical and Applied Genetics.*, 93: 1006-1010.
- Graham, S., Dayal, H., Swanson, M., Mittelman, A. and Wilkinson, G. (1978). Diet in the epidemiology of cancer of the colon and rectum. *Journal of the National Cancer Institute.*, 61: 709-714.
- Guo, L. and Poulton, J.E. (1994). Partial purification and characterization of *Arabidopsis thaliana* A UDPG: thiohydroximate glucosyltransferase. *Phytochemistry.*, 36: 1133-1138.
- Halkier, B.A. (1999). Glucosinolates. *Naturally Occurring Glycosides: Chemistry Distribution and Biological Properties*, ed R. Ikan. John Wiley & Sons Ltd., 193-223.
- Hartel, F.V. and Brandt, A. (2002). Characterization of a *Brassica napus* myrosinase expressed and secreted by *Pichia pastoris*. *Protein Expr Purif.*, 24(2): 221-226.
- Hawksworth, D.L. (1991). The fungal dimension of biodiversity: magnitude, significance and conservation. *Mycol. Res.*, 95: 641-655.
- Heaney, R.K. and Fenwick, G.R. (1989). The Analysis of Anti-nutritional Factors. *Aspects of Applied Biology.*, 19: 1-20.
- Hecht, A. and Stephen, S. (1999). Chemoprevention of Cancer by Isothiocyanates, Modifiers, of Carcinogen Metabolism. *Am Soc. Of Nut. Sciences.*, 768s-774s.
- Henderson, H.M. and McEwen, T.J. (1972). Effect of ascorbic acid on thioglucosidases from different crucifers. *Phytochemistry.*, 11: 3127-3133.
- Hull, A.K., Vij, R. and Celenza, J.L. (2000). *Arabidopsis* cytochrome P450s that

catalyze the first step of tryptophan-dependent indole-3-acetic acid biosynthesis. *Proc. Natl. Acad. Sci.*, 97: 2379–2384.

- James, D. and Rossiter, J.T. (1991). Development and characteristics of myrosinase in *Brassica napus* during early seedling growth. *Physiol. Plant.*, 82: 163-170.
- Kohlmeier, L. and Su, L. (1997). *Cruciferous* vegetable consumption and colorectal cancer risk: meta-analysis of the epidemiological evidence. *FASEB Journal.*, 11: 329.
- Kune, S., Kune, G.A. and Watson, L.F. (1987). Case-control study of dietary etiological factor: The Melbourne Colorectal Cancer Study. *Nutrition and Cancer.*, 9: 21-42.
- Lenman, M., Falk, A., Xue, J. and Rask, L. (1993). Characterization of a *Brassica napus* myrosinase pseudogene: myrosinases are members of the BGA family of b-glycosidases. *Plant Mol. Biol.*, 21: 463-474.
- Linscheid, M., Wendisch, D. and Strack, D. (1980). The structures of sinapic acid esters and their metabolism in cotyledons of *Raphanus sativus*. *Naturforsch. Teil.*, 35: 907-914.
- Lonnerdal, B. and Janson, J.C. (1973). Studies on myrosinase II: Purification and characterization of a myrosinase from rapeseed (*Brassica napus* L.). *Biochem. Biophys. Acta.*, 315: 412-429.
- MacGibbon, D.B. and Allison, R.M. (1970). A method for the detection of plant glucosinolates (myrosinase). *Phytochem.*, 9: 541-544.
- MacGibbon, D.B. and Beuzenberg, E.J. (1978). Location of glucosinolase in *Brevicoryne brassicae* and *Lipaphis erysimi* (Aphididae). *New Zealand J. Sci.*, 21: 389-392.
- Machlin, S., Mitchell-Olds, T. and Bradley, D. (1993). Sequence of a *Brassica*

campestris myrosinase gene. *Plant Physiol.*, 102: 1359-1360.

Maheshwari, P.N., Stanley, D.W. and Gray, J.I. (1981). Detoxification of rapeseed production. *J. Food Protect.*, 44: 459-470.

Marsh, R.E. and Waser, J. (1970). Refinement of the crystal structure of sinigrin. *Acta Cryst.*, 26: 1030-1037.

McDanell, R., McLean, A.E., Hanley, A.B., Heaney, R.K. and Fenwick, G.R. (1988). Chemical and biological properties of indole glucosinolates (glucobrassicins): a review. *Food Chem Toxicol.*, 26: 59-70.

Michaud, D.C., Spiegelman, D., Clinton, S.K., Rimm, E.B., Willentt, W.C. and Giocannucci, E.L. (1999). Fruit and vegetable intake and incidence of bladder cancer in a male prospective cohort. *Journal of the National Cancer Institute.*, 91: 605-613.

Miller, G.L. (1959). Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Anal. Chem.*, 31: 426-428.

Miller, R.W., VanEtten, C.H., McGrew, C., Wolff, I.A. and Jones, Q. (1962). Amino acid composition of seed meals from forty-one species of *Cruciferae*. *J. Agric. Food Chem.*, 10: 426-430.

Mithen, R., Dekker, M., Verkerk, R., Rabot, S. and Johnson, I.T. (2000). The nutritional significance, biosynthesis and bioavailability of glucosinolates in human foods. *Journal of the Science of Food and Agriculture.*, 84: 967-984.

Morel, J.L. (1997). Bioavailability of trace elements to terrestrial plants- Chapter 6.

Tarradellas J., Bitton G., Rossel (eds). Soil Ecotoxicology, Lewis Publihers, CRC Press., F1: 141-176.

Nakamura, K., Shiraishi, N., Hosoo, S., Sueyoshi, K., Sugimoto, T., Nanmori, T.,

- Nakagawa, H. and Oji, Y. (2002). A protein kinase activated by darkness phosphorylates nitrate reductase in Komatsuna (*Brassica campestris*) leaves. *Physiol. Plant.*, 115: 496-503.
- Nugon-Baudon, L. and Rabot, S. (1994). Effects of the bacterial status of rats on the changes in some liver cytochrome P450 (EC 1.14.14.1) apoproteins consequent to a glucosinolate-rich diet. *Nutrition Research Reviews.*, 7: 205-231.
- Oginsky, E.L., Stein, A.E. and Greer, M.A. (1965). Myrosinase activity in bacteria as demonstrated by the conversion of progoitrin to goitrin. *Soc. Exp. Biol. Med. Proc.*, 119: 360-364.
- Ohtsuru, M., Tsuruo, I. and Hata, T. (1969). Studies of fungous myrosinase part I: Production, Purification and some characteristics. *Agric. Biol. Chem.*, 33: 1309-1314.
- Ohtsuru, M. and Hata, T. (1972). Molecular properties of multiple forms of plant myrosinase. *Agric. Biol. Chem.*, 36: 2495-2503.
- Ohtsuru, M. and Hata, T. (1973). General characteristics of the intracellular myrosinase from *Aspergillus niger*. *Agric. Biol. Chem.*, 37: 2543-2548.
- Ohtsuru, M., Tsuruo, I. and Hata, T. (1973). The production and stability of intracellular myrosinase from *Aspergillus niger*. *Agric. Biol. Chem.*, 37: 967-971.
- Ohtsuru, M. and Kawatani, H. (1979). Studies of the myrosinase from *Wasabia japonica*: Purification and some properties of Wasabi myrosinase. *Agric. Biol. Chem.*, 43: 2249-2255.
- Osbourn, A. (1996). Preformed antimicrobial compounds and plant defense against fungal attack. *Plant Cell.*, 8(3): 1821-1831.

- Palmieri, S., Leoni, O. and Iori, R. (1982). A steady-state kinetic study of myrosinase with direct ultraviolet spectrophotometric assay. *Anal Biochem.*, 123: 320-324.
- Palop, M.L., Smiths, J.P. and Brink, B.T. (1995). Degradation of sinigrin by *Lactobacillus agilis* strain R16. *Int. J. Food Microbiol.*, 26: 219-229.
- Peterson, G.L. (1977). A simplification of the protein assay method of Lowry *et al.* which is more generally applicable. *Anal. Biochem.*, 83: 346-356.
- Petroski, R.J. and Kwolek, W.F. (1985). Interaction of a fungal thioglucoside glucohydrolase and *Cruciferous* plant epithiospecifier protein to form 1-cyanoepithioalkanes: Implications of an allosteric mechanism. *Phytochem.*, 24: 213-216.
- Petrucchioli, M., Piccioni, P., Federici, F. and Polsinelli, M. (1995). Glucose oxidase overproducing mutants of *Penicillium variable* (P16). *FEMS Microbiol. Lett.*, 128: 107-112.
- Pomazi, A., Wittner, A., Pesti, M. and Hornok, L., (1994). A PCR-generated simple RFLP-probe differentiates three distinct groups within *Fusarium oxysporum* f.sp. *pisi*. *Acta Phytopathologica et Entomologica Hungarica.*, 29: 203-213.
- Poulton, J.E. (1990) Cyanogenesis in plants. *Plant Physiol.*, 94: 401-405.
- Prester, T., Fahey, J.W., Holtzclaw, W.D., Abeygunawardana, C., Kachinski, J.L., and Talalay, P. (1996). Comprehensive chromatographic and spectroscopic methods for the separation and identification of intact glucosinolates. *Anal. Biochem.*, 239: 168-179.
- Raper, K.B. and Fennell, I.D. (1973). The genus *Aspergillus*. Huntington, New York. R.E. Krieger Publishing Company, 686 pp.
- Rakariyatham, N. (2000). The value added by-products from mustard essential oil plants. In: Reports to Thailand Research Fund. Chiang Mai. Thailand.

- Rakariyatham, N. and Sakorn, P. (2002). Biodegradation of glucosinolates in brown mustard seed meal (*Brassica juncea*) by *Aspergillus* sp. NR-4201 in liquid and solid-state cultures. *Biodegradation*., 13: 395-399.
- Rask, L., Andreasson, E., Ekbom, B., Eriksson, S., Pontoppidan, B. and Meijer, J. (2000). Myrosinase: gene family evolution and herbivore defense in *Brassicaceae*. *Plant Mol. Biol.*, 42: 93-113.
- Reese, E.T., Clapp, R.C. and Mandels, M. (1958). A thioglucosidase in fungi. *Arch. Biochem. Biophys.*, 75: 228-242.
- Rodman, J.E. (1981). Divergence, convergence, and parallelism in phytochemical characters: the glucosinolate-myrosinase system. In D.A. Young and D. S. Seigler [eds.], *Phytochemistry and angiosperm phylogeny*., 43-79.
- Rodman, J.E. (1991). A Taxonomic analysis of glucosinolate-producing plants, Part 2: Cladistics. *Systematic Botany*., 16: 619-629.
- Rosa, E.A.S., Heaney, R.K., Fenwick, G.R. and Portas, C.A.M. (1997). Glucosinolates in crop plants. *Horticultural Reviews*., 19: 99-215.
- Roy, R. and Das, A., (1978). The mutagenic action of N-methyl, N'-nitro-N-nitroso-guanidine on *Aspergillus niger*. *J. Sci. Cult.*, 43: 461-463.
- Sakorn, P., Rakariyatham, N., Niamsup, H. and Kovitaya, P. (1999). Sinigrin degradation by *Aspergillus* sp. NR-4201 in liquid culture. *Sci. Asia*., 25: 189-194.
- Sakorn, P., Rakariyatham, N., Niamsup, H. and Nongkunsarn, P. (2002). Rapid detection of myrosinase-production fungi: a plate method based on opaque barium sulphate formation. *World J. Microbiol. Biotechnol.*, 18: 73-74.
- Sakushima, A., Ohnishi, S., Kubo, H. and Maoka, T. (1995). Study of sinapinyl but-3-

enylglucosinolate (boreavan A) and related compounds by mass spectrometry.

Phytochemical Analysis., 8(6): 312-315.

Sang, J.P., Minchinton, I.R., Johnstone, P.K., and Truscott, R.J.W. (1984).

Glucosinolate Profiles in the Seed, Root and Leaf Tissue of Cabbage, Mustard, Rapeseed, Radish and Swede. *Can. J. Plant Sci.*, 64: 77-93.

Shapiro, T.A., Fahey, J.W., Wade, K.L., Stephenson, K.K. and Talalay, P. (1998).

Human metabolism and excretion of cancer chemoprotective glucosinolates and isothiocyanates of *Cruciferous* vegetables. *Cancer Epidemiol Biomarkers Prew.*, 7: 1091-2000.

Smiths, J.P., Knol, W. and Bol, J. (1993). Glucosinolate degradation by *Aspergillus calvatus* and *Fusarium oxysporum* in liquid and solid-state fermentation. *Appl. Microbiol. Biotechnol.*, 38: 696-701.

Steinmetz, K.A. and Potter, J.D. (1991). Vegetables, fruit, and cancer. I.

Epidemiology. *Cancer Cause Control.*, 2: 325-357.

Stoewsand, G. (1995). Bioactive organosulfur phytochemicals in *Brassica oleracea* vegetables a review. *Food and Chemical Toxicology.*, 33(6): 537-543.

Stoner, G.D. and Morse, M.A. (1997). Isothiocyanates and plant polyphenols as inhibitors of lung and esophageal cancer. *Canc. Lett.*, 19: 113-119.

Talalay, P. (1999). The war against cancer: new hope. *Proceedings of the American Philosophical Society.*, 143: 52-72.

Tani, N., Ohtsuru, M. and Hata, T. (1974). Isolation of myrosinase producing microorganism. *Agric. Biol. Chem.*, 38: 1617-1622.

Tawfiq, N., Heaney, R.K., Plumb, J.A., Fenwick, G.R., Musk, S.R. and Williamson,

- G. (1995). Dietary glucosinolates as blocking agents against carcinogenesis: glucosinolates breakdown products assessed by induction of quinine reductase activity in murine hep 1c1c7 cells. *Carcinogenesis*, 16: 1191-1194.
- Thangstad, O.P., Evjen, K. and Bones, A. (1991). Immunogold-EM localization of myrosinase in *Brassicaceae*. *Protoplasma*, 161: 84-93.
- Thangstad, O.P., Winge, P., Husebye, H. and Bones, A. (1993). The thioglucoside glucosylhydrolase (myrosinase) gene family in *Brassicaceae*. *Plant Mol Biol.*, 23: 511-524.
- Underhill, E.W. and Kirkland, D.F. (1972). A New Thioglucoside, 2-methylpropylglucosinolate. *Phytochemistry*, 11: 2085-2088.
- Verhoeven, D.T.H., Goldbohm, R.A., van Poppel, G., Verhagen, H. and van den Brandt, P.A. (1996): Epidemiological studies on *Brassica* vegetables and cancer risk. *Cancer Epidemiol. Biomark. Prev.*, 5: 733-748.
- Wallig, M.A., Gould, D.H., Fettman, M.J. and Willhite, C.C. (1988). Comparative toxicities of the naturally occurring nitrile 1-cyano-3,4-epithiobutane and the synthetic nitrile n-valeronitrile in rats: Differences in target organs, metabolism and toxic mechanisms. *Food Chem. Tox.*, 26: 149-157.
- Wilkinson, A.P., Rhodes, M.J.C. and Fenwick, G.R. (1984). Determination of myrosinase (thioglucoside glucosylhydrolase) activity by a spectrophotometric coupled enzyme assay. *Annal. Biochem.*, 139(2): 284-291.
- Wittstock, U. and Halkier, B.A. (2000). Cytochrome P450 CYP79A2 from *Arabidopsis thaliana* L. catalyzes the conversion of l-phenylalanine to phenylacetaldoxime in the biosynthesis of benzylglucosinolate. *J. Biol. Chem.*, 275: 14659–14666.
- Xue, J., Lenman, M., Falk, A. and Rask, L. (1992). The glucosinolate-degrading

enzyme myrosinase in *Brassicaceae* is encoded by a gene family. *Plant Mol. Biol.*, 18: 387-398.

Xue, J., Philgren, U. and Rask, L. (1993). Temporal, cell-specific, and tissue-preferential expression of myrosinase genes during embryo and seedling development in *Sinapis alba*. *Planta.*, 191: 95-101.

Xue, J., Jørgensen, M., Philgren, U. and Rask, L. (1995). The myrosinase gene family in *Arabidopsis thaliana*: gene organization, expression and evolution. *Plant Mol. Biol.*, 27: 911-922.

Zhang, Y. and Talalay, P. (1994). Anticarcinogenic activities of organic isothiocyanates: chemistry and mechanisms. *Cancer Research.*, 54: 1976s-1981s.

Zhang, Y. and Talalay, P. (1998). Mechanisms of differential potencies of isothiocyanates as inducers of anticarcinogenic Phase 2 enzymes. *Cancer Research.*, 58: 4632-4639.