

TABLE OF CONTENTS

Contents	Page
ACKNOWLEDGEMENT	iii
ABSTRACT	iv
TABLE OF CONTENTS	x
LIST OF TABLES	xii
LIST OF FIGURES	xiv
ABBREVIATIONS	xvi
CHAPTER	
I. INTRODUCTION	1
II. LITERATURE REVIEWS	
A. Tuberculosis (TB)	5
B. History of tuberculosis	7
C. Discovery of tubercle bacilli	8
D. Transmission of <i>Mycobacterium tuberculosis</i>	8
E. Treatment and prevention of tuberculosis	9
F. Drug resistant tuberculosis	10
G. Trend of tuberculosis	11
H. <i>Mycobacterium tuberculosis</i>	12
I. Laboratory diagnosis of <i>Mycobacterium tuberculosis</i>	14
J. Apoptosis	16
K. <i>Mycobacterium tuberculosis</i> and its host cell, the macrophages	19
III. MATERIALS AND METHODS	
A. Subjects	23
B. Isolation and purification of human peripheral blood monocytes	23
C. <i>Mycobacterium tuberculosis</i> strains	25
D. Infection of monocyte-derived macrophages with <i>M. tuberculosis</i>	25

E. Apoptosis assay	25
F. Non-specific esterase staining of cells	27
G. The viability of monocytes/macrophages	27
H. Acid-fast stain	28
I. Statistical analysis	28
IV. RESULTS	
A. Isolation and purification of human peripheral blood monocytes	29
B. Infection of monocyte-derived macrophages with <i>M. tuberculosis</i>	30
C. Determination of the apoptotic cell in <i>Mycobacterium tuberculosis</i> -infected macrophages	31
V. DISCUSSION	67
VI. SUMMARY	72
VII. REFERENCES	75
VIII. APPENDICES	83
IX. CURRICULUM VITAE	87

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่

Copyright© by Chiang Mai University

All rights reserved

LIST OF TABLES


Table	Page
1. Number and differential count of white blood cell in peripheral blood from normal persons	35
2. Number of white blood cells, viable count and non-specific esterase positive cell in peripheral blood mononuclear cells (PBMC) and adherent cells in normal persons	36
3. Number and differential count of white blood cell in peripheral blood from tuberculosis patients	37
4. Number of white blood cells, viable count and non-specific esterase positive cell in peripheral blood mononuclear cells (PBMC) and adherent cells in tuberculosis patients	38
5. Percent recovery of adherent cells compared with number of monocytes in whole blood and the number of non-specific esterase positive cells in peripheral blood mononuclear cell of normal person	39
6. Percent recovery of adherent cells with the number of monocytes in whole blood and the number of non-specific esterase positive cells in peripheral blood mononuclear cell of tuberculosis patients	40
7. Percent phagocytosis of macrophages from normal persons to <i>M. tuberculosis</i> H37Ra at various multiplicity of infections (MOI) and incubation times	41
8. Percent phagocytosis of macrophages from normal persons to <i>M. tuberculosis</i> H37Ra at the multiplicity of infection 10 in 10% pooled human AB serum with variuos incubation times	42
9. Percent phagocytosis of macrophages from normal persons to <i>M. tuberculosis</i> H37Ra and <i>M. tuberculosis</i> H37Rv in 10% pooled human AB serum	43
10. Percent phagocytosis of macrophages from tuberculosis patients to <i>M. tuberculosis</i> H37Ra and <i>M. tuberculosis</i> H37Rv in 10% pooled	

human AB serum	44
11. Percentage of apoptotic cells of <i>M. tuberculosis</i> H37Ra-infected macrophages from normal persons	45
12. Percentage of apoptotic cells of <i>M. tuberculosis</i> H37Ra-infected macrophages or <i>M. tuberculosis</i> H37Rv-infected macrophages from normal persons after 48 hours of incubation	46
13. Percentage of apoptosis of <i>M. tuberculosis</i> H37Ra-infected macrophages or <i>M. tuberculosis</i> H37Rv-infected macrophages from tuberculosis patients	47

LIST OF FIGURES

Figure	Page
1. Non-specific esterase positive cell in adherent cells of normal person and tuberculosis patient	48
2. Phagocytosis of normal macrophages to <i>M. tuberculosis</i> H37Ra in various MOIs and incubation times	49
3. Comparison of percent phagocytosis of macrophages from normal persons to <i>M. tuberculosis</i> H37Ra at various MOI and incubation time	50
4. Percent phagocytosis of macrophages from normal persons to <i>M. tuberculosis</i> H37Ra in 10 percent pooled human AB serum	51
5. Percent phagocytosis of macrophages from normal persons to <i>M. tuberculosis</i> H37Ra in 10 percent pooled human AB serum	52
6. Comparison of percent phagocytosis of <i>M. tuberculosis</i> H37Ra and <i>M. tuberculosis</i> H37Rv in macrophages of normal persons	53
7. Comparison of percent phagocytosis of <i>M. tuberculosis</i> H37Ra and <i>M. tuberculosis</i> H37Rv in macrophages of normal persons and tuberculosis patients	54
8. Apoptosis of Actinomycin D (50µg/ml)-treated macrophages for 19h	55
9. Percent apoptosis of <i>Mycobacterium tuberculosis</i> H37Ra-infected macrophages from normal persons at 48, 72 and 120h	56
10. Comparison of percent apoptosis of <i>M. tuberculosis</i> H37Ra-infected macrophages from normal persons with or without pooled human AB serum	57
11. Comparison of percent apoptosis of <i>M. tuberculosis</i> H37Ra or <i>M. tuberculosis</i> H37Rv-infected macrophages in normal persons	58
12. Apoptosis of <i>M. tuberculosis</i> H37Ra or <i>M. tuberculosis</i> H37Rv-infected macrophages in normal persons	59
13. Comparison of percent apoptosis of <i>M. tuberculosis</i> H37Ra or	

<i>M. tuberculosis</i> H37Rv-infected macrophages in tuberculosis patients	62
14. Apoptosis of <i>M. tuberculosis</i> H37Ra or <i>M. tuberculosis</i> H37Rv-infected macrophages in tuberculosis patients	63
15. Comparison of percent apoptosis of <i>M. tuberculosis</i> H37Ra or <i>M. tuberculosis</i> H37Rv-infected macrophages in normal persons and tuberculosis patients	66




ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
 Copyright© by Chiang Mai University
 All rights reserved

ABBREVIATIONS

ADC	Albumin-dextrose-catalase
AFB	Acid-fast bacilli
AIDS	Acquired immunodeficiency syndrome
ATP	Adenosine triphosphate
BCG	Bacillus Calmette Guerin
bp	Base pair (s)
<i>C. albicans</i>	<i>Candida albicans</i>
Ca ²⁺	Calcium ion
cm	Centimeter
CNS	Central nervous system
CO ₂	Carbon dioxide
CR	Complement receptor
Cu	Copper
Da	Dalton
DD	Death domain
DED	Death effector domain
DOTS	Directly Observed Treatment, Short-course
EDTA	Ethylenediamine tetraacetic acid
ER	Endoplasmic reticulum
FADD	Fas-associated death domain-containing protein
h	Hour (s)
H ₂ O ₂	Hydrogen peroxide
HIV	Human immunodeficiency virus
IFN-γ	Gamma interferon
IL	Interleukin
k	Kilo
LAM	Lipoarabinomannan
LJ	Lowenstein-Jensen

M	Molar
<i>M. bovis</i>	<i>Mycobacterium bovis</i>
<i>M. smegmatis</i>	<i>Mycobacterium smegmatis</i>
<i>M. tuberculosis</i>	<i>Mycobacterium tuberculosis</i>
MBP	Mannose binding protein
MDR-TB	Multidrug resistant tuberculosis
mg	Milligram
ml	Milliliter
min	Minute
mm	Millimeter
MOI	Multiplicity of infection
NO	Nitric oxide
OADC	Oleic acid albumin-dextrose catalase
°C	Degree celsius
ORF	Open reading frame
PBMC	Peripheral blood mononuclear cell
PBS	Phosphate buffer saline
PCR	Polymerase chain reaction
PI	Propidium iodide
PS	Phosphatidyl serine
RFLP	Restriction fragment length polymorphism
RNIs	Reactive nitrogen intermediates
ROIs	Reactive oxygen intermediates
SE	Standard error of the mean
Sp-A	Surfactant protein A
TB	Tuberculosis
TLR	Toll-like receptor
TNF	Tumor necrosis factor
TNFR	Tumor necrosis factor receptor
TRAF	Tumor necrosis factor receptor-associated factor
TRAIL	Tumor necrosis factor-related apoptosis inducing ligand

UV	Ultraviolet
w/v	Weight/volume
WBC	White blood cell
WHO	World Health Organization
Zn	Zinc
μ	Micro
μg	Microgram
μl	Microliter



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
 Copyright© by Chiang Mai University
 All rights reserved