

TABLE OF CONTENTS

Content	Page
ACKNOWLEDGEMENTS	iii
ABSTRACT	iv
LIST OF TABLES	x
LIST OF ILLUSTRATIONS	xii
CHAPTER I INTRODUCTION	1
Principle, theory, and rationale	1
The objective of the study	2
The hypothesis	2
Anticipated benefits	3
Scope of the study	3
CHAPTER II LITERATURE REVIEW	4
I. Effect of orthodontic tooth movement on periodontium	4
II. Gingival crevicular fluid (GCF)	6
III. Glycosaminoglycans (GAGs)	7
IV. Hyaluronic acid (HA)	8
V. ELISA-liked assay for HA	12
CHAPTER III MATERIALS AND METHODS	14
Materials	14
Methods	19
Statistical analyses	21

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่

Copyright © by Chiang Mai University

All rights reserved

Content	Page
CHAPTER IV RESULTS	22
Part I: The descriptions of the patients	22
Part II: The clinical scores	23
Part III: The changes in HA level per unit protein	24
Part IV: Comparisons of the HA level per unit protein during the leveling phase (L-phase), the first eight weeks of canine movement phase (M-phase), and after complete orthodontic canine movement phase (S-phase)	28
Part V: Comparisons of the HA level per unit protein between canines and incisors in every treatment phase	30
CHAPTER V DISCUSSION	32
Limitation of this study	37
Suggestions for further study	39
CHAPTER VI CONCLUSION	40
BIBLIOGRAPHY	41


APPENDIX	46
----------	----

CIRRICULUM VITAE	55
------------------	----

LIST OF TABLES

Table	Page
3.1 A score used to evaluate Gingival Index (GI).	16
3.2 A score used to evaluate Plaque Index (PI).	16
4.1 A summary of the patients' profiles in this study: patients' code, tooth number, age, and sex.	22
4.2 The means (standard deviations) of plaque index (PI) and gingival index (GI) before, during, and after complete orthodontic canine movements.	23
4.3 The mean rank of plaque index (PI) and gingival index (GI) scores of all four patients before, during, and after complete orthodontic canine movements (n = 4).	23
4.4 The non-parametric statistical analysis (Friedman Test) of plaque index (PI) and gingival index (GI) scores.	24
4.5 The medians of HA level per unit protein (ng/ μ g) of canines (n=7) and incisors (n=3) during the leveling phase, the first eight weeks of canine movement phase (M0, M1, M2, M3, ..., and M8), and after complete orthodontic canine movement phase (S1, S2, S3, ..., and S8).	28
4.6 The non-parametric statistical analysis (Friedman Test) of HA level per unit protein of canines and incisors.	30
4.7 The non-parametric statistical analysis (Mann-Whitney U Test) of the medians of HA level per unit protein between canines (n = 7) and incisors (n = 3) in every treatment phase.	31

Table	Page
A.1 Protein levels, HA levels, HA levels per unit protein of GCF samples from C11, C13	47
A.2 Protein levels, HA levels, HA levels per unit protein of GCF samples from D11, D13, and D23	49
A.3 Protein levels, HA levels, HA levels per unit protein of GCF samples from E11, E13, and E23	51
A.4 Protein levels, HA levels, HA levels per unit protein of GCF samples from H13, H23	53



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright © by Chiang Mai University
All rights reserved

LIST OF ILLUSTRATIONS

Figure	Page
4.1 The duration of a longitudinal study.	24
4.2 Intraoral photographs	25
4.3 A linear graph demonstrated HA level per unit protein (ng/ μ g) of tooth C11 (Blue) and C13 (Pink). The red arrows indicated the time when a new closed coil spring was changed and the force was therefore applied to the experimental tooth, C13.	26
4.4 A linear graph demonstrated HA level per unit protein (ng/ μ g) of tooth D11 (Blue), D13 (Pink), and D23 (Green). The red arrows indicated the time when a new closed coil spring was changed and the force was therefore applied to the experimental teeth, D13 and D23.	27
4.5 A Linear graph demonstrated the medians of HA level per unit protein (ng/ μ g) of canines (n=7, Blue) and incisors (n=3, Pink) in the leveling phase, the first eight weeks of canine movement phase (M0, M1, M2, M3, ..., and M8), and the complete canine movement phase (S1, S2, S3, ..., and S8).	29