

REFERENCES

- Anon. (2001): Commission Decision 2001/ 471/EC). *Official Journal of the European Communities* **165**, 48– 53.
- Anon. (2002): Enumeration of *Enterobacteriaceae* by the colony count technique. <http://www.hpa-standardmethods.org.uk/documents/food/pdf/f23.pdf>
- Aoust, J-Y. D., Purvis, U. (1998): Isolation and Identification of *Salmonella* from foods. http://www.hc-sc.gc.ca/food-aliment/mh-dm/mhe-dme/compendium/volume_2/e_mfhpb2001.html
- Baggesen, D.L., Wegener, H.C., Bager, F., Stege, H., Christensen, J. (1996): Herd prevalence of *Salmonella* enterica infections in Danish slaughter pigs determined by microbiological testing. *Prevent. Vet. Med.* **26**, 201-213.
- Berends, B. R., Urlings, H. A. P., Snijders, J. M. A., Van Knapen, F. (1996): Identification and quantification of risk factors in animal management and transport regarding *Salmonella* spp. in pigs. *Int. J. Food Microbiol.* **30**, 37-53
- Borch, E., Nesbakken, T., Christensen, H., (1996): Hazard identification in swine slaughter with respect to foodborne bacteria. *Int. J. Food Microbiol.* **30**, 9-25.
- Botteldoorn, N., Heyndrickx, M., Rijpens, N., Grijspeerdt, K., Herman, L. (2003): *Salmonella* on pigs and cross contamination in slaughterhouse. *J. appl. Microbiol.* **95**, 891-903.
- Byrne, B., Dunne, G., Lyng, J., Bolton, D.J. (2005): Microbiological carcass sampling methods to achieve compliance with 2001/471/EC and new hygiene regulations. *Res. Microbiol.* **156**, 104–106.

CDC. (2003): Foodborne Diseases Active Surveillance Network (FoodNet) Emerging Infections Program, Report on Foodborne Pathogens.

<http://www.cdc.gov/foodnet/pub/publications/2005/FNsurv2003.pdf>

CDC. (2002): Memorandum To: State and Territorial Public Health Laboratory Directors State and Territorial Epidemiologists. Subject: Adoption of the Kauffmann-White scheme for designation of *Salmonella* serotypes.

Crowley, H., Cagney, C., Sheridan, J.J. Anderson, W., McDowll, D.A., Blair, I.S., Bishop, R.H., Duffy, G. (2005): *Enterobacteriaceae* in beef products from retail outlets in the Republic of Ireland and comparison of the presence and counts of *E. Coli*O157:H7 in these products. *Food Microbiol.* **22**, 409-414.

Currier, M., Singleton, M.J., Lee, J., Lee, D.R. (1986): *Salmonella* in swine at slaughter: incidence and serovar distribution at different seasons. *J. Food Prot.* **49**, 366-368.

D' Aoust J-Y, John Maurer, Bailey J., S. (2001): *Salmonella* Species, p. 141-178. In Michael P. Doyle, Larry R. Beuchat, Thamas J. Montville (ed.), Food Microbiology. ASM press Washington D.C.

D' Aoust, J-Y., Maurer, J., Bailey, J.S. (2001): *Salmonella* Species. In: Doyle, M.P., Beuchat, L.R., Montville, T.J. (ed.): Food Microbiology. 2 ed. Washington, D.C.: ASM Press. pp. 141-178.

Dicson, J. S. Hurd, H.S., Rostagno, M.H. (2005): *Salmonella* in the pork production chain a Review article.

<http://www.porkscience.org/documents/Research/review.pdf>

Danilo, M.A., Lo Fo Wong, Tine Hald (2000): *Salmonella* in Pork (SALINPORK): Pre-harvest and Harvest Control Options based on Epidemiologic, Diagnostic

and Economic Research. Final report Contract no: FAIR1 CT95-0400. Copenhagen, Denmark.

Fedorka-Cray, P.J., T.J. Gray, and C. Wray (2000): *Salmonella* Infections in Pigs. In: Wray, C., and A. Wray (eds); *Salmonella* in Domestic Animals. CABI Publishing, Wallingford, UK, pp. 191-207.

Finlay, R.C., Mann, E.D., Horning, J.L. (1986): Prevalence of *Salmonella* and *Campylobacter* contamination in Manitoba swine carcasses. *Can. Vet. J.* 27, 185-187.

Gebreyes, W. A., Davies . R., Turkson, P-K., Morrow, W. E. M., Fink, J.A., Alter, C. (2004): *Salmonella* enterica Serovars from Pigs on Farms and after Slaughter and Validity of Using Bacteriologic Data To Define Herd *Salmonella* Status. *J. Food Protect.* 67 (4), 691-697.

Gill, C. O., Jones, T. (1997): Assessment of the hygienic characteristics of a process for dressing pasteurized pig carcasses. *J. Food Microbiol.* 14, 81-91.

Gill, C. O., Lander, C. (2004): Proximate sources of bacteria on boneless prepared from routinely processes and detained carcasses at a pork packing plant. *Int. J. Food Microbiol.* 97, 171-178.

Gill, C. O., Bryant, J. (1992): The contamination of pork with spoilage bacteria during commercial dressing, chilling and cutting of pig carcasses. *Int. J. Food Microbiol.* 16, 51-62.

Gill, C. O., Mc Ginnis, J. C., Bryant, J. (1998): Microbial contamination of meat during the skinning of beef carcass hind quarters at three slaughtering plants. *Int. J. Food Microbiol.* 42, 175-184.

- Gill, C.O. (2000): HACCP in primary processing: red meat. In: Martyn Brown, HACCP in the Meat Industry CRC Press, pp. 81-122.
- Gill, C.O., Bryant, J. (1993): The presence of *Escherichia coli*, *Salmonella* and *Campylobacter* in pig carcass dehairing equipment. *Food Microbiol.* **10**, 337-344.
- Goodfellow, S.J. (1995): Implementation of the HACCP program by the meat and poultry slaughters. In: Pearson, A.M., and Dutson, T.R., HACCP in Meat, Poultry and Fish Processing. Advance in Meat Research vol.10, pp. 58-71.
- Gray, J.T., Stabel T.J., Fedorka-Gray P.J. (1996): Effect of dose on the immune response and persistence of *Salmonella choleraesuis* infection in swine. *Am. J. Vet. Res.* **57**, 313-319.
- Hinton M., (2000): Microbial control in the meat industry. FLAIR-FLOW EUROPE TECHNICAL MANUAL F-FE 379A/00.
- Holt, J.G., Krieg, N.R., Sneath, P.H.A., Staley, J.T and Williams, S.T. (2000): *Bergey's Manual Of Determinative Bacteriology*. 9th ed. Printed by Lippincott&Wilkins, pp. 175-242.
- Isaacson, R. E., Firkins, L. D., Weigel, R. M., (1999): Effect of transportation and feed withdrawal on shedding of *Salmonella* Typhimurium among experimentally infected pigs. *Am. J. Vet. Res.* **60**, 1155-1158.
- ISO 6579 (2002): International Standard: Microbiology of food and animal feeding stuff-Horizontal method for the detection of *Salmonella* spp. Geneva ISO copyright office.

- Käsbohrer, A., Protz, d., Helmuth, R., Nöckler, K., Blaha, T., Conraths, F.J., Geue, L. (2000): *Salmonella* in slaughter pigs of German origin: An epidemiological study. *Eur. J. Epidemiol.* **16**, 141-146.
- Keteran, K., Brown, J., Shotts, E.B. (1982): *Salmonella* in the mesenteric lymph nodes of healthy sows and hogs. *Am. J. Vet. Res.* **43**, 706-707.
- Krieg, N. R., Holt, J. G. (1984): *Bergey's Manual of Systematic Microbiology*. Volum1. Williams & Wilkins. pp. 427-458.
- Lammerding, A.M., Garcia, M.M., Mann, E.D., Robinson, Y., Dorward, W.J., Truscott, R.B., Tittiger, F. (1988): Prevalence of *Salmonella* and thermophilic *Campylobacter* in fresh pork, beef, veal and poultry in Canada. *J. Food Prot.* **51**, 47-52.
- Lazaro, N.S., Tibana, A., Hofer, E., 1997. *Salmonella* spp. in healthy swine and in abattoir environments in Brazil. *J. Food Prot.* **60**, 1029-1033.
- Letellier, A., Messier, S., Quessy, S. (1999): Prevalence of *Salmonella* spp. and *Yersinia enterocolitica* in finishing swine at Canadian abattoirs. *J. Food Prot.* **62**, 22-25.
- Lo Fo Wong, D.M.A., Hald, T., van der Wolf, P.J., Swanenburg, M., (2002): Epidemiology and control measures for *Salmonella* in pigs and pork. *Livestock Production Science*, **76**, 215-222.
- Mafu, A.A., Higgins, R., Nadeau, M., Cousineau, G. (1989): The incidence of *Salmonella*, *Campylobacter*, and *Yersinia enterocolitica* in swine carcasses and the slaughterhouse environment. *J. Food Prot.* **52**, 642-645.

- McEvoy, J.M., Sheridan, J.J., Blair, I.S., McDowell, D.A. (2004): Microbial contamination on beef in relation to hygiene assessment based on criteria used in EU Decision 2001/471/EC. *Int. J. Microbiol.* **92**, 217–225.
- Miliotis, M.D., Bier, J.W., (2003): International Handbook of Food Pathogens. Marcell Dekker Inc. Pp. 137-165.
- Morgan, I.R., Krautil, F.L., Craven, J.A. (1987): Bacterial populations on dressed pig carcasses. *Epldemiol. Infect.* **98**, 15-24.
- Morse, J.W., Hird, D.W., (1984): Bacteria isolated from lymph nodes of California slaughter swine. *Am. J. Vet. Res.* **45**, 1648–1649.
- Mossel, D.A.A., Weenk, G.H., Morris, G.P., Corry B. Struijk (1998): Identification, assessment and management of food-related microbiological hazards: historical, fundamental and psycho-social essentials *Int. J. Food Microbiol.* **39**, 19–51.
- Narashimha Rao, D., Mair, K.K.S., Sakhare, P.Z. (1998): Meat microbiology and spoilage in tropical countries. In: Anrew Daviea and Rond Board The Microbiology of Meat and Poultry. Published by Blackie Academic and Professional, pp. 220-265.
- Nel, S., Lues, J.F.R., Buys, E.M., Venter, P. (2004): Bacterial populations associated with meat from the deboning room of a high throughput red meat abattoir. *Meat Science*, **66**, 667–674.
- Nortje, G.L., Nel, L., Jordaan, E., Badenhorst, K., Goedhart, E., Holzapfel, W.H., (1990): The aerobic psychrotrophic populations on meat and meat contact surfaces in a meat production system and on meat stored at chill temperatures. *J. Appl. Bacteriol.* **68**, 335–344.

- Oosterom, J., Dekker, R., de Wilde, G.J.A., van Kempen-de Troye, F., Engels, G.B., (1985): Prevalence of *Campylobacter jejuni* and *Salmonella* during pig slaughtering. *Vet. Q.* **7**, 31–34.
- Oosterom, J., Notermans, S. (1983): Further studies on the practicability of fattening and slaughtering pigs free from *Salmonella*. *Tijdschr. Diergeneeskd.* **108**, 177–186.
- Palumbo, S. A., Klein, P., Capra, J., Eblen, S. and A Miller, J. (1999): Comparison of excision and swabbing sampling methods to determine the microbiological quality of swine carcass surfaces. *Food Microb.* **16**, 459-464.
- Pearce, R.A., Bolton, D.J., Sheridan, J.J. McDowel, D.A. Blair, I.S. Harrington, D. (2004): Study to determine critical control points in pork slaughter hazard analysis and critical control points. *Int. J. Food Microbiol.* **90**, 331-339.
- Pearson, A. M., Dutson, T. R. (1995): HACCP in Meat, Poultry and Fish Processing. 1st ed. Blackie Academic & Professional.
- Popoff, M. Y., Le Minor. L. (1997): Antigenic formulas of the *Salmonella* serovars, 7th revision. World Health Organization Collaborating Centre for Reference and Research on *Salmonella*, Pasteur Institute, Paris, France.
- Popoff, M.Y., Bockemühl, J., Brenner, (2000): Supplement 1998 (no. 42) to the Kauffmann–White scheme, *Res. Microbiol.* **151**(1), 63-5.
- Popoff, M.Y., Bockemühl, J., Gheesling, L.L. (2004): Supplement 2002 (no. 46) to the Kauffmann–White scheme, *Res. Microbiol.* **155**, 568–570.
- Qiongzen Li, Julie S. Sherwood, Catherine M. Logue (2004): The prevalence of *Listeria*, *Salmonella*, *Escherichia coli* and *E. coli* O157:H7 on bison carcasses during processing. *Food Microbiol.* **21**, 791–799.

- Quinn, P. J., Carter, M. E., Markey, B. K., Carter, G. R. (1998): Clinical Veterinary Microbiology. Mosby publications.
- Reed, W.M., Olander, H.J., Thacker, H.L. (1986): Studies on the pathogenesis of *Salmonella typhimurium* and *Salmonella choleraesuis* var *kunzendorf* infection in weaning pigs. *Am. J. Vet. Res.* **47**, 75-83.
- Reij, M.W., Den Aantrekker, E.D. (2004): Review article Recontamination as a source of pathogens in processed foods. *Int. J. Food Microbiol.* **91**, 1– 11.
- Rho, M-J., Chung, M-S., Lee, J-H., Park J. (2001): Monitoring of microbial hazards at farms, slaughterhouses and processing lines of swine in Korea. *J. Food Protect.* **64** (9), 1388-1391.
- Rivas, T., Vizcaino, J.A., Herrera, J.F. (2000): Microbial contamination of carcasses and equipment from an Iberian pig slaughterhouse. *J. Food Protect.* **63** (12), 1670-1675.
- Roberts, T.A., Hudson, W.R., Whelehan, O.P. (1944): Number and Distribution of bacteria on some beef carcasses at selected abattoirs in some member of the European communities. *Meat science* **11**, 191-205.
- Rostagno, M.H., Hurd, H.S., McKean, J.D., Ziemer, C.J., Gailey, J.K. and Leyte, R.C. (2003): Preslaughter Holding Environment in Pork Plants is Highly Contaminated with *Salmonella enterica*. *Appl. Envir. Microbiol.* **69** (8), 4489-4494.
- Sammarco, M.L., Ripabelli, G., Ruberto, A., Iannitto, G., Grasso, G.M. (1997): Prevalence of *Salmonellae*, *Listeriae*, and *Yersinia* in the slaughterhouse environment and on work surfaces, equipment and workers. *J. Food Protect.* **60**, 367–371.

Schwartz, K.J. (1998): Salmonellosis. In: Straw, B.E., D'Allaire, S., Mengeling, W.L., Taylor, D.J. (Eds.), *Disease of Swine*. Blackwell Science, Oxford, pp. 535-551.

SIFIN (2004): Screening and Serotyping of Salmonellae with Test Reagents Based Primarily on Monoclonal Antibodies. A Technical Information Booklet.

Sofos, J.N., Kochevar, S.L., Bellinger, G.R., Buege D.R., Hancock, D.D., Ingham, S.C., Morgan, J.B., Reagan, G.O. (1999): Sources and extent of microbial contamination of beef carcasses in seven United States slaughtering plants. *J. Food Protect.* **62**, 140-145.

Søren Kranker, Lis Alban, Jaap Boes, Jan Dahl (2003): Longitudinal Study of *Salmonella enterica* Serotype Typhimurium Infection in Three Danish Farrow-to-Finish Swine Herds. *J. Clinic. Microbiol.* **41**(6), 2282–2288.

Swanenburg, M., Urlings, H.A.P., Snijders, J.M.A., Keuzenkamp, D.A., van Knapen, F. (2001a): *Salmonella* in slaughter pigs: prevalence, serotypes and critical control points during slaughter in two slaughterhouses *Int. J. Food Microbiol.* **70**, 243–254.

Swanenburg, M., van der Wolf, P.J., Urlings, H.A.P., Snijders, J.M.A., van Knapen, F. (2001b): *Salmonella* in slaughter pigs: the effect of logistic slaughter procedures of pigs on the prevalence of *Salmonella* in pork. *Int. J. Food Microbiol.* **70**, 231–242.

Tran, T.P., Ly, T.L., Nguyen, T.T., Akiba, M., Ogasawara, N., Shinoda, D., Okatani, T.A., Hayashidani, H. (2004): Prevalence of *Salmonella* spp. in pigs, chickens and ducks in the Mekong Delta, Vietnam. *J. Vet. Med. Sci.* **66** (8), 1011- 1014

Vaderlinde P.B., Shay B. and Murry J. (1999): Microbial quality of Australian sheep meat. *J. Food Protect*, **62**, 3880-3885.

- Van der Wolf, P.J., Bongers, J.H., Elbers, A.R.W., Franssen, F.M.M.C., Hunneman, W.A., van Exsela, A.C.A. and Tielena, M.J.M. (1999): *Salmonella* infections in finishing pigs in The Netherlands: bacteriological herd prevalence, serogroup and antibiotic resistance of isolates and risk factors for infection. *Vet. Microbiol.* **67**, 263-275.
- Van der Wolf, P.J., Wolbers, W.B., Elbers, A.R.W., van der Heijden, H.M.J.F., Koppen, J.M.C.C., Hunneman, W.A., van Schie, F.W. and Tielens, M.J.M. (2001): Herd level husbandry factors associated with the serological *Salmonella* prevalence in finishing pig herds in The Netherlands. *Vet. Microbiol.* **78**, 205-219.
- Warriner, K., Aldsworth, T. G., Kaur, S., and Dodd, C.E.R. (2002): Cross-contamination of carcasses and equipment during pork processing. *J. Appl. Microbiol.* **93**, 69-177.
- WHO (1995): Surveillance Programme. Sixth Report of WHO Surveillance Programme for Control of Foodborne Infections and Intoxications in Europe. FAO/WHO Collaborating Centre for Research and Training in Food Hygiene and Zoonoses, Berlin.
- WHO (2005): Fact sheet No. 139.
<http://www.who.int/mediacentre/factsheets/fs139/en/>
- Wilcock, B.P., Schwatz, K (1992): Salmonellosis. In: LEMAN, A.D., B.E. STRAW, W.E. MENGELING, S. D'ALLAIRE, and D.J. TAYLOR (eds), Diseases of Swine, 7th edition Iowa State University Press, Ames, Iowa, pp. 570-583.
- Zweifel, C., Baltzer, D. and Stephan, R., (2005): Microbiological contamination of cattle and pig carcasses at five abattoirs determined by swab sampling in accordance with EU Decision 2001/471/EC. *Meat Science*, **69**, 559-566.