TABLE OF CONTENTS

		Page
Ack	knowledgements 82226	iii
Abs	stract (English)	v
Abs	stract (Thai)	viii
List	t of tables	xvi
List	t of illustrations	xviii
Abl	breviations and symbols	xxiii
Cha	apter 1 General introduction	1
Cha	apter 2 Literature review	
2.1	Definition, distribution and habitats of Dracaena lourieri and	5
	Pandanus spp.	
2.2	Importance of Dracaena lourieri and Pandanus spp.	6
2.3	Fungal life strategies (endophytes, parasites and saprobes)	8
2.4	Fungal classification	10
2.5	Ecology and distribution of fungi	10
2.6	Fungal succession	14
2.7	Importance of fungi	15
2.8	Fungi on Pandanaceae and fungi worldwide	16
2.9	Fungi in Thailand	19
2.10	O Phylogenetic relationships of fungi	
Cha	apter 3 Saprobic and pathogenic fungi on Dracaena and Pandanus	
3.1	Introduction	27

		Page
3.2	Materials and methods	27
	3.2.1 Diversity and ecology of saprobic fungi on <i>Dracaena</i> and <i>Pandanus</i>	27
	3.2.2 Fungal succession on <i>Pandanus penetrans</i> leaves	28
	3.2.3 Pathogenic fungi on <i>Dracaena</i> and <i>Pandanus</i> leaves	30
3.3	Results	31
	3.3.1 Diversity and ecology of saprobes	31
	3.3.1A Determination of sample size	31
	3.3.1B Fungal taxonomic composition	32
	3.3.1C Effect of hosts and their habitats on fungal communities	38
	3.3.1D Fungal occurrence on Pandanus penetrans leaves from	40
	different sites and the effect of the stage of decayed	
	3.3.1E Fungal occurrence on different parts of Pandanus	41
	odoratissimus and P. penetrans	
	3.3.1F Seasonal pattern of fungal occurrences on Pandanus	42
	penetrans leaves	
	3.3.2 Fungal succession on Pandanus penetrans leaves	44
	3.3.2A Determination of sample size	44
	3.3.2B Fungal taxonomic composition on Pandanus penetrans leaves	44
	and effect of stages of decay	
	3.3.3 Pathogenic fungi on <i>Dracaena</i> and <i>Pandanus</i> leaves	51
	3.3.3A Collection of Dracaena and Pandanus diseases	51
	3.3.3B Pathogenicity tests	60
3.4	Discussion	63
	3.4.1 Bioiversity of saprobic fungi on Dracaena loureiri and Pandanus spp.	63
	3.4.2 Fungal occurrence on different tissues of <i>Pandanus odoratissimus</i> and <i>P. penetrans</i>	66
	3.4.3 Seasonal pattern of fungi on <i>Pandanus penetrans</i>	66

P	age
3.4.4 Fungal succession on Pandanus penetrans leaves	67
3.4.5 Pathogenic fungi on leaves of <i>Dracaena lourieri</i> and <i>Pandanus</i> spp.	68
Chapter 4 Saprobic and pathogenic fungi on <i>Dracaena</i> and <i>Pandanus</i>	
4.1 Introduction	69
4.2 Descriptions and photographic figures of anamorphic fungi	70
Canalisporium	70
Dictyochaeta	73
Ellisembia	75
Exerohilum	76
Memnoniella	78
Myrothecium	80
Paecilomyces	83
Periconia	85
Periconiella	88
Phaeostalagmus	89
Phialocephala	91
Sporidesmium	93
Stachybotrys	94
Trichothecium	98
Veronaea	99
Zygosporium	101

		Page
4.3	Descriptions and photographic figures of Ascomycetes	107
	Astrosphaeriella (Melanommataceae: Dothideales)	107
	Chaetosphaeria (Chaetomiaceae: Sordariales)	109
	Emericella (Trichocomaceae: Eurotiales)	111
	Linocarpon (Hyponectriaceae: Incertae Sedis)	112
	Melanochaeta (Chaetosphaeriaceae: Sordariales)	120
	Microthyrium (Microthyriaceae: Dothideales)	122
	Ophioceras (Magnaporthaceae: Incertae Sedis)	125
	Ornatispora (Incertae Sedis: Incertae Sedis)	3 131
	Oxydothis (Incertae Sedis: Incertae Sedis)	133
	Paraphaeosphaeria (Phaeosphaeriaceae: Pleosporales)	137
	Phaeonectriella (Halosphaeriaceae: Halosphaeriales)	140
	Phomatospora (Incertae Sedis: Xylariales)	143
	Pseudohalonectria (Magnaporthaceae: Incertae Sedis)	145
	Sordaria (Sordariaceae: Sordariales)	149
	Tubeufia (Tubeufiaceae: Pleosporales)	150
Cha	apter 5 Molecular Phylogeny of Magnaporthaceae (Sordariomycete	s):
	Implication on taxonomy and phylogenetic placement	
5.1	Introduction	153
5.2	Materials and methods	155
	5.2.1 Fungal cultures and DNA extraction	155
	5.2.2 PCR amplification and sequencing of 28S and 18S rDNA	ers 1 ₁₅₆
	5.2.3 Phylogenetic analyses	159

		Page
5.3	Results	160
	5.3.1 Phylogenies based on 28S rDNA dataset (dataset I)	160
	5.3.2 Phylogenies based on 18S rDNA dataset (dataset II)	161
	5.3.3 Phylogenies based on 18S rDNA dataset (dataset III)	161
	5.3.4 Phylogenies based on the combined dataset (dataset IV)	162
5.4	Discussion	166
	5.4.1 Ordinal classification of Magnaporthaceae	166
	5.4.2 Intergeneric relationships within Magnaporthaceae	168
	5.4.3 Placement of recently described taxa in Magnaporthaceae and	171
	Ophioceras chiangdaoensis sp. nov.	
	5.4.4 Magnaporthaceae and their anamorphs	171
	5.4.5 Phylogenies of recently described taxa	173
Cha	apter 6 General discussion and conclusion	
6.1	Bioiversity of pathogenic and saprobic fungi on Dracaena and Pandanus	177
	The relationships between pathogenic and saprobic fungi on <i>Dracaena</i> and <i>Pandanus</i>	179
	Molecular and morphological characters of fungi in family	179
	Magnaporthaceae	179
6.4	Future work	180
	erences 1811 by Chiang Mai Univer	182
App	pendix A	214
Cur	riculum Vitae	217

xvi

LIST OF TABLES

Tal	ole	Page
2.1	Saprobic fungi showing fungal recurrence on different tissues	12
2.2	Number of fungi and myxomycetes published as occurring on the	17
	Pandanaceae (McKenzie et al. 2002)	
2.3	New fungi described from the Pandanaceae since McKenzie and Hyde 1996	18
2.4	The most common fungal genera recorded on various monocotylenonous hosts	20
2.5	New selected fungi described from various host/habitat in Thailand since 1998	21
3.1	Collection details of saprobic fungi study	28
3.2	Frequency and overall percentage occurrence of fungal taxa occurred	33
	on Dracaena lourieri, Pandanus amaryllifolius, P. odoratissimus and P. penetrans	
	Comparison of the most common fungal taxa recorded on naturally samples and baits of <i>Pandanus penetrans</i> leaves	38
3.4	Similarity of fungal taxa composition between <i>Dracaena loureiri</i> and <i>Pandanus</i> species	39
3.5	Fungi occurrence on different tissues of Pandanus odoratissimus.	42
3.6	Frequency and overall percentage occurrence of fungal taxa on Pandanus penetrans leaves during the succession process.	46
3.7	A comparison of the pathogenic taxa recovered from <i>Dracaena loureiri</i> , <i>Pandanus amaryllifolius</i> and <i>P. penetrans</i> .	51
3.8	Results of pathogenicity test of fungal taxa isolated as endophytes, saprobes or pathogens.	60

xvii

LIST OF TABLES (CONTINUED)

ble	Page
Synopsis of ascospore and appendages of <i>Linocarpon</i> spp. for comparison with new taxa (Thongkantha <i>et al.</i> , 2003)	118
Species of Ophioceras found on various habitats	126
A comparision of <i>Ophioceras chiangdaoensis</i> with some other previous described species	128
Comparisons morphological characters of <i>Pseudohalonectria</i> species (Promputtha <i>et al.</i> , 2004)	147
Members of the Magnaporthaceae and their anamorphs	155
Magnaporthaceous taxa and their GenBank accession number used in this study	158
Results of the phylogenetic analysis of 18S and 28S rDNA	162
Magnaporthaceae species member accepted by using morphological taxonomy and molecular taxonomy base on 18S, 28S and ITS rDNA sequences analyses	174
	Synopsis of ascospore and appendages of <i>Linocarpon</i> spp. for comparison with new taxa (Thongkantha <i>et al.</i> , 2003) Species of <i>Ophioceras</i> found on various habitats A comparision of <i>Ophioceras chiangdaoensis</i> with some other previous described species Comparisons morphological characters of <i>Pseudohalonectria</i> species (Promputtha <i>et al.</i> , 2004) Members of the Magnaporthaceae and their anamorphs Magnaporthaceous taxa and their GenBank accession number used in this study Results of the phylogenetic analysis of 18S and 28S rDNA Magnaporthaceae species member accepted by using morphological taxonomy and molecular taxonomy base on 18S, 28S and ITS rDNA

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright[©] by Chiang Mai University All rights reserved

xviii

LIST OF ILLUSTRATIONS

Fig	ure	Page
1.1	Plants selected for the present study	3
1.2	Schematic presentation of the relationships between chapters of the thesis	4
2.1	General structure and use of Pandanus	8
2.2	The different steps in PCR	24
2.3	The exponential amplification of the gene in PCR	24
2.4	Verification of the PCR product on gel	25
2.5	Ribosomal RNA gene (rDNA) diagrams	26
3.1	Isolation and pathogenicity test of fungal pathogens on	30
	Dracaena loureiri and Pandanus spp. leaves	
3.2	Species area curve for fungi collected on Dracaena loureiri	31
	and Pandanus spp. at each sampling times	
3.3	Three-dimensional correspondence analysis of fungal communities	39
	recorded from Dracaena lourieri, Pandanus amaryllifolius, P.	
	odoratissimus and P. penetrans	
3.4	Three-dimensional correspondence ordinations of fungal communities	40
	occurred on Pandanus penetrans leaves that were collected from	
	Chiang Mai and Phayao during 3 seasons	
3.5	Two-dimensional correspondence analysis of fungal communities	41
	occurred on leaves, prop roots and seeds of Pandanus odoratissimus	
	in Rayong Province during cool dry season.	
3.6	Three dimensional correspondence ordinations of fungal communities	43
	occurred on Pandanus penetrans leaves from Suthep Pui National Park	
	that were collected during 3 season of cool dry, hot dry and hot wet as	
	naturally samples, and samples collected during succession stage of	
	decayed at day 0, month 12 and month 18.	
3.7	Species area curve for fungi collected on Pandanus penetrans leaves	45
	at different stages of succession.	

Figu	re	Page
3.8	Three-dimensional correspondence analysis of fungal samples	45
	of Pandanus penetrans at different stages of succession	
3.9	Shannon indices for Pandanus penetrans leaves throughout	49
	the experiment	
3.10	Number of fungal taxa occurring on samples during different stages of decayed	49
3.11	Frequency of dominant fungi occurring on leaves at each sampling time	50
3.12	Appearance of disease symptom on <i>Pandanus penetrans</i> leaves, colony, mycelia, conidiophores and conidia of <i>Acremonium</i> sp.	52
3.13	Appearance of disease symptom on the host leaves: <i>Dracaena loureiri</i> , <i>Pandanus amaryllifolius</i> and <i>P. penetrans</i> . Conidia ooze, conidiophores and conidia of <i>Colletotrichum gloeosporioides</i> . Ascomata, asci and ascospores of <i>Glomerella cingulata</i>	53
3.14	Appearance of disease symptom on the host leaves, conidiophores and conidia of Aspergillus sp.	54
3.15	Appearance of disease symptom on leaf of <i>Dracaena loureiri</i> . Coniophores and conidia of <i>Cladosporium oxysporum</i>	55
3.16	Appearance of disease symptoms on Pandanus penetrans leaf, colony, conidiophores and conidia of Curvularia lunata	56
3.17	Appearance of disease symptoms on the host leaves (Pandanus amaryllifolius and P. penetrans), conidia, chlamydospore, colony, conidiophore and of Fusarium oxysporum	57 Sity
3.18	Appearance of disease symptoms on the host leaves (<i>Dracaena loureiri Pandanus amaryllifolius</i> and <i>P. penetrans</i>) asci, ascospores and colony of. <i>Guignardia</i> sp.	58
3.19	Appearance on <i>Pandanus penetrans</i> leaf, colony, asci and ascospores of <i>Oxydothis</i> sp.	59

Figu	ire	Page
3.20	Appearance of disease symptom on <i>Pandanus penetrans</i> leaf, colonies on culture, conidiophores, conidia and chlamydospores of <i>Penicillium</i> sp.	60
3.21	Appearance of disease symptom on <i>Pandanus amaryllifolius</i> leaf. <i>Colletotrichum gloeosporioides</i> : Isolated culture; No disease symptom on tested leaves that without wound; Disease symptom on wounds of tested leaf; Reisolation of fungus from diseased lesions.	61
3.22	1. Appearance of disease symptom on <i>Pandanus penetrans</i> leaf. 2. Isolated culture of <i>Curvularia lunata</i> . 3. Disease symptom on wounds of tested leaf. 4. Reisolation of fungus from diseased lesions.	62
3.23	1. Appearance of disease symptom on <i>Pandanus amaryllifolius</i> leaf. 2. Isolated culture of <i>Guignardia</i> sp. 3. No disease symptom on tested leaves that without wounding. 4. Disease symptom on wounds of tested leaf. 5. Reisolation of fungi from diseased lesions.	62
Desc	riptions and photographic figures of anamorphic fungi	
4.1	Canalisporium	72
4.2	Dictyochaeta	74
4.3	Ellisembia	76
4.4	Exerohilum	77
4.5	Memnoniella	79
4.6	Myrothecium	82
4.7	Diagrammatic representation of conidiophores, conidiogenous cells and conidia of <i>Myrothecium cintum</i> and <i>M. pandanicola</i>	82

xxi

Figu	re	Page
4.8	Paecilomyces	84
4.9	Periconia	87
4.10	Periconiella	89
4.11	Phaeostalagmus	90
4.12	Phialocephala	92
4.13	Sporidesmium	94
4.14	Stachybotrys	97
4.15	Trichothecium	99
4.16	Veronaea	100
4.17	Zygosporium	106
Desc	riptions and photographic figures of Ascomycetes	
4.18	Astrosphaeriella (Melanommataceae: Dothideales)	108
4.19	Chaetosphaeria (Chaetomiaceae: Sordariales)	110
4.20	Emericella (Trichocomaceae: Eurotiales)	112
4.21	Linocarpon lammiae (Hyponectriaceae: Incertae Sedis)	114
4.22	Linocarpon siamensis Thongkantha, sp. nov.	117
4.23	Linocarpon suthepensis Thongkantha, sp. nov.	117
4.24	Melanochaeta (Chaetosphaeriaceae: Sordariales)	121
4.25	Microthyrium sp. 1 (Microthyriaceae: Dothideales)	123
4.26	Microthyrium sp. 2	124
4.27	Ophioceras (Magnaporthaceae: Incertae Sedis)	129
4.28	Some species of Ophioceras described by Conway and Barr (1977),	130
	Shearer et al. (1999), Teng (1934) and Tsui et al. (2001)	
4.29	Ornatispora (Incertae Sedis, Incertae Sedis)	132
4.30	Oxydothis (Incertae Sedis, Incertae Sedis)	136
4.31	Paraphaeosphaeria (Pleosporales, Phaeosphaeriaceae)	139

xxii

Figu	Figure	
4.32	Phaeonectriella (Halosphaeriales, Halosphaeriaceae)	142
4.33	Phomatospora (Xylariales, Incertae Sedis)	144
4.34	Pseudohalonectria (Incertae Sedis, Magnaporthaceae)	147
4.35	All species of <i>Pseudohalonectria</i> previously described by Cai et al.	148
	(2002), Hyde et al. (1999c), Ono and Kobayashi (2001),	
	Promputta et al. (2004a), and Shearer (1989).	
4.36	Sordaria (Sordariales, Sordariaceae)	150
4.37	Tubeufia (Pleosporales, Tubeufiaceae)	152
5.1	Primers for amplification and sequencing partial 28S rDNA	157
5.2	Primers for amplification and sequencing partial 18S rDNA	157
5.3	One of the best bayesian trees resulting from analysis of alignment I	163
	(28S rDNA-73 taxa) for the major order of the Ascomycetes. The	
	numbers above the branches indicate MP1 bootstrap support proportions	
	from 1000 replicates and numbers below the branches indicate pooled	
	posterior probabilities	
5.4	One of the best bayesian trees resulting from analysis of alignment III	164
	(18S rDNA-65 taxa). The numbers above the branches indicate MP1	
	bootstrap support proportions from 1000 replicates and numbers below	
	the branches indicate pooled posterior probabilities	
5.5	One of the best bayesian trees resulting from analysis of alignment IV	165
	(combine 18S and 28S rDNA-51 taxa). The numbers above the branches	
	indicate MP1 bootstrap support proportions from 1000 replicates and	
	numbers below the branches indicate pooled posterior probabilities	

xxiii

ABBREVIATIONS AND SYMBOLS

gram g L liter

milligram mg minute min milliliter ml

molecular weight MW

nanogram ng No. number

PDA potato dextrose agar rotation per minute rpm

second sec UV ultraviolet

weight by volume w/v

water agar WA

degree Celsius °C

alpha α β beta

μm

microgram μg microlitre μl micrometer

าว**ายาลัยเชีย**อใหม[่]