

## TABLE OF CONTENTS

| Title   | Page |
|---|------|
| ACKNOWLEDGMENTS                                   | iii  |
| ABSTRACT (in English)                             | iv   |
| ABSTRACT (in Thai)                                | v    |
| TABLE OF CONTENTS                                 | vi   |
| LIST OF TABLES                                    | xi   |
| LIST OF ILLUSTRATIONS                             | xiv  |
| ABBREVIATIONS AND SYMBOLS                         | xix  |
| <b>CHAPTER 1 INTRODUCTION</b>                     |      |
| 1.1 Biodiesel fuel                                | 2    |
| 1.2 Methanolysis                                  | 2    |
| 1.2.1 Acid- or alkali-catalyzed alcoholysis       | 3    |
| 1.2.2 Enzymatic alcoholysis by lipase             | 4    |
| 1.2.2.1 Alcoholysis using extracellular lipase    | 5    |
| 1.2.2.2 Methanolysis by intracellular lipase      | 8    |
| 1.2.2.3 Enhancing lipase activity in methanolysis | 9    |
| 1.3 <i>Carica papaya</i> L.                       | 13   |
| 1.3.1 <i>C. papaya</i> latex                      | 14   |
| 1.3.2 Collection of latex                         | 14   |
| 1.3.3 Lipase from <i>C. papaya</i> latex          | 15   |
| 1.3.3.1 Preparation of <i>C. papaya</i> lipase    | 16   |
| 1.3.3.2 Specificity of enzyme                     | 17   |
| 1.4 Objectives                                    | 23   |
| <b>CHAPTER 2 MATERIALS AND METHODS</b>            |      |
| 2.1 Materials                                     | 24   |
| 2.1.1 Chemicals                                   | 24   |



|  | <b>Page</b> |
|--|-------------|
| <b>CHAPTER 3 RESULTS</b>   |             |
| 3.1 Separation of lipase from papaya latex                                     | 36          |
| 3.1.1 Hydrolase activity of CPL  | 37          |
| 3.1.1.1 Hydrolase activity on tributyrin                                       | 37          |
| 3.1.1.2 Hydrolase activity on olive oil  | 37          |
| 3.1.2 Recovery yield of CPL  | 39          |
| 3.1.3 Optimum temperature and optimum pH of CPL                                | 40          |
| 3.2 Optimum enzyme amount for methanolysis of triolein                         | 42          |
| 3.3 CPL specificity for methanolysis and ethanolysis                           | 42          |
| 3.3.1 Methanolysis and ethanolysis of tripalmitin                              | 44          |
| 3.3.2 Methanolysis and ethanolysis of tristearin                               | 44          |
| 3.3.3 Methanolysis and ethanolysis of triolein                                 | 45          |
| 3.4 Improvement of CPL activity in methanolysis of triolein by alcohol washing | 46          |
| 3.5 Specificity of iPCPL on alcoholysis of TAG                                 | 48          |
| 3.5.1 Methanolysis and ethanolysis of trimyristin                              | 48          |
| 3.5.2 Methanolysis and ethanolysis of tripalmitin                              | 49          |
| 3.5.3 Methanolysis and ethanolysis of triolein                                 | 50          |
| 3.5.4 Alcoholysis of tristearin  | 51          |
| 3.6 Methanolysis of triolein catalyzed by iPCPL                                | 52          |
| 3.6.1 Effect of temperature  | 52          |
| 3.6.2 Effect of lipase quantity  | 54          |
| 3.6.3 Effect of substrate ratio  | 55          |
| 3.6.4 Effect of solid buffer (acid and their Na salts)                         | 57          |
| 3.6.5 Effect of water  | 59          |
| 3.6.5.1 Water added to enzyme  | 59          |
| 3.6.5.2 Water added to reaction  | 61          |
| 3.6.6 Effect of organic solvent  | 63          |
| 3.7 Biodiesel fuel from methanolysis of palm oil                               | 64          |
| 3.7.1 Fatty acid composition of palm oil                                       | 64          |

|   | <b>Page</b> |
|---|-------------|
| 3.7.2 Methyl esters from methanolysis of palm oil | 69          |

## **CHAPTER 4 DISCUSSION AND CONCLUSIONS**

|  |    |
|--|----|
| 4.1 Production of CPL from whole latex                         | 72 |
| 4.2 Specificity of CPL on methanolysis and ethanolysis of TAGs | 73 |
| 4.3 Improvement of CPL efficiency on methanolysis              | 75 |
| 4.4 Specificity of iPCPL on alcoholysis of TAGs                | 76 |
| 4.5 Optimum condition of methyl ester synthesis from triolein  | 77 |
| 4.6 Biodiesel fuel synthesis from palm oil                     | 80 |
| 4.7 Conclusions  | 81 |

## **REFERENCES**

82

## **APPENDIX**

93

**Appendix A:** Determination of lipase activity by titrimetric method 93

**Appendix B:** Determination of lipase activity by colorimetry 94

**Appendix C:** Retention time of substrates and products in methanolysis  
of triolein or palm oil 95

**Appendix D:** Standard curves of methyl esters for determination of  
product contents 96

**Appendix E:** GLC chromatograms of reaction mixture from alcoholysis  
of trimyristin 97

**Appendix F:** GLC chromatograms of reaction mixture from alcoholysis  
of tripalmitin 98

**Appendix G:** GLC chromatograms of reaction mixture from  
alcoholysis of tristearin 99

**Appendix H:** GLC chromatograms of reaction mixture from  
alcoholysis of triolein 100

|   | <b>Page</b> |
|---|-------------|
| <b>Appendix I:</b> Standard mass spectrum of methyl esters from analysis of palm oil composition          | 101         |
| <b>Appendix J:</b> Standard curves of ethyl esters and butyl esters for determination of product contents | 102         |
| <b>Appendix K:</b> Standard curves of fatty acids for determination of product contents                   | 103         |
| <b>Appendix L:</b> Standard curves of diacylglycerols for determination of product contents               | 104         |
| <b>Appendix M:</b> Standard curves of monoacylglycerol for determination of product contents              | 105         |
| <b>CURRICULUM VITAE</b>   | 106         |

## LIST OF TABLES

| Table   | Page |
|---|------|
| <b>1.1</b> Comparison between alkali-catalysis and lipase-catalysis processes for biodiesel fuel production.  | 5    |
| <b>1.2</b> Enzymatic alcoholysis reaction using various types of alcohols and lipases.  | 7    |
| <b>1.3</b> Effective methanolysis processes with extracellular lipases.   | 8    |
| <b>1.4</b> Effect of pretreatment on methanolysis of soybean oil by Novozyme 435 under different immersion conditions.  | 12   |
| <b>1.5</b> Composition of <i>C. papaya</i> parts per 100 g.   | 13   |
| <b>1.6</b> Lipolytic activity of crude CPL preparation from various plant varieties in comparison with commercially available crude or purified preparation of papain.  | 16   |
| <b>1.7</b> Hydrolysis activity on tributyroylglycerol of lipase from <i>C. papaya</i> latex.  | 17   |
| <b>1.8</b> Time course of the formations of new TAGs from the interesterification of 1-butyroyl-2-stearoyl-3-palmitoyl- <i>sn</i> -glycerol ( <i>sn</i> -BSP) and trimyristin (MMM). TAGs formed by acyl exchange at the <i>sn</i> 3 position of <i>sn</i> -BSP (BSM+MMP) and the <i>sn</i> 1 position of <i>sn</i> -BSP (BMM+MSP). | 23   |
| <b>3.1</b> Weight of lyophilized lipase obtained from papaya latex.   | 37   |
| <b>3.2</b> Hydrolysis activity on tributyrin of CPL which was determined by titrimetric method.   | 38   |
| <b>3.3</b> Hydrolysis activity of CPL on olive oil by titrimetric and colorimetric methods.   | 39   |
| <b>3.4</b> Recovery yield of CPL obtained from papaya latex.  | 39   |
| <b>3.5</b> Hydrolysis activity of CPL at pH 7 on olive oil at different temperatures.   | 40   |
| <b>3.6</b> Hydrolysis activity of CPL on olive oil in different buffer solutions at 55°C.   | 41   |
| <b>3.7</b> Effect of enzyme amounts on methanolysis of triolein at 37°C for 24 h.   | 43   |

| <b>Table</b>   | <b>Page</b> |
|--|-------------|
| <b>3.8</b> Product contents from methanolysis and ethanolysis of tripalmitin catalyzed by CPL.                                       | 44          |
| <b>3.9</b> Product contents from methanolysis and ethanolysis of tristearin catalyzed by CPL.  | 45          |
| <b>3.10</b> Product contents from methanolysis and ethanolysis of triolein catalyzed by CPL.   | 46          |
| <b>3.11</b> Product contents from methanolysis of triolein catalyzed by native and alcohol washed CPLs.                              | 47          |
| <b>3.12</b> Product contents from alcoholysis of trimyristin catalyzed by iPCPL.   | 48          |
| <b>3.13</b> Product contents from alcoholysis of tripalmitin catalyzed by iPCPL.   | 49          |
| <b>3.14</b> Product contents from alcoholysis of triolein catalyzed by iPCPL.  | 50          |
| <b>3.15</b> Product contents from alcoholysis of tristearin catalyzed by iPCPL.  | 51          |
| <b>3.16</b> The formation of products from methanolysis of triolein catalyzed by iPCPL at different temperatures.                    | 53          |
| <b>3.17</b> Percentage of ME content from methanolysis of triolein catalyzed by different quantities of iPCPL.                       | 54          |
| <b>3.18</b> Product contents from methanolysis of triolein catalyzed by iPCPL at different concentrations of methanol.               | 56          |
| <b>3.19</b> Contents of ME from methanolysis of triolein catalyzed by iPCPL which contained various buffer solutions.                | 58          |
| <b>3.20</b> Product contents from methanolysis of triolein catalyzed by iPCPL in the presence of different solid buffers.            | 58          |
| <b>3.21</b> Effect of water added to iPCPL on product contents from methanolysis of triolein.  | 60          |
| <b>3.22</b> Effect of water adding into substrate mixtures on the product contents from methanolysis of triolein catalyzed by iPCPL. | 62          |
| <b>3.23</b> Product contents from methanolysis of triolein catalyzed by iPCPL in several organic solvents.                           | 63          |
| <b>3.24</b> The identification of fatty acid methyl esters obtained from palm oil.   | 67          |

| <b>Table</b>              |  | <b>Page</b> |
|---------------------------|--|-------------|
| <b>3.25</b>               | Composition of fatty acids in palm oil.  | 68          |
| <b>3.26</b>               | Composition of products from methanolysis of palm oil catalyzed by iPCPL.  | 70          |
| <b>4.1</b>                | Production yield and some properties of lyophilized CPL from papaya latex.                                       | 73          |
| <b>4.2</b>                | ME contents obtained from methanolysis of triolein catalyzed by native and alcohol washed CPLs.                  | 75          |
| <b>4.3</b>                | Condition and maximum ME content obtained from methanolysis of triolein by iPCPL.                                | 78          |
| <b>Tables in Appendix</b> |  | <b>Page</b> |
| <b>C1</b>                 | Retention time of tri-, di- and monoacylglycerols, esters and free fatty acids from chromatogram analyzed by GC. | 95          |

## LIST OF ILLUSTRATIONS

| Figure  | Page |
|---|------|
| 1.1 General equation of alcoholysis. R <sub>1</sub> , R <sub>2</sub> and R <sub>3</sub> represent alkyl groups.   | 3    |
| 1.2 General equation of methanolysis of TAG. R <sub>1</sub> , R <sub>2</sub> and R <sub>3</sub> represent alkyl groups.   | 3    |
| 1.3 Flow diagrams comparing biodiesel production using (a) alkali-catalysis and (b) lipase-catalysis processes.   | 6    |
| 1.4 Comparison of microbial lipase production processes for methanolysis with (a) extracellular and (b) intracellular lipases.  | 9    |
| 1.5 Time course of stepwise methanolysis of vegetable oil by immobilized <i>C. antarctica</i> ; (●) three-step addition and (○) two-step addition.  | 11   |
| 1.6 The effect of methanol concentrations on the initial reaction rate of methanolysis by pretreated Novozyme 435 with (■) <i>i</i> -propanol, (□) soybean oil, (▲) 2-butanol and (◆) <i>t</i> -butanol.                                  | 12   |
| 1.7 Tapping of papaya fruit using a latex collecting equipment.   | 15   |
| 1.8 Hydrolysis of TAGs by CPL.  | 18   |
| 1.9 TAG hydrolase activity of CPL, measured at pH 8.0, with numerous TAGs.  | 19   |
| 1.10 Specificity constant in the esterification of mixtures of myristic acid and individual unsaturated, hydroxy, epoxy and cyclopentenyl fatty acids with 1-butanol in hexane using CPL as a biocatalyst.                                | 20   |
| 1.11 Formation rate of alkyl octanoates relative to formation rate of 1-hexyl octanoate during esterification of octanoic acid, catalyzed by CPL, with equimolar mixtures of various alcohols containing 1-hexanol as reference standard. | 21   |
| 1.12 GLC identification of TAGs in the CPL-catalyzed interesterification of <i>sn</i> -BSP and MMM based on their equivalent carbon number (ECN).   | 22   |

| <b>Figure</b>   | <b>Page</b> |
|---|-------------|
| <b>3.1</b> CPL was separated from whole latex by centrifugation and dried by lyophilization.  | 36          |
| <b>3.2</b> Effect of temperature on hydrolysis of olive oil by CPL at pH 7.   | 41          |
| <b>3.3</b> Effect of pH on hydrolysis of olive oil catalyzed by CPL at 55°C.  | 42          |
| <b>3.4</b> Effect of enzyme amounts on methanolysis of triolein catalyzed by CPL.   | 43          |
| <b>3.5</b> Contents of (■) esters, (▣) palmitic acid and (▤) dipalmitoylglycerol from methanolysis and ethanolysis of tripalmitin catalyzed by CPL.   | 44          |
| <b>3.6</b> Contents of (■) esters, (▣) stearic acid and (▤) distearoylglycerol from methanolysis and ethanolysis of tristearin catalyzed by CPL.  | 45          |
| <b>3.7</b> Contents of (■) esters and (▤) dioleoylglycerol resulted from methanolysis and ethanolysis of triolein catalyzed by CPL.   | 46          |
| <b>3.8</b> Contents of (■) methyl ester, (▣) diacylglycerol and (▤) monoacylglycerol which produced from the methanolysis of triolein catalyzed by native CPL and CPL washing with <i>i</i> -propanol (iP), <i>n</i> -propanol (nP), <i>n</i> -butanol (nB) and <i>t</i> -butanol (tB). | 47          |
| <b>3.9</b> Contents of (■) esters, (▣) dimyristoylglycerol, (▤) monomyristoylglycerol from both reaction types catalyzed by iPCPL.  | 49          |
| <b>3.10</b> Contents of (■) esters, (▣) dipalmitoylglycerol, (▤) monopalmitoylglycerol from both reaction types catalyzed by iPCPL.   | 50          |
| <b>3.11</b> Contents of (■) esters, (▣) dioleoylglycerol, (▤) monooleoylglycerol from both reaction types catalyzed by iPCPL.   | 51          |
| <b>3.12</b> Contents of (■) esters, (▣) distearoylglycerol, (▤) monostearoylglycerol from methanolysis, ethanolysis and butanolysis of tristearin catalyzed by iPCPL.   | 52          |

| <b>Figure</b>  | <b>Page</b> |
|--|-------------|
| <b>3.13</b> Contents of ME from methanolysis of triolein catalyzed by iPCPL at temperatures of (○) 25, (△) 30, (□) 37, (●) 40, (▲) 45 and (■) 50°C.  | 54          |
| <b>3.14</b> Contents of ME obtained from methanolysis of triolein catalyzed by different amounts of iPCPL.   | 55          |
| <b>3.15</b> Influence of methanol concentration on methanolysis of triolein catalyzed by iPCPL using molar ratios of triolein:methanol at (○) 1:1, (△) 1:2, (□) 1:3, (●) 1:4, (▲) 1:5 and (■) 1:6. | 57          |
| <b>3.16</b> Effect of solid buffers on production of ME from methanolysis of triolein catalyzed by iPCPL.  | 59          |
| <b>3.17</b> Effect of water at (△) 5%, (□) 10%, (▲) 15% and (■) 20% added to iPCPL on methanolysis of triolein compared with the (○) control.  | 61          |
| <b>3.18</b> Effect of water (△) 5%, (□) 10%, (▲) 15% and (■) 20% added to reaction mixture on methanolysis of triolein compared with the (○) control.  | 63          |
| <b>3.19</b> Influence of organic solvents on production of (▣) methyl esters, (▤) mono- and (▥) diacylglycerols from methanolysis of triolein catalyzed by iPCPL.                                  | 64          |
| <b>3.20</b> GC-MS chromatogram of reaction mixture from methanolysis of palm oil catalyzed by BF <sub>3</sub> .  | 65          |
| <b>3.21a</b> Mass spectrum of peak a in <b>Figure 3.20</b> which was corresponded to methyl myristate.   | 65          |
| <b>3.21b</b> Mass spectrum of peak b in <b>Figure 3.20</b> which was corresponded to methyl palmitate.   | 66          |
| <b>3.21c</b> Mass spectrum of peak c in <b>Figure 3.20</b> which was corresponded to methyl linoleate.   | 66          |
| <b>3.21d</b> Mass spectrum of peak d in <b>Figure 3.20</b> which was corresponded to methyl oleate.  | 67          |
| <b>3.21e</b> Mass spectrum of peak e in <b>Figure 3.20</b> which was corresponded to methyl stearate.  | 67          |

| <b>Figure</b>   | <b>Page</b> |
|---|-------------|
| <b>3.22</b> GLC chromatogram of reaction mixture from methanolysis of palm oil catalyzed by iPCPL using palm oil:methanol at molar ratio of 1:4 at 24 h of reaction.  | 71          |
| <b>3.23</b> Influence of methanol concentration on methanolysis of palm oil. Reaction was performed at palm oil:methanol molar ratio of (O) 1:1, ( $\Delta$ ) 1:2, ( $\square$ ) 1:3, ( $\bullet$ ) 1:4 and ( $\blacktriangle$ ) 1:5. | 71          |
| <b>4.1</b> Contents of ( $\boxtimes$ ) methyl esters and ( $\boxplus$ ) ethyl esters obtained from methanolysis and ethanolysis, respectively, of tripalmitin (TP), tristearin (TS) and triolein (TO).                                | 74          |
| <b>4.2</b> Contents of ( $\boxtimes$ ) methyl esters, ( $\boxplus$ ) ethyl esters and ( $\square$ ) butyl ester obtained from alcoholysis of the corresponding TAGs in chloroform at 37°C.  | 76          |
| <br><b>Figures in Appendix</b>  |             |
| <b>B1</b> Standard curve of oleic acid.   | 94          |
| <b>D1</b> Standard curves of (a) methyl myristate, (b) methyl palmitate, (c) methyl linoleate, (d) methyl oleate and (e) methyl stearate.   | 96          |
| <b>E1</b> Chromatograms of (a) methanolysis and (b) ethanolysis of trimyristin.   | 97          |
| <b>F1</b> Chromatograms of (a) methanolysis and (b) ethanolysis of tripalmitin.   | 98          |
| <b>G1</b> Chromatograms of (a) methanolysis and (b) ethanolysis of tristearin.  | 99          |
| <b>H1</b> Chromatograms of (a) methanolysis and (b) ethanolysis of triolein.  | 100         |
| <b>I1</b> MS spectrum of (a) methyl myristate, (b) methyl palmitate, (c) methyl linoleate, (d) methyl oleate and (e) methyl stearate.   | 101         |

| <b>Figures in Appendix</b>  | <b>Page</b> |
|---|-------------|
| <b>J1</b> Standard curves of (a) ethyl myristate, (b) ethyl palmitate, (c) ethyl oleate, (d) ethyl stearate and (e) butyl stearate.     | 102         |
| <b>K1</b> Standard curves of (a) palmitic acid, (b) oleic acid and (c) stearic acid.  | 103         |
| <b>L1</b> Standard curves of (a) dimyristoylglycerol, (b) ipalmitoylglycerol, (c) dioleoylglycerol and (d) distearoylglycerol.          | 104         |
| <b>M1</b> Standard curves of (a) monomyristoylglycerol, (b) monopalmitoylglycerol, (c) monooleoylglycerol and (d) monostearoylglycerol. | 105         |

