

TABLE OF CONTENTS

	Page
Acknowledgement	iii
Abstract (in English)	v
Abstract (in Thai)	viii
List of tables	xvi
List of illustrations	xxii
Abbreviations	xxv
Chapter 1 Introduction	
1.1 Statement and significance of the problem	1
1.2 Objectives	2
1.3 Scope of study	2
1.4 Literature review	4
1.4.1 Plants	4
1.4.1.1 Guttiferae	4
1.4.1.1.1 Hypericum	4
1.4.1.1.2 Calophyllum	10
1.4.1.1.3 Cratoxylum	11
1.4.1.1.4 Garcinia	11
1.4.1.2 Schisandraceae	13
1.4.1.3 Plant under studied	16
1.4.2 Biological activity assay	23

1.4.2.1	DPPH free radical scavenging activity	23
1.4.2.2	Sulforhodamine B assay	27
1.4.2.3	Lymphocyte proliferation assay	31
Chapter 2 Materials and methods		
2.1	Materials and equipments	34
2.1.1	Plants	34
2.1.2	Chemicals / tumor cell lines	34
2.1.3	Equipments	36
2.2	Methods	37
Part 1:	Collection of the plants	39
Part 2:	Preparation of the crude extracts and bioactivities screening test	44
2.2.1	Preparation of the crude extracts	44
2.2.2	Free radical scavenging activity study of crude extract (DPPH assay)	46
2.2.3	An antitumor activity study of crude extracts (SRB assay)	46
Part 3:	Extraction, isolation, purification and structure elucidation of the selected plants	47
3.1	Extraction, isolation purification and structure elucidation of <i>Hypericum hookerianum</i>	47
3.2	Extraction, isolation, purification and structure elucidation of <i>Schisandra verruculosa</i>	48

Part 4: Bioactivities of the isolated compounds	49
4.1 Tumor cell growth assay	49
4.2 Human lymphocytes proliferation assay	50
4.3 Cell viability determination by trypan blue exclusion assay	50
4.4 Free radical scavenging activity	50
Chapter 3 Results and discussion	
Part 1: Collection of the plants	51
Part 2: Preparation of the crude extracts and bioactivities screening test	52
2.1 Preparation of the crude extracts	52
2.2 Free radical scavenging activity study of crude extracts (DPPH assay)	53
2.3 An antitumor activity study of crude extracts (SRB assay)	58
Part 3: Extraction, isolation, purification and structure elucidation of the selected plants	62
3.1 Extraction, isolation, purification and structure elucidation of <i>Hypericum hookerianum</i>	66
3.1.1 Preparation of the extract	66
3.1.2 Fractionation of the extract	66
3.1.3 Isolation and Purification of the pure compounds	68
3.1.4 Structure elucidation	71
3.2 Extraction, isolation, purification and structure elucidation of <i>Schisandra verruculosa</i>	91
3.2.1 Preparation of the extract	91

3.2.2 Chlorophyll elimination of the chloroform fraction of the methanol extract	91
3.2.3 Fractionation of the extract	91
3.2.4 Isolation and purification of the constituents	92
3.2.5 Fractionation of the crude extract	93
3.2.6 Isolation and purification of the constituents	94
3.2.7 Structure elucidation	98
Part 4: Bioactivities of the isolated compounds	115
4.1 Tumor cell growth assay	115
4.2 Human lymphocytes proliferation assay	118
4.3 Free radical scavenging activity	120
Chapter 4 Conclusion	122
References	125
Appendix	
Appendix A	138
A.1 Calculation of the percentage yield of the methanol and chloroform fraction of the methanol crude extracts	138
A.2 Calculation of 50% free radical scavenging activity (IC_{50}) of the methanol and chloroform fraction of the methanol extracts	142
Appendix B	145
B.1 Optimum cell density of HeLa, KB and B16F10 cell line in SRB assay	145

B.2 Calculation of the 50% growth inhibition (GI_{50}) of the crude extracts and positive control on cancer cell lines	147
Appendix C	151
C.1 Spectrum of the isolated compounds	151
Appendix D	164
D.1 The 50% cell growth inhibition of isolated compounds and positive control on MCF-7, NCI-H460, SF-268 and UACC-62 cancer cell lines	164
D.2 Lymphocyte proliferation assay and cell viability	168
Curriculum vitae	171

LIST OF TABLES

Table	Page
1.1 Summary of the selected Thai plants in family Guttiferae and Schisandraceae	22
1.2 A list of 60 human cell lines and inoculation density used in the screening and maintenance at NCI	29
1.3 List of additional cell lines evaluated for use in the screening of cytotoxicity	30
3.1 The percentage yield of methanol and chloroform fraction of the methanol extracts from various parts of the selected Thai plants in family Guttiferae and Schisandraceae	52
3.2 The IC ₅₀ values of the selected Thai plant extracts	57
3.3 Effect of methanol and chloroform fraction of the methanol extracts from wood of the selected Thai plants in family Guttiferae and Schisandraceae on the growth of human cancer cell lines	59
3.4 Effect of methanol and chloroform fraction of the methanol extracts from leaves of the selected Thai plants in family Guttiferae and Schisandraceae on the growth of human cancer cell lines	60
3.5 ¹ H NMR spectrum (CDCl ₃ , 300 MHz) of 5-hydroxy-2-methoxyxanthone	73
3.6 COSY spectrum (CDCl ₃ , 300 MHz) of 5-hydroxy-2-methoxyxanthone	72
3.7 ¹³ CNMR and HSQC spectrum (CDCl ₃ , 75.47 MHz) of 5-hydroxy-2-methoxyxanthone	73

3.8	HMBC spectrum (CDCl ₃ , 500 MHz) of 5-hydroxy-2-methoxyxanthone	74
3.9	¹ H NMR spectrum (CDCl ₃ , 300 MHz) of 2-hydroxy-3-methoxyxanthone	75
3.10	¹³ CNMR and HSQC spectrum (CDCl ₃ , 75.47 MHz) of 2-hydroxy-3-methoxyxanthone	76
3.11	HMBC spectrum (CDCl ₃ , 500 MHz) of 5-hydroxy-2-methoxyxanthone	77
3.12	¹ H NMR spectrum (CDCl ₃ , 300 MHz) of <i>trans</i> -kielcorin	78
3.13	COSY spectrum (CDCl ₃ , 300 MHz) of <i>trans</i> -kielcorin	79
3.14	¹³ CNMR and HSQC spectrum (CDCl ₃ , 75.47 MHz) of <i>trans</i> -kielcorin	80
3.15	HMBC spectrum (CDCl ₃ , 500 MHz) of <i>trans</i> -kielcorin	81
3.16	NOSY spectrum (CDCl ₃ , 300 MHz) of <i>trans</i> -kielcorin	82
3.17	¹ H NMR spectrum (CDCl ₃ , 300 MHz) of 4-hydroxy-3-methoxyphenyl ferulate	83
3.18	COSY spectrum (CDCl ₃ , 300 MHz) of 4-hydroxy-3-methoxyphenyl ferulate	84
3.19	¹³ CNMR and HSQC spectrum (CDCl ₃ , 75.47 MHz) of 4-hydroxy-3-methoxyphenyl ferulate	84
3.20	HMBC spectrum (CDCl ₃ , 500 MHz) of 4-hydroxy-3-methoxyphenyl ferulate	85
3.21	¹ H NMR spectrum (CDCl ₃ , 300 MHz) of 3β-O-caffeoylbetulinic acid	87
3.22	¹³ CNMR and HSQC spectrum (CDCl ₃ , 75.47 MHz) of 3β-O-caffeoyl betulinic acid	88
3.23	¹ H NMR spectrum (CDCl ₃ , 300 MHz) of vanillic acid	99
3.24	¹³ CNMR and HSQC spectrum (CDCl ₃ , 75.47 MHz) of vanillic acid	100
3.25	HMBC spectrum (CDCl ₃ , 300 MHz) of vanillic acid	100

3.26	^1H NMR spectrum (CDCl_3 , 300 MHz) of abscisic acid	101
3.27	^{13}C NMR and HSQC spectrum (CDCl_3 , 75.47 MHz) of abscisic acid	102
3.28	^1H NMR spectrum (CDCl_3 , 300 MHz) of methyl 4-hydroxybenzoate	103
3.29	^{13}C NMR and HSQC spectrum (CDCl_3 , 75.47 MHz) of methyl 4-hydroxybenzoate	103
3.30	HMBC spectrum (CDCl_3 , 500 MHz) of methyl 4-hydroxybenzoate	104
3.31	^1H NMR spectrum (CDCl_3 , 300 MHz) of 4-hydroxybenzaldehyde	104
3.32	COSY spectrum (CDCl_3 , 300 MHz) of 4-hydroxybenzaldehyde	105
3.33	^{13}C NMR and HSQC spectrum (CDCl_3 , 75.47 MHz) of 4-hydroxybenzaldehyde	105
3.34	HMBC spectrum (CDCl_3 , 500 MHz) of 4-hydroxybenzaldehyde	105
3.35	^1H NMR spectrum (CDCl_3 , 500 MHz) of 3-methyl 3,4-dihydroxybenzoate	106
3.36	COSY spectrum (CDCl_3 , 300 MHz) of 3-methyl 3,4-dihydroxybenzoate	106
3.37	^{13}C NMR and HSQC spectrum (CDCl_3 , 75.47 MHz) of 3-methyl 3,4-dihydroxybenzoate	107
3.38	HMBC spectrum (CDCl_3 , 300 MHz) of 3-Methyl 3,4-dihydroxybenzoate	107
3.39	^1H NMR spectrum (CDCl_3 , 300 MHz) of 1-(4-hydroxy-3-methoxyphenyl)-3-hydroxy-propan-1-one	108
3.40	COSY spectrum (CDCl_3 , 300 MHz) of 1-(4-hydroxy-3-methoxyphenyl)-3-hydroxy-propan-1-one	109
3.41	^{13}C NMR and HSQC spectrum (CDCl_3 , 125.77 MHz) of 1-(4-hydroxy-3-methoxyphenyl)-3-hydroxy-propan-1-one	109
3.42	HMBC spectrum (CDCl_3 , 500 MHz) of 1-(4-hydroxy-3-methoxyphenyl)-	110

	3-hydroxy-propan-1-one	
3.43	¹ H NMR spectrum (CDCl ₃ , 300 MHz) of 1,2-bis(4-hydroxy-3-methoxy phenyl)-3-hydroxy-propan-1-one	111
3.44	COSY spectrum (CDCl ₃ , 300 MHz) of 1,2-bis(4-hydroxy-3-methoxy phenyl)-3-hydroxy-propan-1-one	112
3.45	¹³ CNMR and HSQC spectrum (CDCl ₃ , 125.77 MHz) of 1,2-bis(4-hydroxy-3-methoxyphenyl)-3-hydroxy-propan-1-one	112
3.46	HMBC spectrum (CDCl ₃ , 300 MHz) of 1,2-bis(4-hydroxy-3-methoxy phenyl)-3-hydroxy-propan-1-one	113
3.47	¹ H NMR spectrum (CDCl ₃ , 300 MHz) of 4-hydroxybenzoic acid	114
3.48	¹³ CNMR and HSQC spectrum (CDCl ₃ , 75.47 MHz) of 4-hydroxybenzoic acid	114
3.49	The IC ₅₀ of compounds from <i>Hypericum hookerianum</i> on the growth of human cancer cell lines	116
3.50	The IC ₅₀ of compounds from <i>Shisandra verruculosa</i> on the growth of human cancer cell lines	117
3.51	The IC ₅₀ of compounds from <i>Hypericum hookerianum</i> on proliferation of human lymphocytes assay	118
3.52	The IC ₅₀ of compounds from <i>Shisandra verruculosa</i> on proliferation of human lymphocytes assay	119
3.53	The IC ₅₀ of compounds from <i>Hypericum hookerianum</i> on DPPH free radical scavenging activity assay	120
3.54	The IC ₅₀ of compounds from <i>Shisandra verruculosa</i> on DPPH free radical scavenging activity assay	121

A.1	Calculation of the percentage yields of methanol and chloroform fraction of the methanol wood extract of <i>Hypericum hookerianum</i>	138
A.2	Calculation of the percentage yield of methanol and chloroform fraction of the methanol leaf extract of <i>Garcinia speciosa</i>	138
A.3	Calculation of the percentage yield of methanol and chloroform fraction Of the methanol wood extract of <i>Garcinia speciosa</i>	139
A.4	Calculation of the percentage yield of methanol and chloroform fraction of the methanol leaf and wood extract of <i>Garcinia xanthochymus</i>	139
A.5	Calculation of the percentage yield of methanol and chloroform fraction of methanol fruit extract of <i>Garcinia xanthochymus</i>	140
A.6	Calculation of the percentage yield of methanol and chloroform fraction of the methanol leaf and wood extract of <i>Cratoxylum formosum</i> ssp. <i>pruniflorum</i>	140
A.7	Calculation of the percentage yield of methanol and chloroform fraction of the methanol leaf and wood extract of <i>Calophyllum polyanthum</i>	141
A.8	DPPH radical scavenging activity (%) of the extracts	143
B.1	Inoculation densities of cell lines	145
B.2	The 50% growth inhibition of the methanol and chloroform fraction of the methanol wood extracts on cell lines	148
B.3	The 50% growth inhibition of the methanol and chloroform fraction of the methanol leave extracts on cell lines	149
B.4	The 50% growth inhibition of doxorubicin on cancer cell lines	151
D.1	The GI ₅₀ of the compounds from <i>H. hookerianum</i> on MCF-7 cell line	164
D.2	The GI ₅₀ of the compounds from <i>H. hookerianum</i> on NCI-H460 cell lines	165

D.3	The GI_{50} of the compounds from <i>H. hookerianum</i> on SF-268 cell line	165
D.4	The GI_{50} of the compounds from <i>H. hookerianum</i> on UACC-62 cell lines	166
D.5	The GI_{50} of the compounds from <i>S. verrucolosa</i> on MCF-7 cell lines	166
D.6	The GI_{50} of the compounds from <i>S. verrucolosa</i> on NCI-H460 cell lines	167
D.7	The GI_{50} of the compounds from <i>S. verrucolosa</i> on SF-268 cell lines	167
D.8	The 50% growth inhibition of doxorubicin on cancer cell lines	168
D.9	The IC_{50} of the compounds from <i>H. hookerianum</i> on human lymphocyte	169
D.10	The IC_{50} of the compounds from <i>S. verrucolosa</i> on human lymphocyte	169
D.11	Cell viability by trypan blue exclusion assay	170


LIST OF ILLUSTRATIONS

Figure	Page
1.1 <i>Hypericum hookerianum</i> Wight & Arn	16
1.2 <i>Garcinia speciosa</i> Wall	17
1.3 <i>Garcinia xanthochymus</i> Hook. F. ex T. Anderson	18
1.4 <i>Cratoxylum formosum</i> ssp. <i>pruniflorum</i> (Kurz) Gogel	19
1.5 <i>Calophyllum polyanthum</i> Wall ex Choisy	20
1.6 <i>Schisandra verruculosa</i> Gagnap	21
1.7 Structure of DPPH	24
1.8 Structure of MTT and their corresponding reaction products	32
2.1 Scope of the Part 1 to Part 4 study	37
2.2 The herbarium specimen of <i>Hypericum hookerianum</i>	38
2.3 The herbarium specimen of <i>Garcinia speciosa</i>	39
2.4 The herbarium specimen of <i>Garcinia xanthochymus</i>	40
2.5 The herbarium specimen of <i>Cratoxylum formosum</i> ssp. <i>pruniflorum</i>	41
2.6 The herbarium specimen of <i>Calophyllum polyanthum</i>	42
2.7 The herbarium specimen of <i>Schisandra verruculosa</i>	43
2.8 Diagram of the preparation of crude extract	44
3.1 Comparison of free radical scavenging activity (%) of the methanol wood extracts of the six selected Thai plants with the standard antioxidants	53
3.2 Comparison of free radical scavenging activity of the chloroform fraction of the methanol wood extracts of the six selected Thai plants with the	54

standard antioxidants	
3.3 Comparison of free radical scavenging activity (%) of the methanol leaves extracts of the six selected Thai plants with the standard antioxidants	55
3.4 Comparison of free radical scavenging activity (%) of the chloroform Fraction of the methanol leaves extracts of the six selected Thai plants with the standard antioxidants	56
3.5 Diagram of the isolation and purification of the compounds from <i>Hypericum hookerianum</i>	63
3.6 Diagram of the isolation and purification of the isolated compounds from <i>S. verruculasa</i>	64
3.6 Diagram of the isolation and purification of the isolated compounds from <i>Schisandra verruculasa</i>	65
3.7 The structure of 5-hydroxy-2-methoxyxanthone	72
3.8 Chemical shifts and coupling constant of 5-hydroxy-2-methoxyxanthone	74
3.9 The structure of 2-hydroxy-3-methoxyxanthone	75
3.10 Chemical shifts and coupling constant of 5-hydroxy-2-methoxyxanthone	77
3.11 The structure of <i>trans</i> -kielcorin	78
3.12 Chemical shifts and coupling constant of <i>trans</i> -kielcorin	72
3.13 The structure of 4-hydroxy-3-methoxyphenyl ferulate	83
3.14 Chemical shifts and coupling constant of 4-hydroxy-3-methoxyphenyl ferulate	86
3.15 The structure of 3 β -O-caffeoylbetulinic acid	87
3.16 Chemical shifts and coupling constant of 3 β -O-caffeoylbetulinic acid	90
3.17 The structure of vanillic acid	99

3.18	The structure of abscisic acid	101
3.19	The structure of methyl 4-hydroxybenzoate	103
3.20	The structure of 4-hydroxybenzaldehyde	104
3.21	The structure of 3-methyl 3,4-dihydroxybenzoate	106
3.22	The structure of 1-(4-hydroxy-3-methoxyphenyl)-3-hydroxy-propan-1-one	107
3.23	The structure of 1,2-bis(4-hydroxy-3-methoxyphenyl)-3-hydroxy-propan-1-one	111
3.24	The structure of 4-hydroxybenzoic acid	114
B.1	Optimum cell density of HeLa cell line in SRB assay	145
B.2	Optimum cell density of KB cell line in SRB assay	146
B.3	Optimum cell density of B16F10 cell line in SRB assay	146
C.1	^1H -, ^{13}C -NMR analysis of 5-hydroxy-2-methoxyxanthone	151
C.2	^1H -, ^{13}C -NMR analysis of 2-hydroxy-3-methoxyxanthone	152
C.3	^1H -, ^{13}C -NMR analysis of <i>trans</i> -kielcorin	153
C.4	^1H -, ^{13}C -NMR analysis of 4-hydroxy-3-methoxyphenyl ferulate	154
C.5	^1H -, ^{13}C -NMR analysis of 3β -O-caffeoylbetulinic acid	155
C.6	^1H -, ^{13}C -NMR analysis of vanillic acid	156
C.7	^1H -, ^{13}C -NMR analysis of abscisic acid	157
C.8	^1H -, ^{13}C -NMR analysis of methyl 4-hydroxybenzoate	158
C.9	^1H -, ^{13}C -NMR analysis of 4-hydroxybenzaldehyde	159
C.10	^1H -, ^{13}C -NMR analysis of 3-methyl 3,4-dihydroxybenzoate	160
C.11	^1H -, ^{13}C -NMR analysis of 1-(4-hydroxy-3-methoxyphenyl)-3-hydroxyl-propan-1-one	161

- C.12 ^1H -, ^{13}C -NMR analysis of 1,2-bis- (4-hydroxy-3-methoxyphenyl)-3-hydroxyl-propan-1-one 162
- C.13 ^1H -, ^{13}C -NMR analysis of 4-hydroxybenzoic acid 163



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

ABBREVIATION

^{13}C NMR	Carbon thirteen Nuclear Magnetic Resonance
CDCl_3	Deuterated chloroform
CHCl_3	Chloroform
CNS	Central nervous system
Co-115	Colon carcinoma cell line: Co-115
COLO-205	Colon carcinoma cell line: COLO-205
COSY	Correlated Spectroscopy
<i>d</i>	doublet
<i>dd</i>	double doublet
DMSO	Dimethylsulfoxide
DPPH	1,1-Diphenyl-2-picrylhydrazyl
EBV-EA	Epstein-Barr virus early antigen
EI-MS	Electron Impact Mass Spectrometry
FBS	Fetal bovine serum
FCS	Fetal calf serum
GI_{50}	Concentration which inhibits cell growth by 50%
HBeAg	Hepatitis B antigen
HeLa	Cervical cancer cell line: HeLa
HEPA	Hepatoma cell line: HEPA
HepG2	Human hepatoma cell line: HepG2
Hep-2	Larynx epithelium carcinoma cell line: Hep-2
^1H NMR	Proton Nuclear Magnetic Resonance

HIV	Human Immunodeficiency Virus
HL-60	Human leukemia cell: HL-60
HMBC	Heteronuclear Multiple Bond Correlation
HRMS	High Resolution Mass Spectrometry
HSQC	Heteronuclear Single Quantum Coherence
Hz	Hertz
IC ₅₀	Concentration giving 50% inhibition
<i>J</i>	Coupling constant in Hz
KB	Human oral epidermoid carcinoma cell line: KB
LL/2	Mouse Lewis lung carcinoma cell line
<i>m</i>	multiplet
<i>m/z</i>	mass per charge
MCF-7	Human breast carcinoma cell line: MCF-7
MeOH	Methanol
MHz	Mega hertz
MTT	3-(4,5-Dimethylthiazole-2-yl)-2,5-dimethyltetrazolium bromide
NBT	Nitroblue tetrazolium
NCI	National Cancer Institute
NCI-H460	Non-small cell lung cancer cell line: NCI-H460
NGF	Nerve growth factor
nm	Nanometer
NMR	Nuclear Magnetic Resonance
NOESY	Nuclear Overhauser and Exchange Spectroscopy
OMe	Methoxy

PAF	Platelet-activating factor
PHA	Phytohemagglutinin
P388	Mouse lymphoid neoplasm cell line: P388
ppm	parts per million
s	singlet
SF-268	CNS cancer cell line: SF-268
sp.	Species (singular)
spp.	Species (plural)
SRB	Sulphorodamine B
T	triplet
TCA	Trichloroacetic acid
TPA	12-o-tetradecanoylphorbol-13-acetate
TK-10	Renal cancer cell line: TK-10
TLC	Thin layer chromatography
Tris	Tris (hydroxymethyl)aminomethane
UACC-62	Melanoma cell line: UACC-62
δ	Chemical shift value in ppm
λ	Wavelength in nanometer
$^{\circ}\text{C}$	Celsius degree
μg	Microgram
μl	Microliter
Wehil 64	Mouse fibrosarcoma: Wehil 64