

## TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS.....	iii
ABSTRACT (IN ENGLISH).....	v
ABSTRACT (IN THAI).....	ix
TABLE OF CONTENTS.....	xiii
LIST OF TABLES.....	xviii
LIST OF FIGURES.....	xx
ABBREVIATIONS.....	xxvi
CHAPTER 1 INTRODUCTION.....	1
1. GENERAL INTRODUCTION.....	1
2. LITERATURE REVIEWS.....	3
2.1 <i>An. dirus</i> complex.....	4
2.2 <i>An. maculatus</i> complex.....	7
2.3 <i>An. minimus</i> complex.....	12
2.4 <i>An. sundaicus</i> complex.....	15
2.5 <i>An. uconitus</i> .....	17

## TABLE OF CONTENTS (cont.)

	Page
3. PURPOSES OF THE STUDY.....	18
4. SIGNIFICANT OF THE RESEARCH.....	19
CHAPTER 2 MATERIALS AND METHODS.....	20
1. MATERIALS.....	20
1.1 For metaphase chromosome preparation.....	20
1.2 For light and scanning electron microscopic studies..	20
1.3 For polytene chromosomes preparation.....	20
1.4 For isoenzyme study.....	20
1.5 For molecular study.....	21
2. METHODS.....	21
2.1 Collection of mosquitoes.....	21
2.2 Rearing mosquitoes.....	24
2.3 Metaphase chromosome preparation.....	25
2.4 Morphometric and morphological studies of eggs, larvae, pupae and adults under light and/or scanning electron microscopy.....	26

## TABLE OF CONTENTS (cont.)

	Page
2.5 Ovarian nurse cell polytene chromosome preparation.	27
2.6 Isoenzyme study.....	27
2.7 Molecular study.....	30
2.7.1 DNA extraction, PCR amplification, cloning and sequencing.....	30
2.7.2 Sequence data analysis.....	31
2.8 Hybridization experiments.....	32
2.9 Establishment of mixed colonies of the three karyotypic forms.....	33
2.10 Malarial susceptibility test.....	33
2.10.1 Laboratory-raised <i>An. aconitus</i> Form B and C	33
2.10.2 Outgroup and ingroup control vectors.....	34
2.10.3 <i>P. falciparum</i> and <i>P. vivax</i> gametocytes.....	34
2.10.4 Infection of mosquitoes with <i>P. falciparum</i> and <i>P. vivax</i> gametocytes.....	35
2.11 Statistical analyses.....	35

### TABLE OF CONTENTS (cont.)

	Page
CHAPTER 3 RESULTS.....	36
1. Metaphase karyotype identification.....	36
2. Morphometric and morphological studies of eggs, larvae, pupae and adults under light and/or scanning electron microscopy.....	38
2.1 Morphometric and morphological studies of larvae, pupae and adults under light microscopy.....	38
2.2 Scanning electron microscopy.....	46
2.2.1 Morphometric measurements and counts of float ribs and tubercles.....	46
2.2.2 Eggs topography of three strains of <i>An.</i> <i>aconitus</i> Form B and C.....	48
3. Ovarian nurse cell polytene chromosome investigation.....	53
4. Isoenzyme study.....	55
5. Molecular study.....	60
5.1 Sequences analysis of ITS2, COI and COII regions.....	60
6. Hybridization study.....	64

## TABLE OF CONTENTS (cont.)

	Page
7. Malarial susceptibility test .....	67
7.1 Oocyst rates of <i>An. dirus</i> B, <i>An. minimus</i> A and C, and <i>An. aconitus</i> Form B and C.....	67
7.2 Oocyst and sporozoite rates of <i>An. dirus</i> B, <i>An. minimus</i> A and C, and <i>An. aconitus</i> Form B and C.....	70
CHAPTER 4 DISCUSSION.....	75
CHAPTER 5 CONCLUSION.....	83
REFERENCES.....	85
CURRICULUM VITAE.....	105

## LIST OF TABLES

Table	Page
1. Electrophoretic procedures.....	29
2. Primer used for PCR amplification.....	31
3. Setal branching on 4 <sup>th</sup> instar larvae of <i>An. aconitus</i> Form B and C.	40
4. Setal branching on pupal skins of <i>An. aconitus</i> Form B and C.....	41
5. Comparative adult female characters of <i>An. aconitus</i> Form B and C.....	42
6. Mean distribution of coeloconic sensillae on antennal segments of <i>An. aconitus</i> Form B (n = 38) and C (n = 32).....	43
7. Comparative mean numbers of coeloconic sensillae per antenna, palp ratios and wing dimensions of <i>An. aconitus</i> Form B (n = 38) and C (n= 32).....	44
8. Morphometric measurements and counts of float ribs and tubercles of eggs of <i>An. aconitus</i> Form B (Chiang Mai and Phetchaburi strains) and C (Chiang Mai and Mae Hong Son strains).....	47
9. Allelic frequencies (allele numbers) observed at 10 isoenzymes 16 loci in 4 <sup>th</sup> instar larvae of <i>An. aconitus</i> Form B (Chiang Mai and Phetchaburi strains) and C (Chiang Mai and Mae Hong Son strains).....	56

### LIST OF TABLES (cont.)

Table	Page
10. Allelic frequencies (allele numbers) observed at 11 isoenzymes 13 loci in adult females of <i>An. aconitus</i> Form B (Chiang Mai and Phetchaburi strains) and C (Chiang Mai and Mae Hong Son strains).....	57
11. Geographical origin of mosquitoes and their GenBank accession numbers.....	61
12. Nucleotide polymorphism within and between isoline sequences of <i>An. aconitus</i> Form B and C.....	63
13. Cross mating among isolines of <i>An. aconitus</i> Form B and C.....	65
14. The oocyst rates of <i>An. dirus</i> B, <i>An. minimus</i> A and C and <i>An. aconitus</i> Form B and C after feeding on blood containing gametocytes of <i>P. falciparum</i> and <i>P. vivax</i> , all dissected 8 days after feeding.....	69
15. The oocyst and sporozoite rates of <i>An. dirus</i> B, <i>An. minimus</i> A and C and <i>An. aconitus</i> Form B and C after feeding on blood containing gametocytes of <i>P. falciparum</i> and <i>P. vivax</i> , all dissected 12 days after feeding.....	72

## LIST OF FIGURES

Figures	Page
1. Map of Thailand showing Chiang Mai (CM), Mae Hong Son (MS) and Phetchaburi (PB) provinces, where mosquito collections were performed. Chiang Mai province is situated on latitude 18° 47' N and longitude 98° 59' E in northern Thailand and is approximately 97 and 647 kilometers away from Mae Hong Son province, northwest Thailand and Phetchaburi province, southwest Thailand, respectively. ....	22
2. Summarized experimental design.....	23
3. Metaphase karyotypes of <i>An. aconitus</i> Form B and C (Giemsa staining). Testis chromosomes; Form B: (A) Chiang Mai strain, showing X <sub>1</sub> , Y <sub>2</sub> -chromosomes; (B) Phetchaburi strain, showing X <sub>2</sub> , Y <sub>2</sub> -chromosomes; Form C: (C) Chiang Mai strain, showing X <sub>1</sub> , Y <sub>3</sub> -chromosomes; (D) Mae Hong Son strain, showing X <sub>2</sub> , Y <sub>3</sub> -chromosomes. Ovary chromosomes: (E) showing homozygous X <sub>1</sub> , X <sub>1</sub> -chromosomes, (F) showing homozygous X <sub>2</sub> , X <sub>2</sub> -chromosomes, (G) showing heterozygous X <sub>1</sub> , X <sub>2</sub> -chromosomes. Note, all types of X-chromosomes were found in all forms and strains of <i>An. aconitus</i> .....	37



## LIST OF FIGURES (cont.)

Figures	Page	
4.	Scatter diagram of coeloconic sensilla and palpal ratios in <i>An. aconitus</i> Form B and C.....	45
5.	Whole eggs: Lateral aspect, anterior end (a), posterior end (p) (x 270)....	50
6.	Whole eggs: Ventral aspect, anterior end (a), posterior end (p) (x 270)....	50
7.	Whole eggs: Dorsal aspect, anterior end (a), posterior end (p)(x 270).....	50
8.	Anterior end, showing irregularly jagged tubercles on the deck and three large, rosette-shaped tubercles (x 3,500).....	50
9.	Posterior end, showing irregularly jagged tubercles on the deck and two large, rosette-shape tubercles (x 3,500).....	50
10.	A higher magnification of the large, rosette-shaped tubercle, surrounded by a sclerotized ridge and raised border (x 10,000).....	50
11.	A higher magnification of the irregularly jagged tubercles on the deck (x 12,000).....	51
12.	Irregularly jagged tubercles on the deck and area covered by the float, and outer chorionic tubercles covered with a membrane-like sheet between the frill (fr) and detachment point of the float (dpf). Note, that the cluster of tubercles adjacent to the detachment point of the float more or less form a wavy border (wb) (x 850).....	51

### LIST OF FIGURES (cont.)

Figures	Page
13. A higher magnification of the irregularly jagged tubercles on the area covered by the float, and outer chorionic tubercles covered with a membrane-like sheet between the frill and detachment point of the float (x 1,500).....	51
14. A higher magnification of the irregularly jagged tubercles on the area covered by the float (x 5,000).....	51
15. A higher magnification of the outer chorionic tubercles covered with a membrane-like sheet between the frill and detachment point of the float (x 8,000).....	51
16. Outer chorionic tubercles from the torn membrane-like sheet between the frill and detachment point of the float (x 540).....	51
17. The inner surface of the frill (fr), showing its sclerotized, ridge-like texture with picket-like ribs (x 8,000).....	52
18. The outer surface of the frill (fr), showing its smooth surface and parallel brick-like texture along its entire length (x 8,000).....	52
19. The anterior end, showing the micropylar orifice surrounded by a smooth collar with an irregular outer margin (x 3,700).....	52

## LIST OF FIGURES (cont.)

Figures	Page
20. Outer chorionic tubercles at the anterior third of the egg, showing irregular bases partially covered with a membrane-like sheet (x 3,500).....	52
21. Outer chorionic tubercles in the middle of the egg, showing irregular bases that were almost completely covered with a membrane-like sheet (x 3,500).....	52
22. Outer chorionic tubercles from the torn membrane-like sheet in the middle of the egg, showing irregular bases and flattened-surface (x 7,000).....	52
23. Standard map of ovarian nurse cell polytene chromosomes of <i>An. aconitus</i> Form B (Chiang Mai strain).....	54
24. Isoenzyme pattern of ACPH, ALP, EST, GPD, GPI, HK, IDH, LAP, LDH, ME and PEP of 4 <sup>th</sup> instar larva <i>An. aconitus</i> Form B and C from Chiang Mai (CM), Mae Hong Son (MS) and Phetchaburi (PB).....	58
25. Isoenzyme pattern of ACPH, ALDOX, EST, FUM, GPI, HK, IDH, LDH, MDH, ME, PGM and XDH of adult <i>An. aconitus</i> Form B and C from Chiang Mai (CM), Mae Hong Son (MS) and Phetchaburi (PB).....	59

## LIST OF FIGURES (cont.)

Figures	Page
26. Variable sites in the sequence alignment of the ITS2, COI and COII sequences. Numbering is relative to the alignment. Only those positions differing from the consensus are shown. Indel (insertion or deletion) sites are not shown. An asterisk shows those sites yielding parsimony informative sites. A dot indicates a base pair identical to that of the consensus sequence. Sequence names are defined in Table 10.....	62
27. Salivary gland polytene chromosome of F <sub>1</sub> -hybrid 4 <sup>th</sup> instar larvae of <i>An. aconitus</i> : (A) Form B female (Chiang Mai strain) x Form B male (Phetchaburi strain), (B) Form B female (Chiang Mai strain) x Form C male (Chiang Mai strain), (C) Form B female (Chiang Mai strain) x Form C male (Mae Hong Son strain). All the crosses yielded complete synapsis in all arms, except floating and heterozygous inversion (INV) on 2L (C), which was found in only one preparation from the F <sub>1</sub> -hybrid 4 <sup>th</sup> instar larvae of Form B female (Chiang Mai strain) x Form C male (Mae Hong Son strain).....	66

## LIST OF FIGURES (cont.)

Figures	Page
28. Salivary glands of <i>An. aconitus</i> Form B. (A) Showing free flow <i>P. vivax</i> sporozoites from the squashed salivary glands. Note, the regular spindle-shaped sporozoites (small arrow). (B) Showing sporozoite-like crystals inside the median lobe of salivary glands (small arrow). (C) Showing free flow sporozoite-like crystals from the squashed salivary glands. Note, the irregular, long or short, crystals with blunt or tapered end(s).....	74

## ABBREVIATIONS

°C	degree Celsius
%	percentage
<	less than
>	more than
cm	centimeter
g	gram
hr	hour
kg	kilogram
μl	microliter
μm	micrometer
M	molar
mg	milligram
min	minute
ml	milliliter
mm	millimeter
mm <sup>2</sup>	square millimeter
n	number
s	second
SD	standard deviation
V	voltage

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่  
 Copyright © by Chiang Mai University  
 All rights reserved