


REFERENCES

- Bundarnsin, T., 2006, Delineation of hydrocarbon potential area in the Gulf of Thailand using seismic attribute analysis, Unpublished M. Sc. Thesis, Chiang Mai University.
- Bunopass, S., and Vella, P., 1983, Opening of the Gulf of Thailand, rifting of continental Southeast Asia, and Late Cenozoic tectonics: Jour. Geol. Soc. Thailand, v. 6, p. 1-12.
- Burri, P., 1989, Hydrocarbon potential of Tertiary intermontane basins in Thailand, in Pitak, T.T., ed., Proceedings of international symposium on intermontane basins: Geology and resources: Chiang Mai, Thailand, Chiang Mai University, p. 3-12.
- Bustin, R.M., and Chonchawalit, A., 1995 Formation and tectonic evolution of the Pattani basin, Gulf of Thailand, International Geology Review, v. 37, p. 866-892.
- Chinbunchorn, N., Praditjan, S., and Sattayarak, N., 1989, Petroleum potential of Tertiary intermontane basins in Thailand, In: Thanasuthipitak, T. and Ounchanum, P. (eds.), International symposium on intermontane basin: Geology and Resources, Chiang Mai, Thailand, p.29-42.
- Chopra, S., and Marfurt, K., 2005, Seismic attributes-a historical perspective, Geophysics, v.70, p.3-28.
- Department of Mineral Fuels, Annual report 2005, Petroleum and coal activities in Thailand.
- Hampson-Russel User's Guide, downloaded from, www.hampson-russell.com

- Hellinger, S.J., and Sclater, J.G., 1983, Some comments on two-layer extensional models for the evolution of sedimentary basins: *Jour. Geophys. Res.*, v. 88, p. 8251-8269.
- Kalkomey, C.T., 1997, Potential risks when using seismic attributes as predictors of reservoir properties: *The Leading Edge*, v. 16, p. 247-251.
- Leloup, P.H., Lasassin, R., Tapponier, P., Scharer, U., Dalai, Z., Xiaohan, L., Liangshang, Z., Shaocheng, J., & Trinh, P.T., 1995, The Ailao Shan-Red River shear zone (Yunnan, China), Tertiary transform boundary of Indochina, *Tectonophysics*, v. 251, p. 3-84.
- Oudom-Ugson, P., Bunopas, S., Achalabhuti, C., & Polachan, S., 1986, Petroleum potential in pre-Tertiary basins of Thailand, *ASCOPE, Indonesia*, October 12, p. 1997.
- Packham, G., 1996, Cenozoic Southeast Asia: reconstructing its aggregation and reorganization, Tectonic evolution of southeast Asia, In: Hall, R. and Blundell, D.J (eds.), v. 106, p. 123-152, London: Geological Society of London.
- Polachan, S., and Sattayarak, N., 1989, Strike-slip tectonics and the development of Tertiary basins in Thailand, *International symposium on intermontane basins: Geology and Resources*, Chiang Mai, Thailand, p. 243-253.
- Pradidtan, S., and Dook, R., 1992, Petroleum geology of the northern part of the Gulf of Thailand, In: Piancharoen, C. (ed.), *National conference on "Geologic resources of Thailand: Potential for future development"*, Department of Mineral Resources, 17-24 November 1992, Bangkok, Thailand, p.235-245.

Schultz, P. S., Ronen, S., Hattori, M., and Corbett, C., 1994, Seismic guided estimation of log properties, parts 1, 2, and 3: The Leading Edge, v.13, p.305–310, 674–678 and 770–776.

Tanner, S., Koehler, F., and Sheriff, R.E., 1979, Complex seismic trace analysis, Geophysics, v.44, p.1041-1063.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright © by Chiang Mai University
All rights reserved