

TABLE OF CONTENTS

	Pages
ACKNOWLEDGEMENT	iii
ABSTRACT IN ENGLISH	iv
ABSTRACT IN THAI	vii
LIST OF TABLES	xi
LIST OF ILLUSTRATIONS	xiii
CHAPTER 1 INTRODUCTION	1
1.1 Introduction	1
1.2 Research Objectives	3
CHAPTER 2 LITERATURE REVIEW	4
2.1 Mango	4
2.2 Chemical composition, nutritive and medicine value	6
2.3 Postharvest Storage	7
2.4 Ripening of mangoes	12
2.5 Chilling injury	16
2.6 Temperature management	27
2.7 Thermal properties	33
2.8 Heat transfer theory	38
2.9 Unsteady state heat transfer	40
2.10 Finite Difference method	41
CHAPTER 3 MATERIALS AND METHODS	43
3.1 Determine thermal properties and chemical composition of mango fruit cv Nam Dok Mai Si Thong.	43
3.2 Prediction of internal temperature in mango fruit cv. Nam Dok Mai Si Thong using thermal properties.	53

	Page
3.3 To determine the relationship between thermal properties of mango and chilling injury during storage at 5 and 13 °C.	60
CHAPTER 4 RESULTS AND DISCUSSION	66
4.1 Chemical composition of peel and flesh of Nam Dok Mai Si Thong	66
4.2 Thermal properties of Nam Dok Mai Si Thong Variety	69
4.3 The comparison of thermal properties obtained from the experiment to the calculated values from the chemical composition	75
4.4 The application of numerical methodology in the prediction of temperature changes within Nam Dok mai Si Thong during thermal processing	80
4.5 Determination of the relationship between the chilling injury and thermal properties of Nam Dok Mai Si Thong during storage at 5 and 13°C	87
CHAPTER 5 CONCLUSION	203
REFERENCES	206
APPENDICES	236
CIRRICULUM VITAE	270

LIST OF TABLES

Tables	Page	
2.1	Difference between Nam Dok Mai Si Thong and Nam Dok Mai No.4.	5
2.2	Analysis of the flesh of green & ripe mangoes gave the following Composition (per 100 gms):	6
2.3	Injuries to mango cultivars induced by disinfestation treatment, applied as hot water, vapor heat or forced hot-air treatments.	30
4.1	Chemical compositions of peel and flesh of Nam Dok Mai Si Thong mango	65
4.2	Chemical compositions of green and ripe mangoes of different variety	67
4.3	Nutrition values of the mango flesh between Thai mango (the variety was not specified) and Haden per 100 g fresh weight	68
4.4	Thermal properties of peel and mango flesh of Nam Dok Mai Si Thong variety at 25.0°C.	69
4.5	Effect of moisture on the variation of thermal conductivity of some fruits.	71
4.6	Effect of temperature on the thermal conductivity of some vegetables and fruits.	72
4.7	The thermal diffusivity of some vegetables and fruits.	74
4.8	The comparison of specific heat capacity of Nam Dok Mai Si Thong peel at 25.0°C between the experiment and equation.	75
4.9	The comparison of specific heat capacity of Nam Dok Mai Si Thong flesh at 25.0°C between the experiment and equation	76

Tables		Page
4.10	The comparison of thermal conductivity of Nam Dok Mai Si Thong peel at 25.0°C between the experiment and equation.	77
4.11	The comparison of thermal conductivity of Nam Dok Mai Si Thong mango flesh at 25.0°C between the experiment and equation.	78
4.12	The comparison of thermal diffusivity of Nam Dok Mai Si Thong peel at 25.0°C between the experiment and equation.	79
4.13	The comparison of thermal diffusivity of Nam Dok Mai Si Thong mango flesh at 25.0°C between the experiment and equation.	80

LIST OF ILLUSTRATIONS

Figures		Page
2.1	Mango cv. Nam Dok Mai Si Thong at A: maturity stage and B: ripening stage.	5
2.2	Pathway of ethylene biosynthesis	14
2.3	Constituent of pectin	15
2.4	A sequence of metabolic events leading from a stress-induced alteration in the properties of membrane to observable macroscopic tissue damage	17
2.5	Membrane injury or acclimation resulting from chilling stress	20
3.1	Sample preparation for determined specific heat and thermal conductivity of mango fruit.	46
3.2	Measurement of density by pycnometer.	47
3.3	Protein Analyzer	50
3.4	Muffle furnace	51
3.5	Fat Analyser	52
3.6	Temperature measurement locations.	54
3.7	Heating test with hot water.	54
3.8	Cooling experiment	55
3.9	The distance between plane 0 and plane 1 equaled to half of the thickness of the seed.	58
3.10	Value range in relation to lightness scales	62
3.11	Color chart of natural color system	63
4.1	Schematic cross-section diagram of the mango used in the investigation of temperature profiling by finite difference method.	81

Figures	Page	
4.2	The comparison between the temperature obtained from the experiment (exp) and model (sim) at the surface ($m = 5$) of Nam Dok Mai Si Thong mango in the incubator ($13.0 \pm 0.5^\circ\text{C}$)	82
4.3	The comparison between the temperature obtained from the experiment (exp) and model (sim) at the center of Nam Dok Mai Si Thong mango flesh in the incubator ($13.0 \pm 0.5^\circ\text{C}$).	83
4.4	The comparison between the temperature obtained from the experiment (exp) and model (sim) at the position of Nam Dok Mai Si Thong mango flesh attached to the seed ($m = 1$) in the incubator ($13.0 \pm 0.5^\circ\text{C}$).	83
4.5	Comparison of temperature profile at the surface of Nam Dok Mai Si Thong between the experiment (exp) and predicted value from the model (sim) in the temperature controlled water bath at $48.4 \pm 0.5^\circ\text{C}$.	85
4.6	Comparison of temperature profile at the center of Nam Dok Mai Si Thong flesh ($m = 3$) between the experiment (exp) and predicted value from the model (sim) in the temperature controlled water bath at $48.4 \pm 0.5^\circ\text{C}$.	85
4.7	The comparison between the temperature obtained from the experiment (exp) and model (sim) at the position of Nam Dok Mai Si Thong mango flesh attached to the seed ($m = 1$) in the temperature controlled incubator ($48.4 \pm 0.5^\circ\text{C}$).	86
4.8	Chilling index score of the mango kept at 5°C .	87
4.9	The characteristics of chilling injury observed in Nam Dok Mai Si Thong mango that was stored at 5°C ($1.5\times$ magnification).	89
4.10	Chilling index score of the mango kept at 13°C .	90

Figures	Page
4.11	91
Profile of the electrolyte leakage value at the mango peel of Nam Dok Mai Si Thong mango being kept at 5, 13, and 25°C.	
4.12	92
The relationship between the leakage of electrolyte from the peel of Nam Dok Mai Si Thong mango and the chilling index score when the mango was kept at 5°C	
4.13	94
The relationship between electrolyte leakage of Nam Dok Mai Si Thong peel and the chilling index score of electrolyte at 13°C.	
4.14	96
The profile of luminosity of the mango peel stored at 5, 13 and 25°C.	
4.15	97
The relationship between the chilling index score and L*-value of Nam Dok Mai Si Thong peel being kept at 5°C.	
4.16	97
The relationship between L*-value and electrolyte leakage of the Nam Dok Mai Si Thong peel kept at 5°C.	
4.17	99
The relationship between the chilling index score and L*-value of Nam Dok Mai Si Thong peel being stored at 13°C.	
4.18	99
The relationship between the percentage of electrolyte leakage and L*-value of Nam Dok Mai Si Thong peel being stored at 13°C.	
4.19	100
The profile of a* value of Nam Dok Mai Si Thong peel being kept in the incubator at 5, 13 and 25°C.	
4.20	101
The profile of b* value of Nam Dok Mai Si Thong peel being kept in the incubator at 5, 13 and 25°C.	
4.21	102
The relationship between the chilling index score and b*-value of Nam Dok Mai Si Thong peel being stored at 5°C.	

Figures	Page
4.22	The relationship between the percentage of electrolyte leakage and b*-value of Nam Dok Mai Si Thong peel being stored at 5°C. 103
4.23	The relationship between the chilling index score and b* value of Nam Dok Mai Si Thong peel maintained at 13°C. 104
4.24	The relationship between the electrolyte leakage and b* value of Nam Dok Mai Si Thong peel maintained at 13°C. 104
4.25	The profile of chroma (C*) value for Nam Dok Mai Si Thong peel maintained in the incubator at 5, 13 and 25°C. 105
4.26	The hue angle profile of Nam Dok Mai Si Thong peel being kept in the incubator at 5, 13 and 25°C. 106
4.27	The color change of Nam Dok Mai Si Thong peel being kept at A:5°C, B:13°C. And C:25°C. 107
4.28	The respiratory rate of Nam Dok Mai Si Thong mango being kept in the incubator at 5, 13 and 25°C. 109
4.29	The change of L*-value of Nam Dok Mai Si Thong mango flesh stored in a temperature controlled incubator at 5, 13 and 25°C. 112
4.30	The change in a* value of Nam Dok Mai Si Thong mango flesh being stored at 5, 13 and 25°C in the temperature controlled incubator. 113
4.31	The profile of b* value of Nam Dok Mai Si Thong mango flesh stored in the temperature controlled incubator at 5, 13 and 25°C. 114
4.32	The profile of C* value for Nam Dok Mai Si Thong mango stored in the incubator at 5, 13 and 25°C. 115
4.33	The profile of hue angle value of Nam Dok Mai Si Thong mango flesh stored in the incubator at 5, 13 and 25°C. 116

Figures	Page
4.34	The change in hue angle value of Nam Dok Mai Si Thong mango flesh stored in the incubator at A:5, B: 13 and C: 25 °C. 118
4.35	The change of Nam Dok Mai Si Thong's pericarp and mango flesh stored in a temperature controlled cabinet at $5 \pm 0.5^\circ\text{C}$. 119
4.36	The change of Nam Dok Mai Si Thong's pericarp and mango flesh stored in a temperature controlled cabinet at $13 \pm 0.5^\circ\text{C}$. 120
4.37	The change of Nam Dok Mai Si Thong's pericarp and mango flesh stored in a temperature controlled cabinet at $25 \pm 0.5^\circ\text{C}$. 121
4.38	Firmness profile of Nam Dok Mai Si Thong mango flesh in the incubator at 5, 13 and 25°C . 122
4.39	The relationship between the change in firmness of mango flesh and the respiratory rate of Nam Dok Mai Si Thong mango stored in the incubator at 5°C . 124
4.40	The relationship between the change in firmness of mango flesh and the respiratory rate of Nam Dok Mai Si Thong mango stored in the incubator at 13°C . 125
4.41	The relationship between the change in firmness of mango flesh and the respiratory rate of Nam Dok Mai Si Thong mango stored in the incubator at 25°C . 125
4.42	The weight loss profile of Nam Dok Mai Si Thong mango stored in the incubator at 5, 13 and $25 \pm 0.5^\circ\text{C}$. 126
4.43	The moisture content profile of Nam Dok Mai Si Thong peel being maintained in the incubator at 5, 13 and 25°C . 128
4.44	The moisture content profile of Nam Dok Mai Si Thong mango flesh being maintained in the incubator at 5, 13 and 25°C . 129

Figures	Page
4.45	The profile of total soluble solid of Nam Dok Mai Si Thong mango stored in the incubator at 5, 13 and 25°C. 130
4.46	The relationship between the total soluble solid and the respiratory rate of Nam Si Thong mango being stored at 13°C. 131
4.47	The relationship between the total soluble solid and the respiratory rate of Nam Si Thong mango being stored at 25°C. 132
4.48	The relationship between the total soluble solid and the respiratory rate of Nam Si Thong mango being stored at 5°C. 133
4.49	The profile of titable acidity content of Nam Dok Mai Si Thong mango stored in the incubator at 5, 13 and 25°C. 133
4.50	The pH profile of Nam Dok Mai Si Thong mango stored in the incubator at 5, 13 and 25°C. 134
4.51	The relationship between the titratable citric acid content and the pH of Nam Dok Mai Si Thong mango being stored at 25°C. 135
4.52	The relationship between the titratable citric acid content and the pH of Nam Dok Mai Si Thong mango being stored at 13°C. 136
4.53	The relationship between the titratable citric acid content and the pH of Nam Si Thong mango being stored at 5°C. 137
4.54	The electrolyte leakage profile of Nam Dok Mai Si Thong mango flesh stored in the temperature controlled incubator at 5, 13 and 25°C. 138
4.55	The relationship between the respiratory rate and electrolyte leakage of Nam Si Thong mango flesh being stored at 5°C. 139

Figures		Page
4.56	The relationship between the profile of b* value and electrolyte leakage of Nam Si Thong mango flesh being stored at 5°C.	139
4.57	The relationship between the profile of firmness and electrolyte leakage of Nam Si Thong mango flesh being stored at 5°C.	140
4.58	The relationship between the profile of total soluble solid (TSS) and electrolyte leakage of Nam Si Thong mango flesh being stored at 5°C.	140
4.59	The relationship between the profile of b* value and electrolyte leakage of Nam Si Thong mango flesh being stored at 25°C.	141
4.60	The relationship between the profile of b* value and electrolyte leakage of Nam Si Thong mango flesh being stored at 13°C.	142
4.61	The relationship between the profile of firmness and electrolyte leakage of Nam Si Thong mango flesh being stored at 25°C.	143
4.62	The relationship between the profile of firmness and electrolyte leakage of Nam Si Thong mango flesh being stored at 13°C.	143
4.63	The relationship between the profile of respiratory rate and electrolyte leakage of Nam Si Thong mango flesh being stored at 25°C.	144
4.64	The relationship between the profile of respiratory rate and electrolyte leakage of Nam Si Thong mango flesh being stored at 13°C.	145
4.65	The relationship between the profile of total soluble solid (TSS) and electrolyte leakage of Nam Si Thong mango flesh being stored at 25°C.	146

Figures		Page
4.66	The relationship between the profile of total soluble solid (TSS) and electrolyte leakage of Nam Si Thong mango flesh being stored at 13°C.	146
4.67	The profile of specific heat capacity of Nam Dok Mai Si Thong peel stored in the incubator at 5, 13 and 25°C.	147
4.68	The relationship between the specific heat capacity and b^* value of Nam Dok Mai Si Thong peel being stored at 13°C.	148
4.69	The relationship between the specific heat capacity (C_p) and b^* value of Nam Dok Mai Si Thong peel being stored at 25°C.	149
4.70	The relationship between the specific heat capacity (C_p) of the peel and the respiration rate of Nam Dok Mai Si Thong peel being stored at 13°C.	149
4.71	The relationship between the specific heat capacity (C_p) of the peel and the respiration rate of Nam Dok Mai Si Thong peel being stored at 25°C.	150
4.72	The relationship between the specific heat capacity (C_p) of the peel and the chilling index score of Nam Si Thong peel being stored at 13°C.	151
4.73	The relationship between the specific heat capacity (C_p) of the peel and the electrolyte leakage of Nam Si Thong peel being stored at 13°C.	151
4.74	The relationship between the specific heat capacity (C_p) of the peel and the electrolyte leakage of Nam Si Thong peel being stored at 25°C.	152
4.75	The relationship between the specific heat capacity (C_p) of the peel and the chilling index score of Nam Si Thong peel being stored at 5°C.	153

Figures	Page
4.76	The relationship between the specific heat capacity (C_p) of the peel and b^* value of Nam Dok Mai Si Thong peel being stored at 5°C. 153
4.77	The relationship between the specific heat capacity (C_p) of the peel and respiratory rate of Nam Dok Mai Si Thong peel being stored at 5°C. 154
4.78	The relationship between the specific heat capacity (C_p) of the peel and the electrolyte leakage of Nam Dok Mai Si Thong peel being stored at 5°C. 155
4.79	The profile of specific heat capacity of Nam Dok Mai Si Thong mango flesh stored in the incubator at 5, 13 and 25°C. 156
4.80	The relationship between the specific heat capacity and moisture content of Nam Dok Mai Si Thong mango flesh being stored at 5°C. 157
4.81	The relationship between the specific heat capacity and b^* value of Nam Dok Mai Si Thong mango flesh being stored at 5°C. 158
4.82	The relationship between the specific heat capacity and respiratory rate of Nam Dok Mai Si Thong mango flesh being stored at 5°C. 158
4.83	The relationship between the specific heat capacity and total soluble solid content of Nam Dok Mai Si Thong mango flesh being stored at 5°C. 159
4.84	The relationship between the specific heat capacity and electrolyte leakage of Nam Dok Mai Si Thong mango flesh being stored at 5°C. 159
4.85	The relationship between the specific heat capacity and firmness of Nam Dok Mai Si Thong mango flesh being stored at 5°C. 160

Figures		Page
4.86	The relationship between the specific heat capacity and b^* value of Nam Dok Mai Si Thong mango flesh being stored at 13°C.	161
4.87	The relationship between the specific heat capacity and b^* value of Nam Dok Mai Si Thong mango flesh being stored at 25°C.	161
4.88	The relationship between the specific heat capacity and respiratory rate of Nam Dok Mai Si Thong mango flesh being stored at 13°C.	162
4.89	The relationship between the specific heat capacity and respiratory rate of Nam Dok Mai Si Thong mango flesh being stored at 25°C.	162
4.90	The relationship between the specific heat capacity and total soluble solid of Nam Dok Mai Si Thong mango flesh being stored at 13°C.	163
4.91	The relationship between the specific heat capacity and total soluble solid of Nam Dok Mai Si Thong mango flesh being stored at 25°C.	163
4.92	The relationship between the specific heat capacity and electrolyte leakage of Nam Dok Mai Si Thong mango flesh being stored at 13°C.	164
4.93	The relationship between the specific heat capacity and electrolyte leakage of Nam Dok Mai Si Thong mango flesh being stored at 25°C.	164
4.94	The relationship between the specific heat capacity and firmness of Nam Dok Mai Si Thong mango flesh being stored at 13°C.	165

Figures	Page
4.95	165
The relationship between the specific heat capacity and firmness of Nam Dok Mai Si Thong mango flesh being stored at 25°C.	
4.96	166
The profile of thermal conductivity of Nam Dok Mai Si Thong peel stored in the temperature controlled incubator at 5, 13 and 25°C	
4.97	168
The relationship between the thermal conductivity (k) and chilling injury of Nam Dok Mai Si Thong peel being stored at 5°C.	
4.98	168
The relationship between the thermal conductivity (k) and electrolyte leakage of Nam Dok Mai Si Thong peel being stored at 5°C.	
4.99	169
The relationship between the thermal conductivity (k) and b* value of Nam Dok Mai Si Thong peel being stored at 5°C	
4.100	169
The relationship between the thermal conductivity (k) of the peel and respiratory rate of Nam Dok Mai Si Thong being stored at 5°C.	
4.101	170
The relationship between the thermal conductivity (k) and b* value of Nam Dok Mai Si Thong peel being stored at 13°C.	
4.102	171
The relationship between the thermal conductivity (k) and b*value of Nam Dok Mai Si Thong peel being stored at 25°C.	
4.103	171
The relationship between the thermal conductivity (k) and respiratory rate of Nam Dok Mai Si Thong peel being stored at 13°C.	

Figures		Page
4.104	The relationship between the thermal conductivity (k) and respiratory rate of Nam Dok Mai Si Thong peel being stored at 25°C.	172
4.105	The relationship between the thermal conductivity (k) and chilling injury of Nam Dok Mai Si Thong peel being stored at 13°C.	173
4.106	The relationship between the thermal conductivity (k) and electrolyte leakage of Nam Dok Mai Si Thong peel being stored at 13°C.	173
4.107	The relationship between the thermal conductivity (k) and electrolyte leakage of Nam Dok Mai Si Thong peel being stored at 25°C.	174
4.108	The profile of thermal conductivity of Nam Dok Mai Si Thong of mango peel stored in the incubator at 5, 13 and 25°C.	175
4.109	The relationship between the thermal conductivity (k) and b* value of Nam Dok Mai Si Thong mango flesh being stored at 5°C.	176
4.110	The relationship between the thermal conductivity (k) and respiratory rate of Nam Dok Mai Si Thong mango flesh being stored at 5°C.	176
4.111	The relationship between the thermal conductivity (k) and firmness of Nam Dok Mai Si Thong mango flesh being stored at 5°C.	177
4.112	The relationship between the thermal conductivity and total soluble solid content of Nam Dok Mai Si Thong mango flesh being stored at 5°C.	177

Figures		Page
4.113	The relationship between the thermal conductivity and electrolyte leakage of Nam Dok Mai Si Thong mango flesh being stored at 5°C.	178
4.114	The relationship between the thermal conductivity and b* value of Nam Dok Mai Si Thong mango flesh being stored at 13°C.	179
4.115	The relationship between the thermal conductivity and b* value of Nam Dok Mai Si Thong mango flesh being stored at 25°C.	179
4.116	The relationship between the thermal conductivity and respiratory rate of Nam Dok Mai Si Thong mango flesh being stored at 13°C.	180
4.117	The relationship between the thermal conductivity and respiratory rate of Nam Dok Mai Si Thong mango flesh being stored at 25°C.	180
4.118	The relationship between the thermal conductivity and firmness of Nam Dok Mai Si Thong mango flesh being stored at 13°C.	181
4.119	The relationship between the thermal conductivity and firmness of Nam Dok Mai Si Thong mango flesh being stored at 25°C.	181
4.120	The relationship between the thermal conductivity and total soluble solid content of Nam Dok Mai Si Thong mango flesh being stored at 13°C.	182
4.121	The relationship between the thermal conductivity and total soluble solid content of Nam Dok Mai Si Thong mango flesh being stored at 25°C.	182

Figures	Page
4.122	183
The relationship between the thermal conductivity and electrolyte leakage of Nam Dok Mai Si Thong mango flesh being stored at 13°C.	
4.123	183
The relationship between the thermal conductivity and electrolyte leakage of Nam Dok Mai Si Thong mango flesh being stored at 25°C.	
4.124	184
The density profile of Nam Dok Mai Si Thong peel stored in the incubator at 5, 13 and 25°C.	
4.125	185
The density profile of Nam Dok Mai Si Thong mango flesh stored in the incubator at 5, 13 and 25°C.	
4.126	186
The profile of thermal diffusivity of Nam Dok Mai Si Thong pericarp stored in the temperature controlled incubator at 5, 13 and 25°C.	
4.127	187
The relationship between the thermal diffusivity and chilling injury score of Nam Dok Mai Si Thong of mango peel being stored at 5°C.	
4.128	187
The relationship between the thermal diffusivity and electrolyte leakage of Nam Dok Mai Si Thong of mango peel being stored at 5°C.	
4.129	189
The relationship between the thermal diffusivity and chilling injury score of Nam Dok Mai Si Thong of mango peel being stored at 13°C.	
4.130	189
The relationship between the thermal diffusivity and chilling injury score of Nam Dok Mai Si Thong of mango peel being stored at 25°C.	
4.131	190
The relationship between the thermal diffusivity and b* value of Nam Dok Mai Si Thong of mango peel being stored at 13°C.	

Figures	Page
4.132	190
The relationship between the thermal diffusivity and b* value of Nam Dok Mai Si Thong of mango peel being stored at 25°C.	
4.133	191
The relationship between the thermal diffusivity and respiratory rate of Nam Dok Mai Si Thong mango fruit being stored at 13°C.	
4.134	191
The relationship between the thermal diffusivity and respiratory rate of Nam Dok Mai Si Thong mango fruit being stored at 25°C.	
4.135	192
The relationship between the thermal diffusivity and electrolyte leakage of Nam Dok Mai Si Thong of mango peel being stored at 13°C.	
4.136	192
The relationship between the thermal diffusivity and electrolyte leakage of Nam Dok Mai Si Thong of mango peel being stored at 25°C.	
4.137	193
The profile of thermal diffusivity of Nam Dok Mai Si Thong mango flesh stored in the temperature controlled incubator at 5, 13 and 25°C.	
4.138	194
The relationship between the thermal diffusivity and firmness of Nam Dok Mai Si Thong mango flesh being stored at 5°C.	
4.139	195
The relationship between the thermal diffusivity and firmness of Nam Dok Mai Si Thong mango flesh being stored at 13°C.	
4.140	195
The relationship between the thermal diffusivity and firmness of Nam Dok Mai Si Thong mango flesh being stored at 25°C.	

Figures	Page
4.141	196
The relationship between the thermal diffusivity and b^* value of Nam Dok Mai Si Thong mango flesh being stored at 5°C.	
4.142	196
The relationship between the thermal diffusivity and b^* value of Nam Dok Mai Si Thong mango flesh being stored at 13°C.	
4.143	197
The relationship between the thermal diffusivity and b^* value of Nam Dok Mai Si Thong mango flesh being stored at 25°C.	
4.144	197
The relationship between the thermal diffusivity and total soluble solid content of Nam Dok Mai Si Thong mango flesh being stored at 5°C.	
4.145	198
The relationship between the thermal diffusivity and total soluble solid content of Nam Dok Mai Si Thong mango flesh being stored at 13°C.	
4.146	198
The relationship between the thermal diffusivity and total soluble solid content of Nam Dok Mai Si Thong mango flesh being stored at 25°C.	
4.147	199
The relationship between the thermal diffusivity and respiratory rate of Nam Dok Mai Si Thong mango fruit being stored at 5°C.	
4.148	199
The relationship between the thermal diffusivity and respiratory rate of Nam Dok Mai Si Thong mango fruit being stored at 13°C.	
4.149	200
The relationship between the thermal diffusivity and respiratory rate of Nam Dok Mai Si Thong mango fruit being stored at 25°C.	

Figures		Page
4.150	The relationship between the thermal diffusivity and electrolyte leakage of Nam Dok Mai Si Thong mango flesh being stored at 5°C.	201
4.151	The relationship between the thermal diffusivity and electrolyte leakage of Nam Dok Mai Si Thong mango flesh being stored at 13°C.	201
4.152	The relationship between the thermal diffusivity and electrolyte leakage of Nam Dok Mai Si Thong mango flesh being stored at 25°C.	202