

REFERENCES

- Adema, GJ, Boer, A.J., Hullenaar, P., Denijn, R., Ruiter, M., Vogel, D.J., Figdor, AM. 1993. CG Melanocyte lineage-specific antigens recognized by monoclonal antibodies NKI-beteb, HMB-50 and HMB-45 are encoded by a single cDNA. *Am. j pathol.* **143**, 1579–1585.
- ATSDR., 1999. Toxicological Profile for Formaldehyde (Final Report). NTIS Accession No. PB99-166654. Atlanta, GA: Agency for Toxic Substances and Disease Registry. 451.
- Ballacha, J., Greuter B., Schultza E., and Jaeschke W. ,1999. Variations of uptake rates in benzene diffusive sampling as a function of ambient conditions. *Sci. total environ.* **243/244**, 203-217.
- Balmat, J.L. and Meadows G.W., 1985. Monitoring of Formaldehyde in Air. *AIHA.* **46**, 578 – 584.
- Bertoni, G., Ciuchini C., Di Palo V. and Possanzini M., 2005. Development of a Passive Sampler for Long-Term Measurements of Formaldehyde and Total Oxidants in Air. *Chromatographia.* **61**, 385–389.
- Brace, S. and Peterson, D.L., 1998. Spatial patterns of tropospheric ozone in the Mount Rainier region of the Cascade Mountains. *Atmos. Environ.* **32 (21)**, 3629 – 3637.
- California environmental protective agency (CEPA). 2004. *Indoor Air Quality Guideline*. California Air Resources Board, No.1
- Cao X.L., and Hewitt C.N. 1991. Application of passive samplers to the monitoring of low concentration organic vapours in indoor and, ambient air: A review, *Environ. Technol.* **12**, 1055-1062.

Chansa M. K. and James K. H., 1991. Permeation-solid adsorbent sampling and GC analysis of formaldehyde. *Talanta.* **38**, 1381-1386.

ComiteÂ EuropeÂen de Normalisation, 1999. Ambient Air Quality Diffusive Samplers for the Determination of Concentrations of Gases or Vapours Part 3: Guide for Selection, Use and Maintenance, prEN 13528-3, CEN, Brussels,

Current Intelligence Bulletin No.1981. 34, Formaldehyde: Evidence of Carcinogenicity", April 15, U.S. Dept. of Health and Human Services, Public Health Service, Center for Disease Control, NIOSH

De Santis, F., Dogeroglu, T.S., Menichelli, Vazzana, C. and Allegrini, I., 2001. The use of a new passive sampler for ozone and nitrogen oxides monitoring in ecological effects research. In: Proceedings of the International Symposium on Passive Sampling of Gaseous Air Pollutants in Ecological Research. The Scientific World 1, 475–482.

Dillon, G.C.G. 1994. Compost quality Objectives Study. Final report to Alberta environmental protection, Alberta agriculture, Alberta health, and Edmonton Board of Health. *Canada and E&A Environmental Consultants, Inc.* Canton, MA.

Fox. C. B. 1873. Ozone and Ant ozone. J. and A. Cherchill. London.

Freeland, L.T.1977. Industrial Hygiene Calibration Manifold. *Am. Ind. Hyg. Assoc. J.* **38**,712-720.

Fritz. F.,1966. Spot Tests in Organic Analysis. 7th Ed. Newyork. American Elsevier Publishing Co.

Fuller, E. N.; Schettler, P. D.; Giddings, J. C. 1966. A new method for prediction of binary gas phase diffusion coefficients. *Ind. Eng. Chem.* **58** (5), 18-27.

- Gair, A.J. and Penkett, S.A., 1995. The effects of wind speed and turbulence on the performance of diffusion tube samplers. *Atmos. Environ.* **29**, 2529–2533.
- Geisling, K.L. and Rappaport, S.M., 1982. A passive sampling device for determining formaldehyde in indoor air. *Environ. Int.* **8**, 153-158.
- Gerboles, M., Buzica, D., Amantini, L., Plaisance, H. and Marfaing, H., 2004. Intercomparison exercises for the monitoring of ambient nitrogen dioxide using diffusive samplers. *Pollut. Atmos.* **183**, 345–359.
- Gibson, L.T. and Brokerhof, A.W., 2001. A passive tube-type sampler for the determination of formaldehyde vapours in museum enclosures. *Stud. Conserv.* **46**, 289-303.
- Gibbons, R. D. 1996."The Problem with U.S. EPA's Method Detection Limit", *Am. Environ. Lab.* February.
- Gillettr . W., Kreibich,. H. and Ayers . G . P. 2000. Measurement of Indoor Formaldehyde Concentrations with a Passive Sampler. *Environ. Sci. Technol.* **34**, 2051-2056.
- Gilliland, E. R., 1934. Diffusion Coefficients in Gaseous Systems. *Ind. Eng. Chem.* **26**, 681-685.
- Gorecki, T. and Namiesnik, J., 2002. Passive sampling. *Trac. Anal. Chem.* **21**, 276-291.
- Grosjean, D. and Williams, E.L. 1992. A passive sampler for airborne formaldehyde. *Atmos. Environ.* **26**, 2923-2928.
- Heal, M.R., and Cape, J.N. 1997. A numerical evaluation of chemical interferences in the measurement of ambient nitrogen dioxide by passive diffusion samplers. *Atmos. Environ.* **31 (13)**, 1911-1924.

HSDB, 2001: Hazardous Substances Data Bank, National Library of Medicine.

TOXNET (Toxicology Data Network). *Ammonia*. Retrieved from:

<http://toxnet.nlm.nih.gov/>.

IARC. 1995. Wood Dust and Formaldehyde. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, Lyon, France: International Agency for Research on Cancer. **262**, 405.

IARC. 1987. Overall Evaluations of Carcinogenicity. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, Supplement 7. Lyon, France: International Agency for Research on Cancer. 440.

IARC. 1982. Monograph on the Evaluation of the Carcinogenic Risk of Chemical to Humans, Some Industrial Chemicals and Dyestuffs" International Agency for Research on Cancer: Lyon. **29**, 345-389.

IARC. 1982. Some Industrial Chemicals and Dyestuffs. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans. Lyon, France: International Agency for Research on Cancer. **29**, 416.

IARC. 1973. Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Man, some Aromatic Amines Hydrazine and Related Substances, N-nitroso Compounds and Miscellaneous Alkylating Agents", International Agency for Research on Cancer: Lyon. **4**, 231-237.

Jones. A.P. 1999, Indoor air quality and Health. *Atmos. Environ.* **33**, 4535–4564.

Kinyanta. C. M. and James H. K. 1991, Permeation-solid adsorbent sampling and GC analysis of formaldehyde. *Talanta*. **38**, 1381-1386.

Koutrakis, P. Sioutas, C. Ferguson S.T. and Wolfson, J.M. 1993. Development and evaluation of a glass honeycomb denuder/filter pack sampler to collect atmospheric gases and particles, *Environ. Sci. Technol.* **27**, 2497–2501.

Kring, E.V., Thornley, G.D., Dessenberger, C., Lautenberger, W.J. and Ansul, G.R., 1982. *Am. Ind. Hyg. Assoc.*, **43** 786-795.

Krochmal, D. and Goarski, L. 1991. Determination of nitrogen dioxide in ambient air by use of a passive sampling technique and triethanol- amine as absorbent. *Environ. Sci. Technol.* **25**, 531-535.

Krochmal, D. and Kalina, A. 1997. A method of nitrogen dioxide and sulphur dioxide determination in ambient air by use of passive samplers and ion chromatography. *Atmos Environ.* **31**, 3473 – 348.

Krupa S.V. and Legge. A.H. 2000. Passive sampling of ambient, gaseous air pollutants: an assessment from an ecological perspective. *Environ. Pollut.* **107**, 31-45.

Levin, J. O., Lindahl, R., and Andersson, K. 1986. A passive sampler for formaldehyde in air using 2,4-dinitrophenylhydrazine-coated glass Fiber Filters.

Environ. Sci. Technol. **20**, 1273-1276.

Lodge, J. P. 1989. Methods of Air Sampling and Analysis, Third Edition, Ed..274.

Monn, C., Hangartner, M., 1990. Passive sampling for ozone. *J Air. Waste. Manage.* **40(3)**, 357-358.

Namiesnik J., Gorecki, T. and E. Kozlowski, 1984. Passive dosimeters and approach to atmospheric pollutants analysis. *Sci. Total. Environ.* **38**, 225–258.

- Nosal, M., 1984. Atmosphere±biosphere Interface: Analytical Design and a Computerized Regression Model for Lodgepole Pine Response to Chronic, Atmospheric SO₂ Exposure (RMD Report 83/26 and 83/27). Research Management Division, Alberta Environment, Edmonton, Alberta, Canada.
- Palmes, E.D., Gunnison, A.F., DiMattio, J. and Tomczyk, C., 1976. Personal sampler for nitrogen dioxide. *Am. Ind. Hyg. Assoc. J.* **37**, 570–577.
- Plaisance, H., Minguy, A., Garcia-Fouque, S. and Galloo, J.C. 2004. Influence of meteorological factors on the NO₂ measurements by passive diffusion tube. *Atmos. Environ.* **38**, 573–580.
- Potter W. and Karst K, 1996. *Anal. Chem.* **68**, 3354-3358.
- Reeuwijk, V., Fisher, H., Harssema, P. H., Briggs, H., Smallbone, D. J. And Labret, E. 1998. Field comparison of two NO₂ Sampler to asses special variation. *Environ. Monit. Assess.* **50**, 37-51.
- Runeckles, V.C. and Bowen, P.A. 1999. The use of calibrated passive monitors to assess crop loss due to ozone in rural locations. In: Agrawal, S.B., Agrawal, M. (Eds.), Environmental Pollution and Plant Responses. Lewis Publishers, Boca Raton, FL (in press).
- Sexton, Ken, Kai-Shen, Liu, Myrto and Petreas X., 1986 . Formaldehyde concentrations inside private residences: A mail-out approach to indoor air Monitoring. *J Air. Pollut. Assoc.* **36**, 698-704.
- Shih W. T. and Shane S. Q. H., 1999. A new passive sampler for aldehydes. *Am. Ind. Hyg. Assoc. J.* **60**, 463–473.

Thammakhet C., Muneesawang V., Thavarungkul and P. Kanatharana P 2006. Cost effective passive sampling device for volatile organic compounds monitoring.

Atmos. Environ. **40**, 4589–4596.

Wilke, C. R., Lee, C. Y. 1955. *Ind. Eng. Chem.*, **47/6**, 1253-1257.

Willems, J. J. H. 1993. Low-cost methods for measuring air pollutants (a training program). Report R-635. Wageningen (NI) Agricultural University, Department of Air Quality, Wageningen, Netherlands.

Zhou, J., Smith, S., 1997. Measurement of ozone concentrations in ambient air using a badge-type passive monitor. *J Air. Waste. Manit. Assoc.* **47**, 697-703.

<http://www.apvma.gov.au>

http://www.healthgoods.com/Education/Healthy_Home_Information/Indoor_Air_Quality/indoor_pollutant_formaldehyde.htm

<http://en.wikipedia.org/wiki/Formaldehyde>)

<http://www.urbandictionary.com/define.php?term=cross-link>

จัดทำโดย คณิตศาสตร์ มหาวิทยาลัยเชียงใหม่
 Copyright[©] by Chiang Mai University
 All rights reserved