

TABLE OF CONTENTS

TITLE	PAGE
Acknowledgement	iii
Abstract in English	v
Abstract in Thai	vii
Table of contents	ix
List of tables	xiii
List of figures	xiv
CHAPTER 1 INTRODUCTION	1
1.1 Background	1
1.2 Research objectives	2
CHAPTER 2 LITERATURE REVIEWS	3
2.1 The longan fruit	3
2.2 Fruit growth and development	4
2.3 Longan cultivars	5
1. Kra Loke longan	6
2. Kra-Dook longan	8
3. Pseudo-Kra Loke longan	8
4. Sai Nam Peung longan	8

TABLE OF CONTENTS (CONTINUED)

	PAGE
5. Other longan cultivars	9
2.4 Postharvest disease of longan	9
1. Disease-causing microorganisms	9
2. Storage temperature	10
3. Wound and contusion during transportation and storage	10
2.5 Fruit infection	12
2.6 Changes in ultrastructure	13
CHAPTER 3 MATERIAL AND METHODS	16
3.1 Characteristic and surface appearance of the longan fruit	16
3.1.1 Microscopic level	16
3.1.2 Ultrastructure level using the scanning electron microscopy (SEM)	17
3.2 Isolation and Identification of longan pathogenic fungi	17
3.3 Pathogenicity of the isolated fungi and disease development	18
3.4 Identification of a rot causing fungal isolate	19
3.5 Infection process of <i>Pestalotiopsis</i> sp. MLP on a pericarp and stem- end of harvested longan fruits	20
3.6 Sulfur dioxide fumigation effect on disease development of harvested longan fruit	23
3.6.1 Surface appearance changes of longan fruit fumigated with SO ₂	23

TABLE OF CONTENTS (CONTINUED)

	PAGE
3.6.2 Effect of SO ₂ on disease development on the inoculated longan fruits	24
3.6.3 Effect of SO ₂ on fungal growth <i>in vitro</i>	24
CHAPTER 4 RESULTS	26
4.1 Characteristic and surface appearance of the longan fruit	26
4.2 Isolation and Identification of longan pathogenic fungi	32
4.3 Pathogenicity of the isolated fungi and disease development	42
4.4 Identification of a rot causing fungal isolates	45
4.5 Infection process of a selected fungus on a pericarp and stem-end of harvested longan fruits	47
4.6 Sulfur dioxide fumigation effect on disease development of harvested longan fruits	55
4.6.1 Surface appearance changes of longan fruit fumigated with SO ₂	55
4.6.2 Effect of SO ₂ on disease development on the inoculated longan fruits	56
4.6.3 Effect of SO ₂ on fungal growth <i>in vitro</i>	57
CHAPTER 5 DISCUSSIONS	58
CHAPTER 6 CONCLUSIONS	63

TABLE OF CONTENTS (CONTINUED)

	PAGE
REFERENCES	64
APPENDIX A: Chemical reagents	75
APPENDIX B: A Simplified key to <i>Pestalotiopsis</i> and Related Genera	76
APPENDIX C: Paraffin section method	79
CURRICULUM VITAE	82

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
 Copyright© by Chiang Mai University
 All rights reserved

LIST OF TABLES

TABLE	PAGE
2.1 Major postharvest pathogens of longan fruit	11
2.2 Longan Fruit Composition at Harvest.	12
4.1 Genera of fungi isolated from the pericarp and stem-end of longan fruit (<i>Dimocarpus longan</i> Lour. cv. Daw.) using tissue transplanting method incubate for 2-3 days	33
4.2 The pathogenicity of fungi isolated from longan fruits (<i>Dimocarpus longan</i> Lour. cv. Daw)	44

LIST OF FIGURES

FIGURE	PAGE
2.1 Sigmoid curve of longan fruit growth and development.	5
4.1 Scanning electron micrographs of longan pericarp surface structure at selected intervals during fruit development.	28
4.2 Longan fruit surface	33
4.3 Scanning electron micrographs of longan fruit surface	29
4.4 Cross-section of mature longan fruit's pericarp 194 days after anthesis	30
4.5 Cross-section of mature longan fruit's pericarp 194 days after anthesis	31
4.6 <i>Lasiodiplodia</i> sp. (a) colony on PDA, 3 days, 25 °C; (b) conidia	36
4.7 <i>Fusarium</i> sp. (a) colony on PDA, 7 days, 25 °C; (b) macroconidia and microconidia	37
4.8 <i>Penicillium</i> sp (a) colony on PDA, 7 days, 25 °C; (b) penicilli and conidia	37
4.9 <i>Aspergillus</i> sp.LK1 (a) colony on PDA, 7 days, 25 °C; (b) heads and conidia	38
4.10 <i>Aspergillus</i> sp. LP2 (a) colony on PDA, 7 days, 25 °C; (b) heads and conidia	38
4.11 <i>Cladosporium</i> sp. (a) colony on PDA, 7 days, 25 °C; (b) conidia	39
4.12 <i>Colletotrichum</i> sp. (a) colony on MYA, 7 DAYS, 25°C ; (b) conidia	39
4.13 <i>Pestalotiopsis</i> sp. (a) colony on PDA, 7 days, 25 °C; (b) conidia	40

LIST OF FIGURES (CONTINUED)

	PAGE
4.14 <i>Phomopsis</i> sp(a) colony on PDA, 7 days, 25 °C; (b) conidia	40
4.15 <i>Trichoderma</i> sp. (a) colony on PDA, 7 days, 25 °C; (b) conidia	41
4.16 <i>Verticillium</i> sp. (a) colony on PDA, 7 days, 25 °C; (b) conidia	41
4.17 <i>Mucor</i> sp. (a) and <i>Rhizopus</i> sp. (b) colonies on PDA, 7 days, 25 °C	42
4.18 Disease symptoms on detached fruit 48 hours after being inoculated with <i>L. theobromae</i> LP20 on the fruit surface	43
4.19 Disease symptoms on detached fruit 48 hours after being inoculated with <i>L. theobromae</i> LP20 on the stem-end	43
4.20 Disease symptoms developed on detached longan fruits after being inoculated with <i>Pestalotiopsis</i> sp.MLP	45
4.21 Colonial growth of <i>L. theobromae</i> LP20	46
4.22 Light micrographs of <i>L. theobromae</i> LP20.	46
4.23 Scanning electron micrographs of <i>L. theobromae</i> LP20	47
4.24 Light micrographs of germinated spores of <i>Pestalotiopsis</i> MLP	48
4.25 Stereo micrographs of longan fruit peel surfaces infected with <i>Pestalotiopsis</i> sp. MLP	48
4.26 Spores germination of <i>Pestalotiopsis</i> sp. MLP was found germ tubes at 5 hours after the inoculation	49
4.27 Cross-section view of the inoculated longan fruit peel with conidia of <i>Pestalotiopsis</i> sp. MLP	49

LIST OF FIGURES (CONTINUED)

	PAGE
4.28 Scanning micrographs of germinated spore of <i>Pestalotiopsis</i> sp. MLP on the surface of longan fruit sampled	50
4.29 The conidia of <i>Pestalotiopsis</i> sp. MLP were observed on the longan fruit surfaces	51
4.30 Scanning micrographs shows early infection stages of the <i>Pestalotiopsis</i> sp. MLP	51
4.31 Scanning micrographs showed infection stages of the <i>Pestalotiopsis</i> sp. MLP	52
4.32 Scanning micrograph showed (a, b) droplet-like spore masses of <i>Pestalotiopsis</i> sp. MLP on the skin surface of infected longan fruit, 48 hours after the inoculation	53
4.33 Scanning micrograph showed the skin surface of longan fruit well-covered with abundant mycelia of <i>Pestalotiopsis</i> sp. MLP at 48 hours after the inoculation	53
4.34 Scanning micrographs showed the processes of internal breakdown of fruit tissue after inoculated with <i>Pestalotiopsis</i> sp. MLP	54
4.35 Scanning micrographs of the epiphytic fungal hypha on the outer skin texture of the peel of the SO ₂ – fumigated longan fruit	55
4.36 Scanning micrographs shows the blistered spores of <i>Pestalotiopsis</i> sp. MLP presented on the inoculated fruits with SO ₂ -fumigation	56
4.37 Scanning micrographs showed the filamentous hypha of <i>Pestalotiopsis</i> sp. MLP on cultured PDA	57