

## TABLE OF CONTENTS

	Page
<b>ACKNOWLEDGMENTS</b>	iii
<b>ABSTRACT (ENGLISH)</b>	iv
<b>ABSTRACT (THAI)</b>	vi
<b>LIST OF TABLES</b>	xi
<b>LIST OF FIGURES</b>	xiv
<b>ABBREVIATIONS AND SYMBOLS</b>	xvii
<b>CHAPTER I: INTRODUCTION</b>	
1.1 Solid-phase extraction (SPE)	1
1.1.1 The sample preparation	1
1.1.2 Solid-phase extraction	2
1.1.3 Advantages of solid phase extraction over liquid-liquid extraction	4
1.1.4 Step of solid phase extraction procedure	5
1.1.4.1 Pretreatment of the samples	5
1.1.4.2 Conditioning of the cartridges	5
1.1.4.3 Sample loading step	6
1.1.4.4 Washing step	6
1.1.4.5 Eluting step	6
1.2 Molecularly imprinted polymers (MIPs)	8
1.2.1 Theory of MIPs	8
1.2.2 Molecular imprinting procedure	9

1.2.2.1 Covalent imprinting	10
1.2.2.2 Non-covalent imprinting	11
1.2.3 Constraints of MIPs synthesis	13
1.2.3.1 Template	13
1.2.3.2 Functional monomer	14
1.2.3.3 Cross-linker	14
1.2.3.4 Initiators	17
1.2.3.5 Solvents (Prorogens)	18
1.3 Molecularly imprinted solid-phase extraction (MISPE)	19
1.4 Nevirapine	23
1.5 Aim of this research	25
 <b>CHAPTER II: SYNTHESIS AND EVALUATION OF MOLECULARLY IMPRINTED POLYMERS FOR THE BINDING AFFINITY TO NVP</b>	
2.1 Introduction	26
2.2 Experimentals	28
2.2.1 Chemicals	28
2.2.2 Instrumentals	29
2.2.3 Synthesis of MIPs	29
2.2.4 Equilibrium binding study of MIPs with the corresponding templates	32
2.2.5 Equilibrium binding study of MIPs with NVP	33
2.2.6 Competitive binding study	33

2.2.7 Chromatographic conditions	34
2.3 Results and discussions	35
2.3.1 Synthesis of MIPs	35
2.3.2 Equilibrium binding study of MIPs with the corresponding templates	36
2.3.3 Equilibrium binding study of MIPs with NVP	40
2.3.4 The effect of amount of polymer to the binding efficiency of NAM to P(NAM)	43
2.3.5 The competitive binding study	45
<b>CHAPTER III: MOLECULARLY IMPRINTED SOLID-PHASE EXTRACTION FOR NVP</b>	
3.1 Introduction	47
3.2 Experimentals	48
3.2.1 Chemicals	48
3.2.2 Instrumentals	49
3.2.3 General procedure for MISPE	49
3.2.4 Optimization of the extraction conditions	50
3.2.4.1 Optimization of the washing condition	50
3.2.4.2 Optimization of the eluting condition	52
3.2.5 MISPE with the real plasma sample	53
3.2.6 Effect of NAM in MISPE process	53
3.2.7 Method validation	53


3.2.8 Plasma sample preparation	54
3.2.9 Chromatographic conditions	54
3.3 Results and discussions	55
3.3.1 Optimization of the washing condition	55
3.3.2 Optimization of the eluting condition	57
3.3.3 MISPE with the real plasma sample	58
3.3.4 Effect of NAM to recovery of NVP	60
3.3.5 Method Validation	61
<b>CHAPTER IV: CONCLUSIONS</b>	62
<b>REFERENCES</b>	66
<b>APPENDICES</b>	73
<b>APPENDIX A</b>	74
<b>APPENDIX B</b>	88
<b>CURRICULUM VIVATE</b>	97

## LIST OF TABLES

Table	Page
1.1 Advantages and disadvantages of LLE and SPE technique	4
1.2 Advantages and disadvantages of covalent and non-covalent imprinting	12
1.3 Various application of MIP in SPE	20
2.1 Conditions of MIPs preparation	30
2.2 Chromatographic condition for analysis	33
2.3 The NVP selectivity factors of all polymers in 0.2 mM NVP in 0.01 M phosphate buffer pH 7 containing 0.05% Tween 20	40
2.4 The imprinting factors of P(NAM) in 0.2 mM NVP in 0.01 M phosphate buffer pH 7 containing 0.05% Tween 20	44
3.1 The washing conditions	50
3.2 The eluting conditions	51
3.3 Chromatographic conditions for analysis	54
A.1 Calibration curve of NAM in 0.01 M phosphate buffer pH 7 containing 0.05% Tween 20	73
A.2 Calibration curve of NVP in 0.01 M phosphate buffer pH 7 containing 0.05% Tween 20	74
A.3 Calibration curve of BZM in 0.01 M phosphate buffer pH 7 containing 0.05% Tween 20	75
A.4 Calibration curve of BZP in 0.01 M phosphate buffer pH 7 containing 0.05% Tween 20	76

<b>Table</b>	<b>Page</b>
A.5 The binding study of 0.2 mM NVP in 0.01 M phosphate buffer pH 7 containing 0.05% Tween 20 with P(NVP)	77
A.6 The binding study of 0.2 mM NAM in 0.01 M phosphate buffer pH 7 containing 0.05% Tween 20 with 5 mg of P(NAM)	77
A.7 The binding study of 0.2 mM BZM in 0.01 M phosphate buffer pH 7 containing 0.05% Tween 20 with 5 mg of P(BZM)	78
A.8 The binding study of 0.2 mM BZP in 0.01 M phosphate buffer pH 7 containing 0.05% Tween 20 with 5 mg of P(BZP)	78
A.9 The binding study of 0.2 mM NVP in 0.01 M phosphate buffer pH 7 containing 0.05% Tween 20 with P(NVP), P(NAM), P(BZM) and P(BZP)	79
A.10 The binding study of 0.2 mM NAM in 0.01 M phosphate buffer pH 7 containing 0.05% Tween 20 with P(NAM)	80
A.11 The competitive binding study of mix solution of NAM and NVP in 0.01 M phosphate buffer pH 7 containing 0.05% Tween 20 with P(NAM)	80
A.12 Calibration curve of NVP in plasma sample	82
A.13 Percentage of NVP retained in MISPE cartridge after washing step in various washing conditions	83
A.14 Percentage of NVP recovery after MISPE process in various eluting conditions	84
A.15 The effect of NAM to recovery of NVP in plasma sample after MISPE process.	85

A.16 The precision study	85
A.17 The accuracy of the recovery of 10 µg/ml NVP in plasma sample after MISPE process	86



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่  
 Copyright© by Chiang Mai University  
 All rights reserved

## LIST OF FIGURES

Figure	Page
1.1 The general SPE procedure	7
1.2 The schematic of MIP preparation	9
1.3 The schematics of MIP preparation; covalent and non-covalent approach	12
1.4 The functional monomers most commonly used in MIPs	15
1.5 The cross-linker most commonly used in MIPs	16
1.6 The initiator most commonly used in MIPs	17
1.7 The structure of Nevirapine	23
2.1 Structure of templates, functional monomer, cross-linker and initiator used in MIP synthesis	29
2.2 The imprinting factor and the percentage bound of MIP with corresponding template	37
2.3 Schematic representation of the synthesis of P(NVP)	38
2.4 The percentage bound of P(NVP), P(NAM), P(BZM) and P(BZP) to NVP (0.2 mM) in 0.01 M phosphate buffer pH 7 containing 0.05% Tween 20	40
2.5 The schematic presentation the P (NAM) synthesis	41
2.6 The effect of amount of P(NAM) to the binding affinity with corresponding template	43
2.7 The competitive binding study of P(NAM) with mix solution of NAM:NVP in 0.01M phosphate buffer pH 7 containing 0.05% Tween 20 in ratio 0:1, 1:1, 5:1 and 10:1	45
3.1 MISPE method	49



<b>Figure</b>	<b>Page</b>
3.2 The percentage of retained NVP after washing step in various washing conditions	56
3.3 The percentage of recovery NVP after eluting step in various eluting conditions	57
3.4 HPLC chromatogram of MISPE for NVP in plasma sample	58
3.5 Effect of NAM for recovery of NVP	59
A.1 Calibration curve of NAM in 0.01 M phosphate buffer pH 7 containing 0.05% Tween 20	73
A.2 Calibration curve of NAM in 0.01 M phosphate buffer pH 7 containing 0.05% Tween 20	74
A.3 Calibration curve of BZP in 0.01 M phosphate buffer pH 7 containing 0.05% Tween 20	75
A.5 Calibration curve of BZP in 0.01 M phosphate buffer pH 7 containing 0.05% Tween 20	76
A.4 Calibration curve of NVP in plasma sample	82
B.1 The HPLC chromatogram of competitive binding study of P(NAM) in mix solution of NAM:NVP (1:1) in 0.01 M phosphate buffer pH7 containing 0.05% Tween 20	87
B.2 The HPLC chromatogram of competitive binding study of P(NAM) in mix solution of NAM:NVP (5:1) in 0.01 M phosphate buffer pH7 containing 0.05% Tween 20	88

<b>Figure</b>	<b>Page</b>
B.3 The HPLC chromatogram of competitive binding study of P(NAM) in mix solution of NAM:NVP (10:1) in 0.01 M phosphate buffer pH7 containing 0.05% Tween 20	89
B.4 The HPLC chromatogram of the study of MISPE with mix solution of NAM:NVP (6:1) in 0.01 M phosphate buffer pH7 containing 0.05% Tween 20	90
B.5 The HPLC chromatogram of the study of MISPE with mix solution of NAM:NVP (1:1) in 0.01 M phosphate buffer pH7 containing 0.05% Tween 20	91
B.6 The HPLC chromatogram of the study of MISPE with mix solution of NAM:NVP (12:1) in 0.01 M phosphate buffer pH7 containing 0.05% Tween 20	92
B.7 The HPLC chromatograms of 0.2 mM NVP in 0.01 M phosphate buffer pH 7 containing 0.05% Tween 20 after washing step	93
B.8 The HPLC chromatograms of 0.2 mM NVP in 0.01 M phosphate buffer pH 7 containing 0.05% Tween 20 after eluting step	95

## ABBREVIATIONS AND SYMBOLS

<b>MISPE</b>	Molecularly imprinted solid-phase extraction
<b>NVP</b>	Nevirapine
<b>MIPs</b>	Molecularly imprinted polymers
<b>NAM</b>	Nicotinamide
<b>SPE</b>	Solid-phase extraction
<b>HPLC</b>	High-performance liquid chromatography
<b>GC</b>	Gas chromatography
<b>LC-MS-MS</b>	Liquid chromatography tandem mass spectrometry
<b>RIA</b>	Radio immunoassay
<b>LLE</b>	Liquid-liquid extraction
<b>UV</b>	Ultraviolet
<b>ELISA</b>	Enzyme-link immunoassay
<b>NNRTI</b>	Non-nucleoside reverses transcriptase inhibitor
<b>BZM</b>	Benzamide
<b>BZP</b>	Benzophenone
<b>NIP</b>	Non-imprinted polymer
<b>TRIM</b>	Trimethylopropane trimethacrylate
<b>MAA</b>	Methacrylic acid
<b>MeOH</b>	Methanol
<b>THF</b>	Tetrahydrofuran
<b>TEA</b>	Triethylamine

<b>SD</b>	Standard deviation
<b>%RA</b>	Percentage of relative accuracy
<b>%RSD</b>	Percentage of relative standard deviation
<b>M</b>	Molar
<b>mM</b>	Millimolar
<b>ml</b>	Milliliter
<b>mg</b>	Milligram
<b>nm</b>	Nanometer
<b>mg/ml</b>	Milligram per milliliter
<b>µg/ml</b>	Microgram per milliliter
<b>ml/min</b>	Milliliter per minute
<b>v/v</b>	Volumn by volumn
<b>λ</b>	Wavelength
<b>λ<sub>max</sub></b>	The wavelength of maximum absorbance
<b>α</b>	The imprinting factor
<b>α'</b>	The NVP selectivity factor
<b>mAU</b>	milli area unit