

REFERENCES

- Abbott, T.P., Nabetani, H., Sessa, D.J., Wolf, W.J., Liebman, M.N., and Dukor, R.K. (1996). Effect of bound water on FTIR spectra of glycinin. *Journal of Agricultural and Food Chemistry*, 44(8), 2220-2224.
- Anderson, R.L., and Wolf, J.W. (1995). Compositional changes in trypsin inhibitors, phytic acid, saponins and isoflavones related to soybean processing. *Journal of Nutrition*, 125(3), 581-588.
- AOAC (2000). *Official methods of analysis of AOAC international*, 17th ed. Gaithersburg, MD, USA.
- Apichartsrangkoon, A. (2002). Dynamic viscoelastic properties of heated gluten/soy protein gels. *Journal of Food Science*, 67(2), 637-657.
- Apichartsrangkoon, A. (2003). Effects of high pressure on rheological properties of soy protein gels. *Food Chemistry*, 80(1), 55-60.
- Apichartsrangkoon, A., and Ledward, D.A. (2002). Dynamic viscoelastic behaviour of high pressure treated gluten-soy mixtures. *Food Chemistry*, 77(3), 317-323.
- Auty, M.A.E., Twomey, M., Guinee, T.P., and Mulvihill, D.H. (2001). Development and application of confocal scanning laser microscopy methods for studying the distribution of fat and protein in selected dairy products. *Journal of Dairy Research*, 68(3), 417-427.
- Badley, R.A., Atkinson, D., Hauser, H., Oldani, D., Green, J.P., and Stubbs, J.M. (1975). The structure, physical and chemical properties of the soy bean protein glycinin. *Biochimica et Biophysica Acta*, 412(2), 214-228.
- Balci, T., and Wilbey, A. (1999). High pressure processing of milk-the first 100 years in the development of a new technology. *International Journal of Dairy Technology*, 52(4), 149-155.
- Balny, C., Mozhaev, V.V., and Lange, R. (1997). Hydrostatic pressure and proteins: Basic concepts and new data. *Comparative Biochemistry and Physiology Part A: Physiology*, 116A(4), 299-304.

- Beddows, C.G., and Wong, J. (1987). b. Optimization of yield and properties of silken tofu from soybeans. II. Heat processing. *International Journal of Food Science and Technology*, 22, 23-27.
- Bell, A.E. (1989). Gel Structure and Food Biopolymers. in T.M. Hardman (Ed.), *Water and Food Quality* (pp.251-275), London, UK: Elsevier Applied Science.
- Belyakova, L.E., Semenova, M.G., and Antipova, A.S. (1999). Effect of small molecule surfactants on molecular parameters and thermodynamic properties of legumin in a bulk and at the air-water interface depending on a protein structure in an aqueous medium. *Colloids and Surfaces B: Biointerfaces*, 12(3), 271-285.
- Berk, Z. (1992). Technology of production of edible flours and protein products from soybean. *FAO Agricultural service bulletin*, 97, 127-134.
- Birk, Y. (1985). The Bowman-Birk inhibitor. Trypsin- and chymotrypsin-inhibitor from soybeans. *International Journal of Peptide Protein Research*, 25(2), 113-131.
- Boonyaratankornkit, B.B., Park, C.B., and Clark, D.S. (2002). Pressure effects on intra- and intermolecular interactions within proteins. *Biochimica et Biophysica Acta*, 1595(1-2), 235-249.
- Cai, T.D., and Chang, K.C. (1998). Characteristics of production-scale tofu as affected by soymilk coagulation method: propeller blade size, mixing time and coagulant concentration. *Food Research International*, 31(4), 289-295.
- Cai, T.D., Chang, K.C., Shih, M.C., Hou, H.J., and Ji, M. (1997). Comparison of bench and production scale methods for making soymilk and tofu from 13 soybean varieties. *Food Research International*, 30(9), 659-668.
- Chang, S.K.C., and Yuan, S. (2005). Ultrastructural properties of various types of tofu. IFT Annual Meeting, New Orleans, Louisiana.
- Cheng, Y., Shimizu, N., and Kimura, T. (2005). The viscoelastic properties of soybean curd (tofu) as affected by soymilk concentration and type of coagulant. *International Journal of Food Science and Technology*, 40(4), 385-390.

- Clark, A.H., Kavanagh, G.M., and Ross-Murphy, S.B. (2001). Globular protein gelation-theory and experiment. *Food Hydrocolloids*, 15(4-6), 383-400.
- Damodaran, S. (1996). Amino Acid, Peptides, and Proteins. in O.R. Fennema (Ed.), *Food Chemistry*, 3rd ed. (pp.321-429), New York: Marcel Dekker Inc.
- deMan, J.M., deMan, L., and Gupta, S. (1986). Texture and microstructure of soybean curd (tofu) as affected by different coagulants. *Food Microstructure*, 5, 83-89.
- DiPietro, C.M., and Liener, I.E. (1989). Heat interaction of the Kuniz and Bowman-Birk soybean protease inhibitors. *Journal of Agricultural and Food Chemistry*, 37(1), 39-44.
- Dumay, E.M., Kalichevsky, M.T., and Cheftel, J.C. (1998). Characteristics of pressure-induced gels of β -lactoglobulin at various times after pressure release. *LWT-Food Science and Technology*, 31(1), 10-19.
- Dumoulin, M., Ozawa, S., and Hayashi, R. (1998). Textural properties of pressure-induced gels of food proteins obtained under different temperatures including subzero. *Journal of Food Science*, 63(1), 92-95.
- Dunn, M.J. (1993). *Gel Electrophoresis: Proteins*. Harefield, UK: Bios Scientific Publishers.
- Emerton, V. (2003). *Essential Guide to Food Additives*, 2nd ed. Surrey, UK: Leatherhead Publishing.
- Estrada-Giron, Y., Swanson, B.G., and Barbosa-Canovas, G.V. (2005). Advances in the use of high hydrostatic pressure for processing cereal grains and legumes. *Trends in Food Science and Technology*, 16(5), 194-203.
- Ferry, J.D. (1980). *Viscoelastic properties of polymers*, 3rd ed. New York: John Wiley & Sons.
- Fukushima, D. (1981). Soy proteins for foods centering around soy sauce and tofu. *Journal of the American Oil Chemists' Society*, 58(3), 346-354.
- Fukushima, D. (1991). Recent process of soybean protein foods: Chemistry, Technology, and Nutrition. *Food Research International*, 7(3), 323-351.
- Galazka, V.B., Dickinson, E., and Ledward, D.A. (2000). Influence of high pressure processing on protein solutions and emulsions. *Current Opinion in Colloid and Interface Science*, 5(3-4), 182-187.

- Gaucheron, F., Famelart, M.H., Mariette, F., Raulot, K., Michel, F., and Le Graet, Y. (1997). Combined effects of temperature and high-pressure treatments on physicochemical characteristics of skim milk. *Food Chemistry*, 59(3), 439-447.
- German, B., Damodaran, S., and Kinsella, J.E. (1982). Thermal dissociation and association behavior of soy proteins. *Journal of Agricultural and Food Chemistry*, 30(5), 807-811.
- Gomes, M.R.A. (1997). *Effect of high pressure treatment on polyphenoloxidases, papain and amylases*. Ph.D. thesis, Department of Food Science and Technology, The University of Reading, Reading, UK.
- Gould, G.W. (2001). The evolution of high pressure processing of foods. in M.E.G. Hendrickx, and D. Knorr (Eds.). *Ultra High Pressure Treatments of Foods* (pp.3-21). New York: Kluwer Academic/Plenum Publishers.
- Gross, M., and Jaenicke, R. (1994). Proteins under pressure. The influence of high hydrostatic pressure on structure, function and assembly of proteins and protein complexes. *European Journal of Biochemistry*, 221(2), 617-630.
- Guo, S.-T., and Ono, T. (2005). The role of composition and content of protein particles in soymilk on tofu curdling glucono- δ -lactone or calcium sulfate. *Journal of Food Science*, 70(4), C258-262.
- Hackler, L.R., Van Buren, J.P., Streinkraus, K.H., El Rawi, I., and Hand, D.B. (1965). Effect of heat treatment on nutritive value of soymilk protein fed to weanling rats. *Journal of Food Science*, 30(4), 723-728.
- Hamerstrand, G.E., Black, L.T., and Glover, J.D. (1981). Trypsin inhibitors in soy products: Modification of the standard analytical procedure. *Cereal Chemistry*, 58(1), 42-45.
- Hayakawa, S., and Nakai, S. (1985). Relationships of hydrophobicity and net charge to the solubility of milk and soy proteins. *Journal of Food Science*, 50(2), 486-491.
- Hendrickx, M., Ludikhuyze, L., Van den Broeck, I., and Weemaes, C. (1998). Effects of high pressure on enzymes related to food quality. *Trends in Food Science and Technology*, 9(5), 197-203.

- Heremans, K. (1982). High pressure effects on proteins and other biomolecules. *Annual review of biophysics and bioengineering*, 11, 1-21.
- Heremans, K. (1995). High Pressure effects on Biomolecules. in D.A. Ledward, D.E. Johnston, R.G. Earnshaw, and A.P.M. Hasting (Eds.). *High Pressure Processing of Foods* (pp.81-97). Reading: Nottingham University Press.
- Heremans, K. (2001). The effects of high pressure on biomaterials. in M.E.G. Hendrickx, and D. Knorr (Eds.). *Ultra High Pressure Treatments of Foods* (pp.23-46). New York, USA: Kluwer Academic/Plenum Publishers.
- Heremans, K., and Smeller, L. (1998). Protein structure and dynamics at high pressure. *Biochimica et Biophysica Acta*, 1386(2), 353-370.
- Hermansson, A.M. (1978). Physico-chemical aspects of soy proteins structure formation. *Journal of Texture Studies*, 9, 33-58.
- Hermansson, A.-M. (1986). Soy protein gelation. *Journal of the American Oil Chemists' Society*, 63(5), 658-666.
- Hermansson, A.-M. (1994). Microstructure of protein gels related to functionality. in R.Y. Yada, R.L. Jackman, and J.L. Smith (Eds.). *Protein Structure-Function Relationships in Foods* (pp.22-42). Glasgow, UK: Black Academic & Professional.
- Huppertz, T., Kelly, A.L., and Fox, P.F. (2002). Effects of high pressure on constituents and properties of milk. *International Dairy Journal*, 12(7), 561-572.
- Kajiyama, N., Isobe, S., Uemura, K., and Noguchi, A. (1995). Change of soy protein under ultra-high hydraulic pressure. *International Journal of Food Science and Technology*, 30(2), 147-158.
- Kakade, M.L., Rackis, J.J., McGhee, J.E., and Puski, G. (1974). Determination of trypsin inhibitor activity of soy products: A collaborative analysis of an improved procedure. *Cereal Chemistry*, 51, 376-382.
- Kakade, M.L., Simon, N.R., and Liener, I.E. (1969). An evaluation of natural vs. synthetic substrates for measuring the antitryptic activity of soybean samples. *Cereal Chemistry*, 46, 518-526.

- Kao, F.-J., Su, N.-W., and Lee, M.-H. (2003). Effect of calcium sulfate concentration in soymilk on the microstructure of firm tofu and protein constitutions in tofu whey. *Journal of Agricultural and Food Chemistry*, 51(21), 6211-6216.
- Kao, F.-J., Su, N.-W., and Lee, M.-H. (2004). Effect of water-to-bean ratio on the contents and compositions of isoflavones in tofu. *Journal of Agricultural and Food Chemistry*, 52(8), 2277-2281.
- Kato, A., Osako, Y., Matsudomi, N., and Kobayashi, K. (1983). Changes in the emulsifying and foaming properties of proteins during heat denaturation. *Agricultural and Biological Chemistry*, 47(1), 33-37.
- Kennedy, A.R. (1994). Prevention of carcinogenesis by protease inhibitors. *Cancer Research*, 54, 1999s-2005s.
- Kennedy, A.R. (1998). Chemopreventive agents: Protease inhibitors. *Pharmacology and Therapeutics*, 78(3), 167-209.
- Ker, Y.-C., Chen, R.-H., and Wu, C.-S. (1993). Relationships of secondary structure, microstructure, and mechanical properties of heat-induced gel of soy 11S globulin. *Bioscience, Biotechnology Biochemistry*, 57(4), 536-541.
- Kinsella, J.E. (1979). Functional properties of soy proteins. *Journal of the American Oil Chemists' Society*, 56, 242-258.
- Kinsella, J.E., Rector, D.J., and Philips, L.G. (1994). Physicochemical properties of proteins: Texturization via gelation, glass and film formation. in R.Y. Yada, R.L. Jackman, and J.L. Smith (Eds.), *Protein Structure-Function Relationships in Foods* (pp.1-21). Glasgow, UK: Blackie Academic & Professional.
- Kitamura, K., Takagai, T., and Shibasaki, K. (1976). Subunit structure of soybean 11S globulin. *Agricultural Biological Chemistry*, 40(9), 1837-1844.
- Knorr, D., Heinz, V., and Buckow, R. (2006). High pressure application for food biopolymers. *Biochimica et Biophysica Acta*, 1764(3), 619-631.
- Kohyama, K., Yoshida, M., and Nishinari, K. (1992). Rheological study on gelation of soybean 11S protein by glucono- δ -lactone. *Journal of Agricultural and Food Chemistry*, 40(5), 740-744.

- Kohyama, K., and Nishinari, K. (1993). Rheological study on the gelation process of soybean 7S and 11S protein in the presence of Glucono- δ -lactone. *Journal of Agricultural and Food Chemistry*, 41(1), 8-14.
- Kohyama, K., Sano, Y., and Doi, E. (1995). Rheological characteristics and gelation mechanism of tofu (soybean curd). *Journal of Agricultural and Food Chemistry*, 43(7), 1808-1812.
- Koshiyama, L. (1971). Some aspects of subunit structure of 7S protein in soybean globulin. *Agricultural and Biological Chemistry*, 35, 385-392.
- Kwok, K.-C., Liang, H.-H., and Niranjana, K. (2002). Optimizing conditions for thermal processes of soy milk. *Journal of Agricultural and Food Chemistry*, 50(17), 4834 - 4838.
- Laemmli, U.K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, 227, 680-685.
- Law, A.M., and Kelton, W.D. (2000). *Simulation Modeling and Analysis*. Singapore: McGraw-Hill Higher Education.
- Ledward, D.A. (1995). High pressure processing-The potential. in D.A. Ledward, D.E. Johnston, R.G. Earnshaw, and A.P.M. Hasting (Eds.), *High Pressure Processing of Foods* (pp.1-7), UK: Nottingham University Press.
- Liener, I.E. (1981). Factors affecting the nutritional quality of soya products. *Journal American Oil Chemists' Society*, 58(3), 406-415.
- Liu, K.S. (1999). *Soybeans chemistry, technology and utilization*. Gaithersburg, Maryland: Aspen Publishers.
- Liu, Z.-S., and Chang, K.C. (2003). Development of rapid titration method for predicting optimal coagulant concentration for filled tofu. *Journal of Agricultural and Food Chemistry*, 51(18), 5214-5221.
- Liu, Z.-S., and Chang, S.K.-C. (2004). Effect of soy milk characteristics and cooking conditions on coagulant requirements for making filled tofu. *Journal of Agricultural and Food Chemistry*, 52(11), 3405-3411.
- Lopez-Fandino, R. (2006). High pressure-induced changes in milk proteins and possible applications in dairy technology. *International Dairy Journal*, 16(10), 1119-1131.

- Lullien-Pellerin, V., and Balny, C. (2002). High-pressure as a tool to study some proteins' properties: conformational modification, activity and oligomeric dissociation. *Innovative Food Science and Emerging Technologies*, 3(3), 209-221.
- Maltais, A., Remondetto, G.E., Gonzalez, R., and Subirade, M. (2005). Formation of soy protein isolate cold-set gels: Protein and salt effects. *Journal of Food Science*, 70(1), C67-73.
- Matser, A.M., Krebbers, B., van den Berg, R.W., Bartels, P.V. (2004). Advantages of high pressure sterilisation on quality of food products. *Trends in Food Science and Technology*, 15, 79-85.
- Matsuura, M. (1985). Process for producing an aseptic packed tofu, *U.S. Patent 4514433* [Online]. Available: <http://www.freepatentsonline.com/4514433.html> [2007, 18 June].
- Matsuura, M., Sasaki, M., Sasaki, J., and Takeuchi, T. (2000). Process for producing packed tofu, *U.S. Patent 6042851* [Online]. Available: <http://www.patentstorm.us/patents/6042851.html> [2007, 19 June].
- Messens, W., Van Camp, J., and Huyghebaert, A. (1997). The use of high pressure to modify the functionality of food proteins. *Trends in Food Science and Technology*, 8(4), 107-112.
- Messina, M.J. (1999). Soyfoods: Their Role in Disease Prevention and Treatment. in K.S. Liu, *Soybeans Chemistry, Technology, and Utilization*, (pp.442-476), Gaithersburg, Maryland: Aspen Publishers, Inc.
- Min, S., Yu, Y., and St. Martin, S. (2005). Effect of soybean varieties and growing locations on the physical and chemical properties of soymilk and tofu. *Journal of Food Science*, 70(1), C8-12.
- Molina, E., and Ledward, D.A. (2003). Effects of combined high-pressure and heat treatment on the textural properties of soya gels. *Food Chemistry*, 80(3), 367-370.
- Molina, E., Defaye, A.B., and Ledward, D.A. (2002). Soy protein pressure-induced gels. *Food Hydrocolloids*, 16(6), 625-632.


- Molina, E., Papadopoulou, A., and Ledward, D.A. (2001). Emulsifying properties of high pressure treated soy protein isolate and 7S and 11S globulins. *Food Hydrocolloids*, 15(3), 263-269.
- Moure, A., Sineiro, J., Dominguez, H. and Parajo, J.C. (2006). Functionality of oilseed protein products: A review. *Food Research International*, 39(9), 945-963.
- Murphy, P.A., and Resurrection, A.P. (1984). Varietal and environmental differences in soybean glycinin and β -conglycinin content. *Journal of Agricultural and Food Chemistry*, 32(4), 911-915.
- Neuhoff, V., Arold, N., Taube, D., and Ehrhardt, W. (1988). Improved staining of proteins in polyacrylamide gels including isoelectric focusing gels with clear background at nanogram sensitivity using Coomassie Brilliant Blue G-250 and R-250. *Electrophoresis*, 9, 255-262.
- Oakenfull, D., Pearce, J., and Burley, R.W. (1997). Protein gelation. in S. Damodaran and A. Paraf (Eds.), *Food Proteins and Their Applications* (pp.110-142), New York: Marcel Dekker, Inc.
- Obata, A., and Matsuura, M. (1993). Decrease in the gel strength of tofu caused by an enzyme reaction during soybean grinding and its control. *Bioscience, Biotechnology Biochemistry*, 57(4), 542-545.
- Okamoto, M., Kawamura, Y., and Hayashi, R. (1990). Application of high pressure to food processing: Textural comparison of pressure- and heat-induced gels of food proteins. *Agricultural and Biological Chemistry*, 54(1), 183-189.
- Ono, T., Katho, S., and Mothizuki, K. (1993). Influences of calcium and pH on protein solubility in soybean milk. *Bioscience, Biotechnology and Biochemistry*, 57(1), 24-28.
- Pandey, P.K., Ramaswamy, H.S., and St-Gelais, D. (2000). Water-holding capacity and gel strength of rennet curd as affected by high-pressure treatment of milk. *Food Research International*, 33(8), 655-663.
- Pares, D., Sagner, E., Saurina, J., Sunol, J.J., and Carretero, C. (1998). Functional Properties of heat induced gels from liquid and spray-dried porcine blood plasma as influenced by pH. *Journal Food Science*, 63(6), 958-961.

- Pedrosa, C., and Ferreira, S.T. (1994). Deterministic pressure-induced dissociation of vicilin, the 7S storage globulin from pea seeds: Effects of pH and cosolvents on oligomer stability. *Biochemistry*, 33(13), 4046-4055.
- Prestamo, G., and Fontecha, J. (2007). High pressure treatment on the tofu fatty acids and acylglycerols content. *Innovative Food Science and Emerging Technologies*, 8(2), 188-191.
- Prestamo, G., Lasuncion, M.A., and Arroyo, G. (2002). Response of rat to the intake of tofu treated under high-pressure. *Innovative Food Science and Emerging Technologies*, 3(2), 149-155.
- Prinyawiwatkul, W., Eitenmiller, R.R., Beuchat, L.R., McWatters, K.H., and Philips, R.D. (1996). Cowpea flour vitamins and trypsin inhibitor affected by treatment and fermentation with *Rhizopus microsporus*. *Journal of Food Science*, 61(5), 1039-1042.
- Puppo, C., Chapleau, N., Speroni, F., de Lamballerie-Anton, M., Michel, F., Anon, C., and Anton, M. (2004). Physicochemical modifications of high-pressure-treated soybean protein isolates. *Journal of Agricultural and Food Chemistry*, 52(6), 1564 -1571.
- Puppo, M.C., and Anon, M.C. (1998). Structure properties of heat-induced soy protein gels as affected by ionic strength and pH. *Journal of Agricultural and Food Chemistry*, 46, 3583-3589.
- Rao, M.A. (1999). *Rheology of Fluid and Semisolid Foods*. Gaithersburg, Maryland : Aspen Publishers.
- Renkema, J.M.S., Knabben, J.H.M., and van Vliet, T. (2001). Gel formation by β -conglycinin and glycinin and their mixtures. *Food Hydrocolloids*, 15(4-6), 407-414.
- Riblett, A.L., Herald, T.J., Schmidt, K.A., and Tilley, K.A. (2001). Characterization of β -Conglycinin and glycinin soy protein fractions from four selected soybean genotypes. *Journal of Agricultural and Food Chemistry*, 49(10), 4983-4989.
- Ruegg, M., Moor, U., and Blanc, B. (1975). Hydration and thermal denaturation of β -lactoglobulin. A calorimetric study. *Biochimica et Biophysica Acta*, 400(2), 334-342.

- Saio, K. (1979). Tofu-relationships between texture and fine structure. *Cereal Foods World*, 24(8), 342-345, 352-354.
- Saio, K., Kajikawa, M., and Watanabe, T. (1971). Food processing characteristics of soybean proteins part II. Effect of sulfhydryl groups on physical properties of tofu-gel. *Agricultural and Biological Chemistry*, 35(6), 890-898.
- Schwertfeger, M., and Buchheim, W. (1999). Coagulation of skim milk under high hydrostatic pressure with acidification by glucono- δ -lactone. *International Dairy Journal*, 9(7), 487-492.
- Shen, C.F., De Man, L., Buzzell, R.I., and De Man, J.M. (1991). Yield and quality of tofu as affected by soybean and soymilk characteristics: glucono-delta-lactone coagulant. *Journal of Food Science*, 56(1), 109-112.
- Shih, M.C., Hou, H.J., and Chang, K.C. (1997). Process optimization for soft tofu. *Journal of Food Science*, 62(4), 833-837.
- Shun-Tang, G., Ono, T., and Mikami, M. (1999). Incorporation of soymilk lipid into protein coagulum by addition of calcium chloride. *Journal of Agricultural and Food Chemistry*, 47(3), 901-905.
- Sorgentini, D.A., Wagner, J.R., and Anon, M.C. (1995). Effects of thermal treatment of soy protein isolate on the characteristics and structure-function relationship of soluble and insoluble fractions. *Journal of Agricultural and Food Chemistry*, 43(9), 2471-2479.
- Steffe, J.F. (1996). *Rheological methods in food process engineering*, 2nd ed. East Lansing, USA: Freeman Press.
- Sun, N., and Breene, W.M. (1991). Calcium sulfate concentration influence on yield and quality of tofu from five soybean varieties. *Journal of Food Science*, 56(6), 1604-1607, 1610.
- Tay, S.L., and Perera, C.O. (2004). Physicochemical properties of 7S and 11S protein mixtures coagulated by glucono-delta-lactone. *Journal of Food Science*, 69(4), E139-43.
- Tay, S.L., Xu, G.Q., and Perera, C.O. (2005). Aggregation profile of 11S, 7S and 2S coagulated with GDL. *Food Chemistry*, 91(3), 457-462.

- Torrezan, R., Tham, W.P., Bell, A.E., Frazier, R.A., and Cristianini, M. (2007). Effects of high pressure on functional properties of soy protein. *Food Chemistry*, 104(1), 140-147.
- Tsai, S.-J., Lan, C.Y., Kao, C.S., and Chen, S.C. (1981). Studies on the yield and quality characteristics of tofu. *Journal of Food Science*, 46, 1734-1737.
- U.S. FDA. (1999). FDA approves new health claim for soy protein and coronary heart disease, In *FDA Talk Paper T99-48* [Online]. Available: <http://www.fda.gov/bbs/topics/ANSWERS/ANS00980.html> [2007, 26 June].
- Utsumi, S., and Kinsella, J.E. (1985). Forces involved in soy protein gelation: Effects of various reagents on the formation, hardness and solubility of heat-Induced gels made from 7S, 11S, and soy isolate. *Journal of Food Science*, 50(5), 1278-1282.
- van der Ven, C., Matser, A.M., and van den Berg, R.W. (2005). Inactivation of soybean trypsin inhibitors and lipoxygenase by high-pressure processing. *Journal of Agricultural and Food Chemistry*, 53(4), 1087-1092.
- Walstra, P., Geurts, T.J., Noomen, A., Jellema, A., and van Boekel, M.A.J.S. (1999). *Dairy Technology*. New York: Marcel Dekker, Inc.
- Weaver, C.M., Heaney, R.P., Connor, L., Martin, B.R., Smith, D.L., and Nielson, S. (2002). Bioavailability of calcium from tofu as compared with milk in premenopausal woman. *Journal of Food Science*, 67(8), 3144-3147.
- Windholz, M., Budavari, S., Blumetti, R.F., Otterbein, E.S. (Eds.) (1983). *The Merck index: an encyclopedia of chemicals, drugs, and biologicals*, 10th ed. Rahway, N.J., U.S.A.: Merck & Co., Inc.
- Wu, W.U., Hettianachy, N.S., and Qi, M. (1998). Hydrophobicity, solubility, and emulsifying properties of soy protein peptides prepared by papain modification and ultrafiltration. *Journal of the American Oil Chemists' Society*, 75(7), 845-850.
- Wu, Y.V., and Sessa, D.J. (1994). Conformation of Bowman-Birk inhibitor. *Journal of Agricultural and Food Chemistry*, 42(10), 2136-2138.
- Yamauchi, F., Yamagishi, T., and Iwabuchi, S. (1991). Molecular understanding of heat-induced phenomena of soybean protein. *Food Reviews International*, 7(3), 283-322.

- Zarkadas, C.G., and Yu, Z. (1993). Assessment of the protein quality of a new high-protein soybean cultivar by amino acid analysis. *Journal of Agricultural and Food Chemistry*, 41(4), 616-623.
- Zhang, H., Li, L., Tatsumi, E., and Isobe, S. (2005). High-pressure treatment effects on proteins in soy milk. *LWT-Food Science and Technology*, 38(1), 7-14.
- Zhang, H., Li, L., Tatsumi, E., and Kotwal, S. (2003). Influence of high pressure on conformational changes of soybean glycinin. *Innovative Food Science and Emerging Technologies*, 4(3), 269-275.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved