

## REFERENCES

- Adhikari, S., Joshi, R., & Gopinathan, C. (1998). Bilirubin as an anti precipitant against copper mediated denaturation of bovine serum albumin: formation of copper-bilirubin complex. *Biochim Biophys Acta*, 1380(1), 109-114.
- Akin, E., Clower, B., Tibbs, R., Tang, J., & Zhang, J. (2002). Bilirubin produces apoptosis in cultured bovine brain endothelial cells. *Brain Res*, 931(2), 168-175.
- Ahlfors, C. E., & Wennberg, R. P. (2004). Bilirubin-albumin binding and neonatal jaundice. *Semin Perinatol*, 28(5), 334-339.
- Amit, Y., & Brenner, T. (1993). Age-dependent sensitivity of cultured rat glial cells to bilirubin toxicity. *Exp Neurol*, 121(2), 248-255.
- Armstrong, C., Leong, W., & Lees, G. J. (2001). Comparative effects of metal chelating agents on the neuronal cytotoxicity induced by copper (Cu<sup>+2</sup>), iron (Fe<sup>+3</sup>) and zinc in the hippocampus. *Brain Res*, 892(1), 51-62.
- Asad, S. F., Singh, S., Ahmad, A., & Hadi, S. M. (1999). Bilirubin-Cu(II) complex degrades DNA. *Biochim Biophys Acta*, 1428(2-3), 201-208.
- Asad, S. F., Singh, S., Ahmad, A., & Hadi, S. M. (2000). Inhibition of L-DOPA-Cu(II)-mediated DNA cleavage by bilirubin. *Toxicol In Vitro*, 14(5), 401-404.
- Asad, S. F., Singh, S., Ahmad, A., Khan, N. U., & Hadi, S. M. (2001). Prooxidant and antioxidant activities of bilirubin and its metabolic precursor biliverdin: a structure-activity study. *Chem Biol Interact*, 137(1), 59-74.
- Asad, S. F., Singh, S., Ahmad, A., & Hadi, S. M. (2002). Bilirubin/biliverdin-Cu(II) induced DNA breakage; reaction mechanism and biological significance. *Toxicol Lett*, 131(3), 181-189.
- Bakalbassis, E. G., Paschalidis, D. G., Raptopoulou, C. P., & Tangoulis, V. (1998). Synthesis, Crystal Structure, and Magnetic Properties of a One-Dimensional Polymeric Copper(II) Complex, Containing Unicoordinated Phthalato Bridges. *Inorg Chem*, 37(18), 4735-4737.
- Baranano, D. E., & Snyder, S. H. (2001). Neural roles for heme oxygenase: contrasts to nitric oxide synthase. *Proc Natl Acad Sci U S A*, 98(20), 10996-11002.
- Baranano, D. E., Rao, M., Ferris, C. D., & Snyder, S. H. (2002). Biliverdin reductase: a major physiologic cytoprotectant. *Proc Natl Acad Sci U S A*, 99(25), 16093-16098.

- Beckman, J. S., Carson, M., Smith, C. D., & Koppenol, W. H. (1993). ALS, SOD and peroxy nitrite. *Nature*, 364(6438), 584.
- Berg, J. M., & Shi, Y. (1996). The galvanization of biology: a growing appreciation for the roles of zinc. *Science*, 271(5252), 1081-1085.
- Bhat, R., & Hadi, S. M. (1994). DNA breakage by tannic acid and Cu(II): sequence specificity of the reaction and involvement of active oxygen species. *Mutat Res*, 313(1), 39-48.
- Brodersen, R. (1979). Bilirubin. Solubility and interaction with albumin and phospholipid. *J Biol Chem*, 254(7), 2364-2369.
- Burkitt, M. J., Milne, L., Nicotera, P., & Orrenius, S. (1996). 1,10-Phenanthroline stimulates internucleosomal DNA fragmentation in isolated rat-liver nuclei by promoting the redox activity of endogenous copper ions. *Biochem J*, 313 ( Pt 1), 163-169.
- Casanova, J., Alzuet, G., Latorre, J., & Borras, J. (1997). Spectroscopic, Magnetic, and Electrochemical Studies of a Dimeric N-Substituted-Sulfanilamide Copper(II) Complex. X-ray and Molecular Structure of the Cu(2)(sulfathiazolato)(4) Complex. *Inorg Chem*, 36(10), 2052-2058.
- Celec, P., & Prihodova, M. (2003). Pathogenesis of Alzheimer's disease and the role of heme oxygenase: new perspectives and hypothesis. *Folia Neuropathol*, 41(3), 155-160.
- Chen, Z., Zhang, T., Han, Y., & Zhu, L. (2006). Resonance light scattering spectroscopy study of interaction between norfloxacin and calf thymus DNA and its analytical application. *Spectrochim Acta A Mol Biomol Spectrosc*, 65(3-4), 919-924.
- Chowdhury JR, W. A., Chowdhary NR, Arias IM. (2001). *Hereditary jaundice and disorders of bilirubin metabolism*. Paper presented at the The metabolic basis of inherited disease.
- Conde-Martel, A., Gonzalez-Reimers, E., Santolaria-Fernandez, F., Castro-Aleman, V., Galindo-Martin, L., Rodriguez-Moreno, F., & Martinez-Riera, A. (1992). Combined effects of ethanol and protein deficiency on hepatic iron, zinc, manganese, and copper contents. *Alcohol*, 9(5), 341-348.
- Connor, J. R., Pavlick, G., Karli, D., Menzies, S. L., & Palmer, C. (1995). A histochemical study of iron-positive cells in the developing rat brain. *J Comp Neurol*, 355(1), 111-123.
- Cousins, R. J. (1985). Absorption, transport, and hepatic metabolism of copper and zinc: special reference to metallothionein and ceruloplasmin. *Physiol Rev*, 65(2), 238-309.

- Cowger, M. L., Igo, R. P., & Labbe, R. F. (1965). The mechanism of bilirubin toxicity studied with purified respiratory enzyme and tissue culture systems. *Biochemistry*, 4(12), 2763-2770.
- Deibel, M. A., Ehmann, W. D., & Markesberry, W. R. (1996). Copper, iron, and zinc imbalances in severely degenerated brain regions in Alzheimer's disease: possible relation to oxidative stress. *J Neurol Sci*, 143(1-2), 137-142.
- Dendrinou-Samara, C., Psomas, G., Raptopoulou, C. P., & Kessissoglou, D. P. (2001). Copper(II) complexes with phenoxyalkanoic acids and nitrogen donor heterocyclic ligands: structure and bioactivity. *J Inorg Biochem*, 83(1), 7-16.
- Dennerly, P. A., Seidman, D. S., & Stevenson, D. K. (2001). Neonatal hyperbilirubinemia. *N Engl J Med*, 344(8), 581-590.
- DiDonato, M., & Sarkar, B. (1997). Copper transport and its alterations in Menkes and Wilson diseases. *Biochim Biophys Acta*, 1360(1), 3-16.
- Dostert, P., Strolin Benedetti, M., & Dordain, G. (1988). Dopamine-derived alkaloids in alcoholism and in Parkinson's and Huntington's diseases. *J Neural Transm*, 74(2), 61-74.
- Drevensek, P., Leban, I., Turel, I., Giester, G. and E. Tillmanns. (2003). A novel copper(II) complex with 1,10-phenanthroline and ciprofloxacin. *Acta Crystallographica*, 59, m376-m378
- Efthimiadou, E. K., Sanakis, Y., Katsarou, M., Raptopoulou, C. P., Karaliota, A., Katsaros, N., & Psomas, G. (2006). Neutral and cationic mononuclear copper(II) complexes with enrofloxacin: structure and biological activity. *J Inorg Biochem*, 100(8), 1378-1388.
- Fazal F, Rahman A, Greensill J, Ainley K, Hadi SM, Parish JH. Strand scission in DNA by quercetin and Cu(II). Identification of free radical intermediates and biological consequence of scission. *Carcinogenesis* 1990 ;11 : 2005– 8.
- Feig, D. I., & Loeb, L. A. (1993). Mechanisms of mutation by oxidative DNA damage: reduced fidelity of mammalian DNA polymerase beta. *Biochemistry*, 32(16), 4466-4473.
- Ferri, A., Duffard, R., Sturtz, N., & Evangelista de Duffard, A. M. (2003). Iron, zinc and copper levels in brain, serum and liver of neonates exposed to 2,4-dichlorophenoxyacetic acid. *Neurotoxicol Teratol*, 25(5), 607-613.
- Fraga, C. G. (2005). Relevance, essentiality and toxicity of trace elements in human health. *Mol Aspects Med*, 26(4-5), 235-244.

- Gagne, R. R., Allison, J. L., Gall, R. S., & Koval, C. A. (1977). Models for copper-containing proteins: structure and properties of novel five-coordinate copper(I) complexes. *J Am Chem Soc*, 99(22), 7170-7178.
- Glaser, K. L. (2006). *Jaundice and Hyperbilirubinemia in the Newborn* from <http://www.medstudents.com.br/pedia/pedia3.htm>
- Gordon, N. (2003). Iron deficiency and the intellect. *Brain Dev*, 25(1), 3-8.
- Gourley, G. R. (1997). Bilirubin metabolism and kernicterus. *Adv Pediatr*, 44, 173-229.
- Grojean, S., Koziel, V., Vert, P., & Daval, J. L. (2000). Bilirubin induces apoptosis via activation of NMDA receptors in developing rat brain neurons. *Exp Neurol*, 166(2), 334-341.
- Grojean, S., Lievre, V., Koziel, V., Vert, P., & Daval, J. L. (2001). Bilirubin exerts additional toxic effects in hypoxic cultured neurons from the developing rat brain by the recruitment of glutamate neurotoxicity. *Pediatr Res*, 49(4), 507-513.
- Hahm, J. S., Ostrow, J. D., Mukerjee, P., & Celic, L. (1992). Ionization and self-association of unconjugated bilirubin, determined by rapid solvent partition from chloroform, with further studies of bilirubin solubility. *J Lipid Res*, 33(8), 1123-1137.
- Halliwell, B., & Gutteridge, J. M. (1990). The antioxidants of human extracellular fluids. *Arch Biochem Biophys*, 280(1), 1-8.
- Halliwell, B., Gutteridge, J.M.C. (1999). *Free Radicals in Biology and Medicine* (4 th ed.): Oxford University Press,Oxford.
- Hanko, E., Hansen, T. W., Almaas, R., Lindstad, J., & Rootwelt, T. (2005). Bilirubin induces apoptosis and necrosis in human NT2-N neurons. *Pediatr Res*, 57(2), 179-184.
- Hansen, T. W., Bratlid, D., & Walaas, S. I. (1988). Bilirubin decreases phosphorylation of synapsin I, a synaptic vesicle-associated neuronal phosphoprotein, in intact synaptosomes from rat cerebral cortex. *Pediatr Res*, 23(2), 219-223.
- Hansen, T. W. (2000). Kernicterus in term and near-term infants--the specter walks again. *Acta Paediatr*, 89(10), 1155-1157.
- Harish Prashanth, K. V., Dharmesh, S. M., Jagannatha Rao, K. S., & Tharanathan, R. N. (2007). Free radical-induced chitosan depolymerized products protect calf thymus DNA from oxidative damage. *Carbohydr Res*, 342(2), 190-195.

- Hartard, C., Weisner, B., Dieu, C., & Kunze, K. (1993). Wilson's disease with cerebral manifestation: monitoring therapy by CSF copper concentration. *J Neurol*, 241(2), 101-107.
- Howell, G. A., Welch, M. G., & Frederickson, C. J. (1984). Stimulation-induced uptake and release of zinc in hippocampal slices. *Nature*, 308(5961), 736-738.
- Imlay, J. A., & Linn, S. (1988). DNA damage and oxygen radical toxicity. *Science*, 240(4857), 1302-1309.
- Iztok Turel, K. G., Ivan Leban and Nata Bukovec. (1996). Synthesis, crystal structure, and characterization of three novel compounds of the quinolone family member *Journal of Inorganic Biochemistry* 61(3), 197-212.
- Kapitulnik, J. (2004). Bilirubin: an endogenous product of heme degradation with both cytotoxic and cytoprotective properties. *Mol Pharmacol*, 66(4), 773-779.
- Karp, W. B. (1979). Biochemical alterations in neonatal hyperbilirubinemia and bilirubin encephalopathy: a review. *Pediatrics*, 64(3), 361-368.
- Kishi, R., Ikeda, T., Miyake, H., Uchino, E., Tsuzuki, T., & Inoue, K. (1982). Regional distribution of lead, zinc, iron and copper in suckling and adult rat brains. *Brain Res*, 251(1), 180-182.
- Labieniec, M., & Gabryelak, T. (2006). Interactions of tannic acid and its derivatives (ellagic and gallic acid) with calf thymus DNA and bovine serum albumin using spectroscopic method. *J Photochem Photobiol B*, 82(1), 72-78.
- Lai, J. C., & Blass, J. P. (1984). Neurotoxic effects of copper: inhibition of glycolysis and glycolytic enzymes. *Neurochem Res*, 9(12), 1699-1710.
- Leist, M., & Nicotera, P. (1998). Apoptosis, excitotoxicity, and neuropathology. *Exp Cell Res*, 239(2), 183-201.
- Li, D. (1998). Effects of iron deficiency on iron distribution and gamma-aminobutyric acid (GABA) metabolism in young rat brain tissues. *Hokkaido Igaku Zasshi*, 73(3), 215-225.
- Liu, B., Gao, H. M., Wang, J. Y., Jeohn, G. H., Cooper, C. L., & Hong, J. S. (2002). Role of nitric oxide in inflammation-mediated neurodegeneration. *Ann N Y Acad Sci*, 962, 318-331.
- Lubec G. The hydroxyl radical: from chemistry to human disease. *J Investig Med* 1996 ; 44 ::324-46. Review.
- Macias, B., Villa, M. V., Rubio, I., Castineiras, A., & Borras, J. (2001). Complexes of Ni(II) and Cu(II) with ofloxacin. Crystal structure of a new Cu(II) ofloxacin complex. *J Inorg Biochem*, 84(3-4), 163-170.

- Macyk W, F. A., Stochel G, . (2005). Metal compounds and small molecules activation – case studies. *Coordination Chemistry Reviews*, 249, 2437–2457.
- Madhavan, M., Wattigney, W. A., Srinivasan, S. R., & Berenson, G. S. (1997). Serum bilirubin distribution and its relation to cardiovascular risk in children and young adults. *Atherosclerosis*, 131(1), 107-113.
- McDonald, J. W., Shapiro, S. M., Silverstein, F. S., & Johnston, M. V. (1998). Role of glutamate receptor-mediated excitotoxicity in bilirubin-induced brain injury in the Gunn rat model. *Exp Neurol*, 150(1), 21-29.
- McGeary, R. P., Szyczew, A. J., & Toth, I. (2003). Biological properties and therapeutic potential of bilirubin. *Mini Rev Med Chem*, 3(3), 253-256.
- Mediavilla, M. G., Pascolo, L., Rodriguez, J. V., Guibert, E. E., Ostrow, J. D., & Tiribelli, C. (1999). Uptake of [(3)H]bilirubin in freshly isolated rat hepatocytes: role of free bilirubin concentration. *FEBS Lett*, 463(1-2), 143-145.
- Meuwissen, J. A. T. P. H., & Heirwegh, K. P. M. (1982). *Bilirubin* (Vol. 2). Florida: CRC, Boca Raton.
- Mustafa, M. G., Cowger, M. L., & King, T. E. (1969). Effects of bilirubin on mitochondrial reactions. *J Biol Chem*, 244(23), 6403-6414.
- Neu, H. C. (1987). Ciprofloxacin: an overview and prospective appraisal. *Am J Med*, 82(4A), 395-404.
- Ostrow, J. D., Pascolo, L., Shapiro, S. M., & Tiribelli, C. (2003a). New concepts in bilirubin encephalopathy. *Eur J Clin Invest*, 33(11), 988-997.
- Ostrow, J. D., Pascolo, L., & Tiribelli, C. (2003b). Reassessment of the unbound concentrations of unconjugated bilirubin in relation to neurotoxicity in vitro. *Pediatr Res*, 54(1), 98-104.
- Park, W. S., Chang, Y. S., & Lee, M. (2002). Effect of 7-nitroindazole on bilirubin-induced changes in brain cell membrane function and energy metabolism in newborn piglets. *Biol Neonate*, 82(1), 61-65.
- Pascolo, L., Fornetti, C., Garcia-Mediavilla, M. V., Ostrow, J. D., & Tiribelli, C. (2001). Mechanisms for the transport of unconjugated bilirubin in human trophoblastic BeWo cells. *FEBS Lett*, 495(1-2), 94-99.
- Ponting, C. P. (2001). Domain homologues of dopamine beta-hydroxylase and ferric reductase: roles for iron metabolism in neurodegenerative disorders? *Hum Mol Genet*, 10(17), 1853-1858.

- Porter, M. L., & Dennis, B. L. (2002). Hyperbilirubinemia in the term newborn. *Am Fam Physician*, 65(4), 599-606.
- Psomas, G., Raptopoulou, C. P., Iordanidis, L., Dendrinou-Samara, C., Tangoulis, V., & Kessissoglou, D. P. (2000). Structurally diverse copper(II)-carboxylato complexes: neutral and ionic mononuclear structures and a novel binuclear structure. *Inorg Chem*, 39(14), 3042-3048.
- Purnell, L. G., Shepherd, J. C., & Hodgson, D. J. (1975). Interaction of metal ions with 8-azapurines. Synthesis and structure of tetrachlorobis-2-((5-amino-4-carboxamidinium)(1,2,3)triazole)copper(II) monohydrate. *J Am Chem Soc*, 97(9), 2376-2380.
- Quinlan, G. J., & Gutteridge, J. M. (1987). Oxygen radical damage to DNA by rifamycin SV and copper ions. *Biochem Pharmacol*, 36(21), 3629-3633.
- Rahman, A., Shahabuddin, Hadi, S. M., Parish, J. H., & Ainley, K. (1989). Strand scission in DNA induced by quercetin and Cu(II): role of Cu(I) and oxygen free radicals. *Carcinogenesis*, 10(10), 1833-1839.
- Reisman, Y., Gips, C. H., Lavelle, S. M., & Wilson, J. H. (1996). Clinical presentation of (subclinical) jaundice--the Euricterus project in The Netherlands. United Dutch Hospitals and Euricterus Project Management Group. *Hepatogastroenterology*, 43(11), 1190-1195.
- Rodrigues, C. M., Sola, S., & Brites, D. (2002a). Bilirubin induces apoptosis via the mitochondrial pathway in developing rat brain neurons. *Hepatology*, 35(5), 1186-1195.
- Rodrigues, C. M., Sola, S., Brito, M. A., Brites, D., & Moura, J. J. (2002b). Bilirubin directly disrupts membrane lipid polarity and fluidity, protein order, and redox status in rat mitochondria. *J Hepatol*, 36(3), 335-341.
- Rodrigues, C. M., Sola, S., Silva, R. F., & Brites, D. (2002c). Aging confers different sensitivity to the neurotoxic properties of unconjugated bilirubin. *Pediatr Res*, 51(1), 112-118.
- Rojkind, M., & Dunn, M. A. (1979). Hepatic fibrosis. *Gastroenterology*, 76(4), 849-863.
- Ruiz, M., Perello, L., Ortiz, R., Castineiras, A., Maichle-Mossmer, C., & Canton, E. (1995). Synthesis, characterization, and crystal structure of [Cu(cinoxacinate)2].2H<sub>2</sub>O complex: a square-planar CuO<sub>4</sub> chromophore. Antibacterial studies. *J Inorg Biochem*, 59(4), 801-810.

- Saha, D. K., Sandbhor, U., Shirisha, K., Padhye, S., Deobagkar, D., Anson, C. E., & Powell, A. K. (2004). A novel mixed-ligand antimycobacterial dimeric copper complex of ciprofloxacin and phenanthroline. *Bioorg Med Chem Lett*, 14(12), 3027-3032.
- Seubert, J. M., Darmon, A. J., El-Kadi, A. O., D'Souza, S. J., & Bend, J. R. (2002). Apoptosis in murine hepatoma hepa 1c1c7 wild-type, C12, and C4 cells mediated by bilirubin. *Mol Pharmacol*, 62(2), 257-264.
- Shamsi, F. A., & Hadi, S. M. (1995). Photoinduction of strand scission in DNA by uric acid and Cu(II). *Free Radic Biol Med*, 19(2), 189-196.
- Shapiro, S. M. (2003). Bilirubin toxicity in the developing nervous system. *Pediatr Neurol*, 29(5), 410-421.
- Sheline, C. T., Behrens, M. M., & Choi, D. W. (2000). Zinc-induced cortical neuronal death: contribution of energy failure attributable to loss of NAD(+) and inhibition of glycolysis. *J Neurosci*, 20(9), 3139-3146.
- Schenker, S., Hoyumpa, AM., & McCandless D.W. (1986). Bile pigments and jaundice. In J. D. Ostrow (Ed.), *Bile pigments and jaundice* (Vol. 2). New York: Marcel Dekker.
- Schneider, W. C. (1957). Determination of nucleic acids in tissues by pentose analysis *Methods of Enzymology*, 3, 680 - 684
- Silva, R., Mata, L. R., Gulbenkian, S., Brito, M. A., Tiribelli, C., & Brites, D. (1999). Inhibition of glutamate uptake by unconjugated bilirubin in cultured cortical rat astrocytes: role of concentration and pH. *Biochem Biophys Res Commun*, 265(1), 67-72.
- Silva, R. F., Rodrigues, C. M., & Brites, D. (2001). Bilirubin-induced apoptosis in cultured rat neural cells is aggravated by chenodeoxycholic acid but prevented by ursodeoxycholic acid. *J Hepatol*, 34(3), 402-408.
- Silva, R. F., Rodrigues, C. M., & Brites, D. (2002). Rat cultured neuronal and glial cells respond differently to toxicity of unconjugated bilirubin. *Pediatr Res*, 51(4), 535-541.
- Sorenson, J. R. (1989). Copper complexes offer a physiological approach to treatment of chronic diseases. *Prog Med Chem*, 26, 437-568.
- Stanley, F. L., Doumas, BT & Ashwood, ER (2004). General Clinical Performance of Bilirubin Determinations in US Laboratories—Revisited. *Clin Chem*, 50(1), 190-194.

- Strausak, D., Mercer, J. F., Dieter, H. H., Stremmel, W., & Multhaup, G. (2001). Copper in disorders with neurological symptoms: Alzheimer's, Menkes, and Wilson diseases. *Brain Res Bull*, 55(2), 175-185.
- Tangoulis, V., Malamatari, D. A., Soulti, K., Stergiou, V., Raptopoulou, C. P., Terzis, A., Kabanos, T. A., & Kessissoglou, D. P. (1996). Manganese(II/II/II) and Manganese(III/II/III) Trinuclear Compounds. Structure and Solid and Solution Behavior. *Inorg Chem*, 35(17), 4974-4983.
- Van Houten, B., Woshner, V., & Santos, J. H. (2006). Role of mitochondrial DNA in toxic responses to oxidative stress. *DNA Repair (Amst)*, 5(2), 145-152.
- Wallis, S. C., Gahan, L. R., Charles, B. G., Hambley, T. W., & Duckworth, P. A. (1996). Copper(II) complexes of the fluoroquinolone antimicrobial ciprofloxacin. Synthesis, X-ray structural characterization, and potentiometric study. *J Inorg Biochem*, 62(1), 1-16.
- Walravens, P. A. (1980). Nutritional importance of copper and zinc in neonates and infants. *Clin Chem*, 26(2), 185-189.
- Wang X, C. J., Chowdhury NR, . (2006). Bilirubin metabolism: Applied physiology. *Current Pediatrics*(16), 70-74.
- Wiegand, R. C., Godson, G. N., & Radding, C. M. (1975). Specificity of the S1 nuclease from Aspergillus oryzae. *J Biol Chem*, 250(22), 8848-8855.
- Wennberg, R. P. (2000). The blood-brain barrier and bilirubin encephalopathy. *Cell Mol Neurobiol*, 20(1), 97-109.
- Wood, S. G., Gedik, C. M., & Collins, A. R. (2000). Controlled oxidation of calf thymus DNA to produce standard samples for 8-oxodeoxyguanosine analysis; effects of freeze-drying, storage and hydrolysis conditions. *Free Radic Res*, 32(4), 327-332.
- Wu, L. J., Leenders, A. G., Cooperman, S., Meyron-Holtz, E., Smith, S., Land, W., Tsai, R. Y., Berger, U. V., Sheng, Z. H., & Rouault, T. A. (2004). Expression of the iron transporter ferroportin in synaptic vesicles and the blood-brain barrier. *Brain Res*, 1001(1-2), 108-117.
- Wu, T. W., Fung, K. P., & Yang, C. C. (1994). Unconjugated bilirubin inhibits the oxidation of human low density lipoprotein better than Trolox. *Life Sci*, 54(25), P477-481.
- Yamada, N., Sawasaki, Y., & Nakajima, H. (1977). Impairment of DNA synthesis in Gunn rat cerebellum. *Brain Res*, 126(2), 295-307.
- Youdim, M. B. (2000). Nutrient deprivation and brain function: iron. *Nutrition*, 16(7-8), 504-508.

Zong, W. X., & Thompson, C. B. (2006). Necrotic death as a cell fate. *Genes Dev*, 20(1), 1-15.

Zucker, S. D., Goessling, W., & Hoppin, A. G. (1999). Unconjugated bilirubin exhibits spontaneous diffusion through model lipid bilayers and native hepatocyte membranes. *J Biol Chem*, 274(16), 10852-10862.

Zecca, L., Stroppolo, A., Gatti, A., Tampellini, D., Toscani, M., Gallorini, M., Giaveri, G., Arosio, P., Santambrogio, P., Fariello, R. G., Karatekin, E., Kleinman, M. H., Turro, N., Hornykiewicz, O., & Zucca, F. A. (2004). The role of iron and copper molecules in the neuronal vulnerability of locus caeruleus and substantia nigra during aging. *Proc Natl Acad Sci U S A*, 101(26), 9843-9848.

จิฬิสิทธิ์มหาวิทยาลัยเชียงใหม่  
Copyright<sup>©</sup> by Chiang Mai University  
All rights reserved