

## TABLE OF CONTENTS

	Pages
ACKNOWLEDGEMENTS	iii
ABSTRACT (ENGLISH)	iv
ABSTRACT (THAI)	vi
LIST OF TABLES	ix
LIST OF FIGURES	xi
LIST OF SCHEMES	xiv
ABBREVIATIONS AND SYMBOLS	xv
CHAPTER I : INTRODUCTION	
- Review of the Genus <i>Kaempferia</i>	1
- Historical background of <i>K. parviflora</i>	3
CHAPTER II : REVIEW OF LITERATURES	
- General knowledge	8
- Flavonoid biosynthesis	9
- Flavonoid antioxidants: Chemistry, Structure-activity relationship and mechanism	14
- The chemistry and biological activities of <i>K. parviflora</i> Wall. ex Bak.	22
CHAPTER III : MATERIALS AND METHODS	
- Source and authentication of the plant materials	28
- General techniques	28
- Isolation the <i>n</i> -hexane extract of <i>K. parviflora</i>	34
- Isolation the ethyl acetate extract of <i>K. parviflora</i>	40

- Isolation the methanol extract of <i>K. parviflora</i>	44
CHAPTER IV : RESULTS AND DISCUSSION	
- Structure elucidation of the isolated compounds	49
- Antioxidant activity of crude extracts and isolated compounds of <i>K.parviflora</i>	62
CHAPTER V : CONCLUSIONS	77
REFERENCES	80
APPENDIX	90
VITA	116

## LIST OF TABLES

Tables	Page
1.1 The occurrence of <i>Kaempferia</i> in Thailand	1
2.1 List of enzymes leading to various flavonoid classes	12
2.2 Chemical constituents and Biological activities of <i>K. parviflora</i>	22
3.1 Antioxidant activity of fraction KP3-001 to KP3-005 by ABTS <sup>o+</sup> assay	45
3.2 Antioxidant activity of fraction KP3-005-1 to KP3-005-7 by ABTS <sup>o+</sup> assay	47
4.1 The amount and percentage yield of extracts from <i>K. parviflora</i>	48
4.2 The antioxidant activity of crude extracts of <i>K. parviflora</i>	49
4.3 NMR data of compound 5-hydroxy-3,7-dimethoxyflavone, 5-hydroxy-7,4'-dimethoxyflavone and 5-hydroxy-7-methoxyflavone	57
4.4 NMR data of compound 3,5,7-trimethoxyflavone and 5,7-dimethoxyflavone	58
4.5 The physical properties of crude extracts and isolated compounds from <i>n</i> -hexane, ethyl acetate and methanol fractions from rhizomes of <i>K. parviflora</i>	61
4.6 The data used to construct calibration curve for ascorbic acid standard	62
4.7 The data used to construct calibration curve for quercetin standard	63
4.8 The data used to construct calibration curve for pyrogallol standard	63

- 4.9 The antioxidant activity by ABTS<sup>0+</sup> assay of isolated compounds from *n*-hexane, ethyl acetate and methanol extracts of *K. parviflora* as a function of concentration of standard expressed as mg/g of ascorbic acid (a), quercetin (b) and pyrogallol (c) solutions 65
- 4.10 The data used to construct calibration curve for Fe<sub>2</sub>SO<sub>4</sub> standard 66
- 4.11 The antioxidant activity by ABTS<sup>0+</sup> assay and FRAP assay of isolated compounds from *n*-hexane, ethyl acetate and methanol extracts of *K. parviflora*. 68

## LIST OF FIGURES

Figure	Page
1.1 Leaves and flowers of <i>K. parviflora</i>	4
1.2 Rhizomes of <i>K. parviflora</i>	5
2.1 The flavonoid skeleton and its numbering system	10
2.2 Flavonoid biosynthesis pathway	13
2.3 Chain reaction of lipidperoxidation	21
4.1 The flavone skeleton and its numbering system	50
4.2 The chemical structure of compound KP1-002P, KP1-004-5P and KP1-010P	51
4.3 The chemical structure of compound KP1-005-2P and KP1-007-6P	52
4.4 The chemical structure of compound KP1-006-5P and KP2-007-2-02P	53
4.5 The chemical structure of compound KP1-012P and KP2-007-1P	54
4.6 The chemical structure of compound KP2-005P and KP2-006-2P	55
4.7 Isolated compounds from <i>n</i> -hexane and ethyl acetate fractions (5 compounds)	56
4.8 The LC chromatogram of KP3-005-2P from methanol extract of <i>K. parviflora</i>	59
4.9 The mass spectrm of KP3-005-2P from methanol extract of <i>K. parviflora</i>	60
4.10 Concentration-response curves for the absorbance at 743 nm for ABTS <sup>0+</sup> as a function of concentration of standard ascorbic acid (a), quercetin (b) and pyrogallol (c) solutions	64

4.11	Calibration curve for the absorbance at 593 nm for FRAP assay as a function of concentration of standard $\text{Fe}_2\text{SO}_4$	67
4.12	Quercetin	69
4.13	Summary of isolated compounds obtained from <i>n</i> -hexane and ethyl acetate fractions	71
4.14	Structure of flavonoids which reduce metal ion	73
A-1	The IR spectrum of compound KP1-002P, KP1-004-5P and KP1-010P	91
A-2	The $^1\text{H}$ -NMR spectrum of compound KP1-002P, KP1-004-5P and KP1-010P (in $\text{CDCl}_3$ )	92
A-3	The $^1\text{H}$ -NMR expanded spectrum of compound KP1-002P, KP1-004-5P and KP1-010P (in $\text{CDCl}_3$ )	93
A-4	The $^{13}\text{C}$ -NMR spectrum of compound KP1-002P, KP1-004-5P and KP1-010P (in $\text{CDCl}_3$ )	94
A-5	The mass spectrum of compound KP1-002P, KP1-004-5P and KP1-010P	95
B-1	The IR spectrum of compound KP1-005-2P and KP1-007-6P	96
B-2	The $^1\text{H}$ -NMR spectrum of compound KP1-005-2P and KP1-007-6P (in $\text{CDCl}_3$ )	97
B-3	The $^1\text{H}$ -NMR expanded spectrum of compound KP1-005-2P and KP1-007-6P (in $\text{CDCl}_3$ )	98
B-4	The $^1\text{H}$ -NMR expanded spectrum of compound KP1-005-2P and KP1-007-6P (in $\text{CDCl}_3$ )	99
B-5	The $^{13}\text{C}$ -NMR spectrum of compound KP1-005-2P and KP1-007-6P (in $\text{CDCl}_3$ )	100
B-6	The mass spectrum of compound KP1-005-2P and KP1-007-6P	101
C-1	The IR spectrum of compound KP1-006-5P and KP2-007-2-02P	102
C-2	The $^1\text{H}$ -NMR spectrum of compound KP1-006-5P and KP2-007-2-02P (in $\text{CDCl}_3$ )	103

C-3	The $^1\text{H}$ -NMR expanded spectrum of compound KP1-006-5P and KP2-007-2-02P (in $\text{CDCl}_3$ )	104
C-4	The $^{13}\text{C}$ -NMR spectrum of compound KP1-006-5P and KP2-007-2-02P (in $\text{CDCl}_3$ )	105
C-5	The mass spectrum of compound KP1-006-5P and KP2-007-2-02P	106
D-1	The IR spectrum of compound KP1-012P and KP2-007-1P	107
D-2	The $^1\text{H}$ -NMR spectrum of compound KP1-012P and KP2-007-1P (in $\text{CDCl}_3$ )	108
D-3	The $^{13}\text{C}$ -NMR spectrum of compound KP1-012P and KP2-007-1P (in $\text{CDCl}_3$ )	109
D-4	The mass spectrum of compound KP1-012P and KP2-007-1P	110
E-1	The IR spectrum of compound KP2-005P and KP2-006-2P	111
E-2	The $^1\text{H}$ -NMR spectrum of compound KP2-005P and KP2-006-2P (in $\text{CDCl}_3$ )	112
E-3	The $^1\text{H}$ -NMR expanded spectrum of compound KP2-005P and KP2-006-2P (in $\text{CDCl}_3$ )	113
E-4	The $^{13}\text{C}$ -NMR spectrum of compound KP2-005P and KP2-006-2P (in $\text{CDCl}_3$ )	114
E-5	The mass spectrum of compound KP2-005P and KP2-006-2P	115

## LIST OF SCHEMES

Schemes	Page
3.1 Isolation scheme of Rhizomes of <i>K. parviflora</i>	33
3.2 Isolation scheme of <i>n</i> -hexane extract of <i>K. parviflora</i> (KP1)	34
3.3 Isolation scheme of compound KP1-004-5P from <i>n</i> -hexane extract	35
3.4 Isolation scheme of compound KP1-005-2P from <i>n</i> -hexane extract	36
3.5 Isolation scheme of compound KP1-006-5P from <i>n</i> -hexane extract	37
3.6 Isolation scheme of compound KP1-007-6P from <i>n</i> -hexane extract	38
3.7 Isolation scheme of Ethyl acetate extract of <i>K. parviflora</i> (KP2)	40
3.8 Isolation scheme of compound KP2-006-2P from Ethyl acetate extract	41
3.9 Isolation scheme of compound KP2-007-1P from Ethyl acetate extract	42
3.10 Isolation scheme of compound KP2-007-2-02P from Ethyl acetate extract	43
3.11 Isolation scheme of Methanol extract of <i>K. parviflora</i> (KP3)	44
3.12 Isolation scheme of compound KP3-005 from Methanol extract	46



# ABBREVIATIONS AND SYMBOLS

ABTS <sup>0+</sup>	Radical form of 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid)
C	Carbon
CC	Column chromatography
°C	Degree Celsius
cm	Centimeter
CDCl <sub>3</sub>	Deuteriochloroform
<i>d</i>	doublet
<i>dd</i>	doublet of doublet
DEPT	Distortionless enhancement by polarization transfer
EIMS	electron impact mass spectra
EtOAc	ethyl acetate
EtOH	Ethanol
FRAP	Ferric reducing ability power
G	Gram
<sup>1</sup> H-NMR	Proton nuclear magnetic resonance
HMBC	<sup>1</sup> H-detected heteronuclear multiple bond coherence
HSQC	<sup>1</sup> H-detected high sensitive quantum coherence
Hz	Hertz
i.d.	Internal diameter
IR	Infrared spectroscopy
<i>J</i>	Coupling constant
LC-MS	liquid chromatography mass spectra
<i>m</i>	Multiplet
MeOH	Methanol
MHz	Megahertz
mM	Milimolar
m.p.	Melting point

$m/z$	A value of mass divided by charge
NMR	Nuclear magnetic resonance spectroscopy
2D-NMR	Two dimensional nuclear magnetic resonance
$R^2$	Coefficient of determination
$s$	Singlet
TLC	Thin layer chromatography
TMS	Tetramethylsilane
UV	Ultraviolet
$\delta$	Chemical shift relative to TMS

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่  
Copyright© by Chiang Mai University  
All rights reserved